



CROSS REFERENCE TO RELATED APPLICATIONS

TECHNICAL FIELD OF THE INVENTION

BACKGROUND OF THE INVENTION

An important and significant goal in healthcare is to discover and make available safer and more effective drugs for the treatment of cancer. Most chemotherapeutic agents act by disrupting DNA metabolism, DNA synthesis, DNA transcription, or microtubule spindle function, or by perturbing chromosomal structural integrity by introducing DNA lesions. These processes affect both normal and tumor cells. The maintenance of DNA integrity is essential to cell viability in normal cells, therefore, anticancer drugs have the lowest therapeutic index of any drug class.

An individual cell creates an exact copy of its chromosomes, and then segregates each copy into two cells by a process called mitosis. The mitotic cycle can be divided into three major events: DNA replication, chromosome segregation, and cell division. Cells have sensing mechanisms to maintain the order of these steps with respect to one another and to ensure that each step is executed with high fidelity. The sensing mechanisms for these processes are referred to as "checkpoints" in L.H. Hartwell et al., *Science*, Nov. 3, 1989, 246(4930):629-34.

Cell cycle checkpoints have been reported to comprise at least three distinct classes of polypeptides. Each class of polypeptides acts sequentially in response to cell cycle signals or defects in chromosomal mechanisms (Carr, (1996) *Science*, 271:314-315). One family of proteins detects or senses DNA damage or abnormalities in the cell cycle. These sensors include Ataxia-Telangiectasia Mutated (Atm) and Ataxia-Telangiectasia Rad-related (Atr) (Keegan et al., (1996) *Genes Dev.*, 10:2423-2437). Another class of polypeptides amplify and transmit the signal detected by the detector and is exemplified by Rad53 (Allen et al. (1994) *Genes Dev.*, 8:2416-2488) and Chk1. In addition, cell cycle effectors, such as p53, mediate a cellular response, including, for example, arrest of mitosis and/or meiosis and apoptosis.

DNA damage can be induced by drugs, radiation, or can be spontaneously generated during the course of normal metabolism. DNA damage checkpoints ensure that cells with unrepaired DNA lesions do not progress into the DNA synthesis phase or mitosis until chromosomal lesions have been removed. Cell

5 This mechanism of regulation of Cdc2 activity stim-
 ulates cell cycle arrest to prevent cells from
 entering mitosis in the presence of DNA damage or
 unreplicated DNA.

10

15

20



(I)

35

- 8 -

X¹ is null, -O-, -S-, -CH₂-, or -N(R¹)-;

X² is -O-, -S-, or -N(R¹)-;

Y is O or S; or =Y represents two hydrogen atoms attached to a common carbon atom;

5 W is selected from the group consisting of heteroaryl, aryl, heterocycloalkyl, cycloalkyl, and C₁₋₃ alkyl substituted with a heteroaryl or aryl group; and

10 Z is selected from the group consisting of hydro, aryl, and heteroaryl;

wherein said aryl groups of W and Z are optionally substituted with one to four substituents represented by R², said heteroaryl groups of W and Z are optionally substituted with one to four substituents represented by R⁵, and said heterocycloalkyl and cycloalkyl groups of W are optionally substituted with one to two substituents represented by R⁶;

R¹ is selected from the group consisting of hydrogen, C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, and aryl;

20 R² is selected from the group consisting of halogen, C₁₋₆alkyl, C₂₋₆alkenyl, OCF₃, NO₂, CN, NC, N(R³)₂, OR³, CO₂R³, C(O)N(R³)₂, C(O)R³, N(R¹)COR³, N(R¹)C(O)OR³, N(R³)C(O)OR³, N(R³)C(O)-C₁₋₃alkylene-C(O)R³, N(R³)C(O)-C₁₋₃alkylene-C(O)OR³, N(R³)C(O)-C₁₋₃alkylene-OR³, N(R³)C(O)-C₁₋₃alkylene-NHC(O)OR³, N(R³)C(O)-C₁₋₃alkylene-SO₂NR³, C₁₋₃alkylene-OR³, and SR³;


25 R³ is selected from the group consisting of hydro, C₁₋₆alkyl, C₂₋₆alkenyl, cycloalkyl, aryl, heteroaryl, SO₂R⁴, C₁₋₆alkyl substituted with one or more of halo, hydroxy, aryl, heteroaryl, heterocycloalkyl, N(R⁴)₂, and SO₂R⁴, C₁₋₃alkylenearyl, C₁₋₃alkyleneheteroaryl, C₁₋₃alkyleneC₃₋₈heterocycloalkyl, C₁₋₃alkyleneSO₂aryl, optionally substituted C₁₋₃alkylene-N(R⁴)₂, OCF₃, C₁₋₃alkyleneN(R⁴)₃⁺, C₃₋₈heterocycloalkyl, and CH(C₁₋₃alkyleneN(R⁴)₂)₂, or two R³ groups are taken

- 9 -

together to form an optionally substituted 3- to 6-membered aliphatic ring;

R^4 is selected from the group consisting of hydro, C_{1-6} alkyl, cycloalkyl, aryl, heteroaryl, C_{1-3} -alkylenearyl, and SO_2C_{1-6} alkyl, or two R^4 groups are taken together to form an optionally substituted 3- to 6-membered ring;


R^5 is selected from the group consisting of C_{1-6} alkyl, aryl, $N(R^3)_2$, OR^3 , halo, N_3 , CN, C_{1-3} alkylenearyl, C_{1-3} alkylene $N(R^3)_2$, $C(O)R^3$, and



R^6 is selected from the group consisting of halo and C_{1-6} alkyl,

and pharmaceutically acceptable salts or solvates thereof.

In another aspect, the present invention is directed to aryl- and heteroaryl-disubstituted urea compounds having a structural formula (II)



(II)


- 10 -

wherein Y' is O or S;


W' is selected from the group consisting

of


5




10



15




20



25


30

- 11 -



5

, and




10

optionally substituted with from one to four substituents selected from the group consisting of C_{1-6} alkyl, aryl, $N(R^7)_2$, OR^7 , N_3 , CN , $C(O)R^7$, C_{1-3} alkylenearyl, C_{1-3} alkylene $N(R^{12})_2$,

15


20




25

Z' is selected from the group consisting of:

30



and



35

- 12 -

wherein:

Q' is selected from the group consisting of hydrogen, OR⁷, SR⁷, and N(R⁷)₂;

5 J' is selected from the group consisting of C-R⁸, N-R⁸, O, and S;

K' is selected from the group consisting of C-R⁹, N-R⁹, O, and S;

L' is selected from the group consisting of C-R¹⁰, N-R¹⁰, O, and S;

10 M' is selected from the group consisting of C-R¹¹, N-R¹¹, O, and S;

wherein:

R⁷, independently, is selected from the group consisting of hydro, C₁₋₆alkyl, C₂₋₆alkenyl, 15 cycloalkyl, aryl, heteroaryl, SO₂R¹², C₁₋₆alkyl substituted with one or more of halo, hydroxy, aryl, heteroaryl, heterocycloalkyl, N(R¹²)₂, and SO₂R¹², C₁₋₃alkylenearyl, C₁₋₃alkyleneheteroaryl, C₁₋₃alkylene-C₃₋₈heterocycloalkyl, C₁₋₃alkyleneSO₂aryl, optionally 20 substituted C₁₋₃alkyleneN(R¹²)₂, OCF₃, C₁₋₃alkylene-N(R¹²)₃⁺, C₃₋₈heterocycloalkyl, and CH(C₁₋₃alkylene-N(R¹²)₂)₂, or two R⁷ groups are taken together to form an optionally substituted 3- to 6-membered aliphatic ring;

25 R⁸, R⁹, and R¹⁰ are each independently selected from the group consisting of null, hydro, halo, optionally substituted C₁₋₆alkyl, C₂₋₆alkenyl, OCF₃, NO₂, CN, NC, N(R⁷)₂, OR⁷, CO₂R⁷, C(O)N(R⁷)₂, C(O)R⁷, N(R¹³)COR⁷, N(R¹³)C(O)OR⁷, N(R⁷)C(O)OR⁷, 30 N(R⁷)C(O)C₁₋₃alkyleneC(O)R⁷, N(R⁷)C(O)C₁₋₃alkylene-C(O)OR⁷, N(R⁷)C(O)C₁₋₃alkyleneOR⁷, N(R⁷)C(O)C₁₋₃alkyleneNHC(O)OR⁷, N(R⁷)C(O)C₁₋₃alkyleneSO₂NR⁷, CF₃, C₁₋₃alkyleneN(R¹²)SO₂aryl, C₁₋₃alkyleneN(R¹²)SO₂heteroaryl, C₁₋₃alkyleneOC₁₋₃alkylenearyl, C₁₋₃alkyleneN(R¹²)C₁₋₃alkylenearyl, 35 C₁₋₃alkyleneN(R¹²)C₁₋₃alkyleneheteroaryl,

- 13 -

C_{1-3} alkyleneN(R¹²)C(O)R⁷, C_{1-3} alkyleneN(R¹²)C(O) C_{1-3} -alkyleneOR², C_{1-3} alkyleneN(R¹²)C(O)aryl, C_{1-3} alkylene-N(R¹²)C(O) C_{1-3} alkyleneN(R¹²)₂, C_{1-3} alkyleneN(R¹²)C(O)-heteroaryl, C_{1-3} alkyleneOR⁷, and SR⁷, wherein R⁷ is as defined above;

R¹¹ is selected from the group consisting of null, hydro, C_{1-6} alkyl, and halo;


R¹² is selected from the group consisting of hydro, C_{1-6} alkyl, cycloalkyl, aryl, and SO₂ C_{1-6} alkyl, or two R¹² groups are taken together to form a 3- to 6-membered ring; and

R¹³ is selected from the group consisting of hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, and aryl;

provided that when Q' is hydrogen or OCH₃, at least one of R⁸, R⁹, and R¹⁰ is not selected from hydrogen, CH₃, OCH₃, or halo,

and pharmaceutically acceptable salts or solvates thereof.

Another aspect of the present invention relates to carbamido-substituted heteroaryl groups having the structural formula (III)



wherein W'' is selected from the group consisting of heteroaryl, aryl, heterocycloalkyl, cycloalkyl, and C_{1-3} alkyl substituted with a heteroaryl or aryl group;

- 14 -

wherein said aryl groups are optionally substituted with one to four substituents represented by R^{14} , said heteroaryl groups are optionally substituted with one to four substituents represented by R^{18} , and said heterocycloalkyl and cycloalkyl groups are optionally substituted with one to two substituents represented by R^{19} ;

R^{14} is selected from the group consisting of halo, C_{1-6} alkyl, C_{2-6} alkenyl, OCF_3 , NO_2 , CN, NC, $N(R^{16})_2$, OR^{16} , CO_2R^{16} , $C(O)N(R^{16})_2$, $C(O)R^{16}$, $N(R^{15})COR^{16}$, $N(R^{15})C(O)OR^{16}$, $N(R^{16})C(O)OR^{16}$, $N(R^{16})C(O)C_{1-3}$ alkylene- $C(O)R^{16}$, $N(R^{16})C(O)C_{1-3}$ alkylene $C(O)OR^{16}$, $N(R^{16})C(O)C_{1-3}$ alkylene OR^{16} , $N(R^{16})C(O)C_{1-3}$ alkyleneNHC(O) OR^{16} , $N(R^{16})C(O)C_{1-3}$ alkylene SO_2NR^{16} , C_{1-3} alkylene OR^{16} , and SR^{16} ;

R^{15} is selected from the group consisting of hydro, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, and aryl;

R^{16} is selected from the group consisting of hydro, C_{1-6} alkyl, C_{2-6} alkenyl, cycloalkyl, aryl, heteroaryl, SO_2R^{17} , C_{1-6} alkyl substituted with one or more of halo, hydroxy, aryl, heteroaryl, heterocycloalkyl, $N(R^{17})_2$, and SO_2R^{17} , C_{1-3} alkylenearyl, C_{1-3} alkyleneheteroaryl, C_{1-3} alkylene C_{3-8} heterocycloalkyl, C_{1-3} alkylene SO_2 aryl, optionally substituted C_{1-3} alkylene $N(R^{17})_2$, OCF_3 , C_{1-3} alkylene $N(R^{17})_3^+$, C_{3-8} heterocycloalkyl, $CH(C_{1-3}alkyleneN(R^{17})_2)_2$;

or two R^{16} groups are taken together to form an optionally substituted 3- to 6-membered aliphatic ring.

R^{17} is selected from the group consisting of hydro, C_{1-6} alkyl, cycloalkyl, aryl, and SO_2C_{1-6} alkyl, or two R^{17} groups are taken together to form an optionally substituted 3- to 6-membered ring;

R^{18} is selected from the group consisting of C_{1-6} alkyl, aryl, $N(R^{15})_2$, OR^{15} , and halo; and

- 16 -


example, Chk1 ensures that cells with unrepaired DNA lesions caused by certain drugs or radiation do not progress through DNA synthesis phase or mitosis until chromosomal lesions have been removed. Accordingly, a tumor cell treated with a Chk1 inhibitor in combination with a DNA damaging agent can kill using lower amounts of DNA damaging agent than tumor cells treated with the DNA damaging agent alone.

Failure of cell cycle checkpoints in normal cells predisposes an individual to, or directly causes, many disease states, such as cancer, ataxia telangiectasia, embryo abnormalities, and various immunological defects associated with aberrant B and T cell development. The latter are associated with the pathological states of lupus, arthritis, and autoimmune diseases. Intense research efforts, therefore, have focused on identifying cell cycle checkpoints and the proteins essential for the function of the checkpoints.

Noncancerous tissue having intact cell checkpoints typically is insulated from temporary disruption of a single checkpoint pathway, such as the Chk1 pathway. Tumor cells, however, have multiple defects in pathways controlling cell cycle progression such that perturbation of the DNA damage checkpoint can render cells particularly sensitive to DNA damaging agents. Therefore, checkpoint inhibitors are expected to enhance the therapeutic index, which is a measure of the probability of tumor control relative to the probability of toxicity to normal tissue to radiation and systemic chemotherapy. In contrast, other classes of inhibitors may not be amenable to combination chemotherapy because both normal and tumor tissue may be similarly sensitized.

- 17 -

One aspect of the present invention is directed to a method of inhibiting Chk1, comprising the step of administering a therapeutically effective amount of a compound of formula (I), or a composition containing the same, to an individual. Compounds of formula (I) have a structural formula



wherein X^1 is null, -O-, -S-, -CH₂-, or -N(R¹)-;

X^2 is -O-, -S-, or -N(R¹)-;

Y is O or S; or =Y represents two hydrogen atoms attached to a common carbon atom;

W is selected from the group consisting of heteroaryl, aryl, heterocycloalkyl, cycloalkyl, and C₁₋₃ alkyl substituted with a heteroaryl or aryl group; and

Z is selected from the group consisting of hydrogen, aryl, and heteroaryl;

wherein said aryl groups of W and Z are optionally substituted with one to four substituents represented by R², said heteroaryl groups of W and Z are optionally substituted with one to four substituents represented by R⁵, and said heterocycloalkyl and cycloalkyl groups of W are optionally substituted with one to two substituents represented by R⁶;

R¹ is selected from the group consisting of hydrogen, C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, and aryl;


- 18 -

R^2 is selected from the group consisting of halogen, C_{1-6} alkyl, C_{2-6} alkenyl, OCF_3 , NO_2 , CN , NC , $N(R^3)_2$, OR^3 , CO_2R^3 , $C(O)N(R^3)_2$, $C(O)R^3$, $N(R^1)COR^3$, $N(R^1)C(O)OR^3$, $N(R^3)C(O)OR^3$, $N(R^3)C(O)C_{1-3}$ alkylene $C(O)R^3$, $N(R^3)C(O)C_{1-3}$ alkylene- $C(O)OR^3$, $N(R^3)C(O)C_{1-3}$ alkylene OR^3 , $N(R^3)C(O)C_{1-3}$ alkylene $NHC(O)OR^3$, $N(R^3)C(O)C_{1-3}$ alkylene- SO_2NR^3 , C_{1-3} alkylene OR^3 , and SR^3 ;

R^3 is selected from the group consisting of hydro, C_{1-6} alkyl, C_{2-6} alkenyl, cycloalkyl, aryl, heteroaryl, SO_2R^4 , C_{1-6} alkyl substituted with one or more of halo, hydroxy, aryl, heteroaryl, heterocycloalkyl, $N(R^4)_2$, and SO_2R^4 , C_{1-3} alkylenearyl, C_{1-3} alkyleneheteroaryl, C_{1-3} alkylene C_{3-8} heterocycloalkyl, C_{1-3} alkylene SO_2 aryl, optionally substituted C_{1-3} alkylene- $N(R^4)_2$, OCF_3 , C_{1-3} alkylene $N(R^4)_3^+$, C_{3-8} heterocycloalkyl, and $CH(C_{1-3}alkyleneN(R^4)_2)_2$, or two R^3 groups are taken together to form an optionally substituted 3- to 6-membered aliphatic ring;

R^4 is selected from the group consisting of hydro, C_{1-6} alkyl, cycloalkyl, aryl, heteroaryl, C_{1-3} alkylenearyl, and SO_2C_{1-6} alkyl, or two R^4 groups are taken together to form an optionally substituted 3- to 6-membered ring;

R^5 is selected from the group consisting of C_{1-6} alkyl, aryl, $N(R^3)_2$, OR^3 , halo, N_3 , CN , C_{1-3} alkylenearyl, C_{1-3} alkylene $N(R^3)_2$, $C(O)R^3$, and




and pharmaceutically acceptable salts or solvates thereof.

Y is 0 or S;

Z is selected from the group consisting

of:



wherein:

J is selected from the group consisting of O, and S;

L is selected from the group consisting of O, and S;

wherein:

- 20 -

R^{20} , R^{21} , and R^{22} are each independently selected from the group consisting of null, hydro, halo, C_{1-6} alkyl, C_{2-6} alkenyl, OCF_3 , NO_2 , CN, NC, $N(R^{25})_2$, OR^{25} , CO_2R^{25} , $C(O)N(R^{25})_2$, $C(O)R^{25}$, $N(R^{24})COR^{25}$, $N(R^{24})-C(O)OR^{25}$, $N(R^{25})C(O)OR^{25}$, $N(R^{25})C(O)C_{1-3}$ alkylene $C(O)R^{25}$, $N(R^{25})C(O)C_{1-3}$ alkylene $C(O)OR^{25}$, $N(R^{25})C(O)C_{1-3}$ alkylene- OR^{25} , $N(R^{25})C(O)C_{1-3}$ alkyleneNHC(O) OR^{25} , $N(R^{25})C(O)C_{1-3}$ alkylene SO_2NR^{25} , CF_3 , C_{1-3} alkylene $N(R^{25})SO_2$ aryl, C_{1-3} alkylene $N(R^{25})SO_2$ heteroaryl, C_{1-3} alkylene OC_{1-3} alkylene-aryl, C_{1-3} alkylene $N(R^{25})C_{1-3}$ alkylenearyl, C_{1-3} alkylene- $N(R^{25})C_{1-3}$ alkyleneheteroaryl, C_{1-3} alkylene $N(R^{25})C(O)R^7$, C_{1-3} alkylene $N(R^{25})C(O)C_{1-3}$ alkylene OR^2 , C_{1-3} alkylene- $N(R^{25})C(O)$ aryl, C_{1-3} alkylene $N(R^{25})C(O)C_{1-3}$ alkylene- $N(R^{25})_2$, C_{1-3} alkylene $N(R^{25})C(O)$ heteroaryl, C_{1-3} alkylene OR^{25} , and SR^{25} ;

R^{23} is selected from the group consisting of null, hydro, C_{1-6} alkyl, and halo;


R^{24} is selected from the group consisting of hydro, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, and aryl;

R^{25} is selected from the group consisting of hydro, C_{1-6} alkyl, C_{2-6} alkenyl, cycloalkyl, aryl, heteroaryl, SO_2R^{26} , and C_{1-6} alkyl substituted with halo, hydroxy, aryl, heteroaryl, heterocycloalkyl, $N(R^{26})_2$, or SO_2R^{26} ; and

R^{26} is selected from the group consisting of hydro, C_{1-6} alkyl, cycloalkyl, aryl, and SO_2C_{1-6} alkyl, or two R^{26} groups are taken together to form an optionally substituted 3- to 6-membered ring.

More preferred compounds of the method are those of structural formula (I) wherein W is selected from the group consisting of pyridazinyl, pyrimidinyl, pyrazinyl, and triazinyl, optionally substituted with from one to four substituents selected from the group consisting of C_{1-6} alkyl,


5



1.0

15

25



30

35

Compounds preferred for use in the method also include those of formula (I) wherein J is se-

- 22 -

lected from the group consisting of CR^{20} and NR^{20} , wherein R^{20} is null, hydro, C_{1-6} alkyl, and halo;

K is selected from the group consisting of CR^{21} and NR^{21} ;

5 L is selected from the group consisting of CR^{22} and NR^{22} ; and

one of R^{21} and R^{22} is hydro and the other is a substituent selected from the group consisting of CO_2R^{25} , $C(O)N(R^{25})_2$, $C(O)R^{25}$, $N(R^{24})COR^{25}$, $N(R^{24})C(O)OR^{25}$,
10 $N(R^{25})C(O)OR^{25}$, $N(R^{25})C(O)C_{1-3}alkyleneC(O)R^{25}$, $N(R^{25})C(O)C_{1-3}alkyleneC(O)OR^{25}$, $N(R^{25})C(O)C_{1-3}alkyleneOR^{25}$, $N(R^{25})C(O)C_{1-3}alkyleneNHC(O)OR^{25}$, $N(R^{25})C(O)C_{1-3}alkyleneSO_2NR^{25}$, $C_{1-3}alkyleneOR^{25}$, and SR^{25} , wherein R^{24} , R^{25} , W, X^1 , X^2 , Y and M are as previously defined.


15 Compounds useful in the method also include structures of formula (I) wherein X^1 is null, X^2 is $-N(H)-$, Y is O, Z is hydro, and W is as previously defined.

The method of inhibiting Chk1 also can be
20 used to sensitize a tumor cell to a chemotherapeutic agent. As such, the present invention also is directed to a method of sensitizing a tumor cell to a chemotherapeutic agent comprising administering a compound of formula (I), or a salt, solvate, or
25 derivative thereof, or a composition comprising the same, to an individual.

In another aspect, the present invention is directed to aryl- and heteroaryl-disubstituted urea compounds having a structural formula (II)

30

- 23 -







(II)

wherein Y' is O or S;


W' is selected from the group consisting

of






5




10



15

20

, and




25


30 optionally substituted with from one to four substituents selected from the group consisting of C₁₋₆alkyl, aryl, N(R⁷)₂, OR⁷, N₃, CN, C(O)R⁷, C₁₋₃alkylenearyl, C₁₋₃alkyleneN(R¹²)₂,

35

- 25 -



10 Z' is selected from the group consisting of:



wherein:

20 Q' is selected from the group consisting of hydrogen, OR^7 , SR^7 , and $N(R^7)_2$;

J' is selected from the group consisting of $C-R^8$, $N-R^8$, O , and S ;

K' is selected from the group consisting of $C-R^9$, $N-R^9$, O , and S ;

25 L' is selected from the group consisting of $C-R^{10}$, $N-R^{10}$, O , and S ;

M' is selected from the group consisting of $C-R^{11}$, $N-R^{11}$, O , and S ;

wherein:

30 R^7 , independently, is selected from the group consisting of hydro, C_{1-6} alkyl, C_{2-6} alkenyl, cycloalkyl, aryl, heteroaryl, SO_2R^{12} , C_{1-6} alkyl substituted with one or more of halo, hydroxy, aryl, heteroaryl, heterocycloalkyl, $N(R^{12})_2$, and SO_2R^{12} ,
 35 C_{1-3} alkylenearyl, C_{1-3} alkyleneheteroaryl, C_{1-3} alkylene-

- 26 -

C₃₋₈heterocycloalkyl, C₁₋₃alkyleneSO₂aryl, optionally substituted C₁₋₃alkyleneN(R¹²)₂, OCF₃, C₁₋₃alkylene-N(R¹²)₃⁺, C₃₋₈heterocycloalkyl, and CH(C₁₋₃alkylene-N(R¹²)₂)₂, or two R⁷ groups are taken together to form
 5 an optionally substituted 3- to 6-membered aliphatic ring;

R⁸, R⁹, and R¹⁰ are each independently selected from the group consisting of null, hydro, halo, optionally substituted C₁₋₆alkyl, C₂₋₆alkenyl,
 10 OCF₃, NO₂, CN, NC, N(R⁷)₂, OR⁷, CO₂R⁷, C(O)N(R⁷)₂, C(O)R⁷, N(R¹³)COR⁷, N(R¹³)C(O)OR⁷, N(R⁷)C(O)OR⁷, N(R⁷)C(O)C₁₋₃alkyleneC(O)R⁷, N(R⁷)C(O)C₁₋₃alkylene-C(O)OR⁷, N(R⁷)C(O)C₁₋₃alkyleneOR⁷, N(R⁷)C(O)C₁₋₃alkyl-eneNHC(O)OR⁷, N(R⁷)C(O)C₁₋₃alkyleneSO₂NR⁷, CF₃, C₁₋₃-
 15 alkyleneN(R¹²)SO₂aryl, C₁₋₃alkyleneN(R¹²)SO₂heteroaryl, C₁₋₃alkyleneOC₁₋₃alkylenearyl, C₁₋₃alkyleneN(R¹²)C₁₋₃-alkylenearyl, C₁₋₃alkyleneN(R¹²)C₁₋₃alkyleneheteroaryl, C₁₋₃alkyleneN(R¹²)C(O)R⁷, C₁₋₃alkyleneN(R¹²)C(O)C₁₋₃alk-yleneOR², C₁₋₃alkyleneN(R¹²)C(O)aryl, C₁₋₃alkyleneN(R¹²)-
 20 C(O)C₁₋₃alkyleneN(R¹²)₂, C₁₋₃alkyleneN(R¹²)C(O)hetero-aryl, C₁₋₃alkyleneOR⁷, and SR⁷, wherein R⁷ is as defined above;

R¹¹ is selected from the group consisting of null, hydro, optionally substituted C₁₋₆alkyl, and
 25 halo;


R¹² is selected from the group consisting of hydro, C₁₋₆alkyl, cycloalkyl, aryl, heteroaryl, C₁₋₃alkylenearyl, and SO₂C₁₋₆alkyl, or two R¹² groups are taken together to form an optionally substituted
 30 3- to 6-membered ring; and

R¹³ is selected from the group consisting of hydro, C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, and aryl; provided that when Q' is hydro or OCH₃, at least one of R⁸, R⁹, and R¹⁰ is different from hydro,
 35 CH₃, OCH₃, and halo,


- 27 -

and pharmaceutically acceptable salts or solvates thereof.


Preferred compounds of structural formula (II) are those wherein W' is selected from the group consisting of



and



optionally substituted with one to four substituents selected from the group consisting of C_{1-6} alkyl, optionally substituted aryl, $N(R^7)_2$, CF_3 , $C(O)R^7$, N_3 , CN, C_{1-3} alkylenearyl, C_{1-3} alkylene $N(R^{12})_2$, OR^7 , halo,



wherein R^7 , Y and Z are as previously defined.

More preferred compounds of formula (II) are those wherein:

- 28 -

J' is selected from the group consisting of CR⁸ and NR⁸, wherein R⁸ is null, hydro, C₁₋₆alkyl, and halo;

5 K' is selected from the group consisting of CR⁹ and NR⁹;

L' is selected from the group consisting of CR¹⁰ and NR¹⁰; and

one of R⁹ and R¹⁰ is hydro and the other is a substituent selected from the group consisting of
 10 CO₂R⁷, C(O)N(R⁷)₂, C(O)R⁷, N(R¹³)COR⁷, N(R¹³)C(O)OR⁷,
 N(R⁷)C(O)OR⁷, N(R⁷)C(O)C₁₋₃alkyleneC(O)R⁷, N(R⁷)C-
 (O)C₁₋₃alkyleneC(O)OR⁷, N(R⁷)C(O)C₁₋₃alkyleneOR⁷,
 N(R⁷)C(O)C₁₋₃alkyleneNHC(O)OR⁷, N(R⁷)C(O)C₁₋₃alkylene-
 SO₂NR⁷, C₁₋₃alkyleneOR⁷, CF₃, C₁₋₃alkyleneN(R¹²)SO₂aryl,
 15 C₁₋₃alkyleneN(R¹²)SO₂heteroaryl, C₁₋₃alkyleneOC₁₋₃alkyl-
 enearyl, C₁₋₃alkyleneN(R¹²)C₁₋₃alkylenearyl, C₁₋₃alk-
 yleneN(R¹²)C₁₋₃alkyleneheteroaryl, C₁₋₃alkyleneN(R¹²)-
 C(O)R⁷, C₁₋₃alkyleneN(R¹²)C(O)C₁₋₃alkyleneOR², C₁₋₃alk-
 yleneN(R¹²)C(O)aryl, C₁₋₃alkyleneN(R¹²)C(O)C₁₋₃alkylene-
 20 N(R¹²)₂, C₁₋₃alkyleneN(R¹²)C(O)heteroaryl, and SR⁷,
 wherein R⁷, R¹³, W', Y', and M' are as previously
 defined.

Yet another aspect of the invention re-
 25 lates to compounds, and compositions containing
 compounds, of structural formula (III)



- 29 -

wherein W'" is selected from the group consisting of heteroaryl, aryl, heterocycloalkyl, cycloalkyl, and C₁₋₃ alkyl substituted with a heteroaryl or aryl group;

5 wherein said aryl groups are optionally substituted with one to four substituents represented by R¹⁴, said heteroaryl groups are optionally substituted with one to four substituents represented by R¹⁸, and said heterocycloalkyl and cycloalkyl groups are optionally substituted with one to
10 two substituents represented by R¹⁹;

R¹⁴ is selected from the group consisting of halo, C₁₋₆alkyl, C₂₋₆alkenyl, OCF₃, NO₂, CN, NC, N(R¹⁶)₂, OR¹⁶, CO₂R¹⁶, C(O)N(R¹⁶)₂, C(O)R¹⁶, N(R¹⁵)COR¹⁶,
15 N(R¹⁵)C(O)OR¹⁶, N(R¹⁶)C(O)OR¹⁶, N(R¹⁶)C(O)C₁₋₃alkylene-C(O)R¹⁶, N(R¹⁶)C(O)C₁₋₃alkyleneC(O)OR¹⁶, N(R¹⁶)C(O)-C₁₋₃alkyleneOR¹⁶, N(R¹⁶)C(O)C₁₋₃alkyleneNHC(O)OR¹⁶, N(R¹⁶)C(O)C₁₋₃alkyleneSO₂NR¹⁶, C₁₋₃alkyleneOR¹⁶, and SR¹⁶;

R¹⁵ is selected from the group consisting
20 of hydro, C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, and aryl;

R¹⁶ is selected from the group consisting of hydro, C₁₋₆alkyl, C₂₋₆alkenyl, cycloalkyl, aryl, heteroaryl, SO₂R¹⁷, and C₁₋₆alkyl substituted with one or more of halo, hydroxy, aryl, heteroaryl, heterocycloalkyl, N(R¹⁷)₂, and SO₂R¹⁷, C₁₋₃alkylenearyl, C₁₋₃alkyleneheteroaryl, C₁₋₃alkyleneC₃₋₈heterocycloalkyl, C₁₋₃alkyleneSO₂aryl, optionally substituted C₁₋₃alkyleneN(R¹⁷)₂, OCF₃, C₁₋₃alkyleneN(R¹⁷)₃⁺, C₃₋₈heterocycloalkyl, CH(C₁₋₃alkyleneN(R¹⁷)₂)₂, or two R¹⁶ groups are
25 taken together to form a 3- to 6-membered aliphatic ring.
30

R¹⁷ is selected from the group consisting of hydro, C₁₋₆alkyl, cycloalkyl, aryl, and SO₂C₁₋₆alkyl, or two R¹⁷ groups are taken together to form
35 an optionally substituted 3- to 6-membered ring;

R¹⁹ is selected from the group consisting of halo and C₁₋₆alkyl.

As used herein, the term "alkyl" includes straight chained and branched hydrocarbon groups containing the indicated number of carbon atoms, typically methyl, ethyl, and straight-chain and branched propyl and butyl groups. Unless otherwise indicated, the hydrocarbon group can contain up to 20 carbon atoms. The term "alkyl" includes "bridged alkyl," i.e., a C₆-C₁₆ bicyclic or polycyclic hydrocarbon group, for example, norbornyl, adamantyl, bicyclo[2.2.2]octyl, bicyclo[2.2.1]heptyl, bicyclo[3.2.1]octyl, or decahydronaphthyl. Alkyl groups can be substituted, for example, with hydroxy (OH), halo, aryl, heteroaryl, cycloalkyl, heterocycloalkyl, amino (N(R^a)₂), and sulfonyl (SO₂R^a), wherein R^a is selected from the group consisting of hydro, C₁₋₆alkyl, cycloalkyl, aryl, and SO₂C₁₋₆alkyl, or two R^a groups are taken together to form an optionally substituted 3- to 6-membered ring.

The term "cycloalkyl" is defined as a cyclic C₃₋₈hydrocarbon group, e.g., cyclopropyl, cyclobutyl, cyclohexyl, and cyclopentyl. "Heterocycloalkyl" is defined similarly as cycloalkyl, and includes bicyclic and polycyclic groups, except the ring contains one to three heteroatoms selected from the group consisting of oxygen, nitrogen, and sulfur. Cycloalkyl and heterocycloalkyl groups can be saturated or partially unsaturated ring systems substituted with, for example, one to three groups, independently selected from C₁₋₄alkyl, C₁₋₃alkyleneOH, C(=O)NH₂, NH₂, oxo (=O), aryl, trifluoroethanoyl, and OH. Heterocycloalkyl groups optionally are further

The term "alkenyl" is defined identically as "alkyl," except the substituent contains a carbon-carbon double bond.

The term "alkylene" refers to an alkyl group having a substituent. For example, the term "C₁₋₃alkyleneC(O)OR" refers to an alkyl group containing one to three carbon atoms substituted with a -C(O)OR group. The alkylene group is optionally substituted with one or more of aryl, heteroaryl, and OR⁷, wherein R⁷ is defined hereafter.

The term "aryl," alone or in combination, is defined herein as a monocyclic or polycyclic aromatic group, preferably a monocyclic or bicyclic aromatic group, e.g., phenyl or naphthyl. Unless otherwise indicated, an "aryl" group can be unsubstituted or substituted, for example, with one or more, and in particular one to four, halo, C₁₋₆alkyl, C₂₋₆alkenyl, OCF₃, NO₂, CN, NC, N(R^a)₂, OR^b, CO₂R^b, C(O)N(R^b)₂, C(O)R^b, N(R^a)COR^b, N(R^a)C(O)OR^b, N(R^a)C(O)-OR^b, N(R^a)C(O)C₁₋₃alkyleneC(O)R^b, N(R^b)C(O)C₁₋₃alkylene-C(O)OR^b, N(R^b)C(O)C₁₋₃alkyleneOR^b, N(R^b)C(O)C₁₋₃alkyleneNHC(O)OR^b, N(R^b)C(O)C₁₋₃alkyleneSO₂NR^b, C₁₋₃alkyleneOR^b, and SR^b, wherein R^b is selected from the group consisting of hydro, C₁₋₆alkyl, C₂₋₆alkenyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, SO₂R^a, and C₁₋₆alkyl substituted with halo, hydroxy, aryl, heteroaryl, heterocycloalkyl, N(R^a)₂, or SO₂R^a,

The term "heteroaryl" is defined herein as a monocyclic or bicyclic ring system containing one or two aromatic rings and containing at least one nitrogen, oxygen, or sulfur atom in an aromatic ring, and that can be unsubstituted or substituted, for example, with one or more, and in particular one to four, substituents, for example, hydrogen, C₁₋₆-alkyl, C₁₋₆alkoxy, aryl, N(R^a)₂, OR^b, and halo, wherein R^a and R^b are as previously defined. Examples of heteroaryl groups include, but are not limited to, thienyl, furyl, pyridyl, oxazolyl, quinolyl, isoquinolyl, indolyl, triazolyl, isothiazolyl, isoxazolyl, imidizolyl, benzothiazolyl, pyrazinyl, pyrimidinyl, thiazolyl, and thiadiazolyl.


The term "3- to 6-membered ring" as used refers to carbocyclic and heterocyclic or aromatic groups, including, but not to, morpholinyl, piperidinyl, phenyl, thioaryl, pyrrolyl, imidazolyl, pyrimidinyl, inyl, optionally substituted with one or in particular one to three, groups as defined above for "aryl" groups.

The carbon atom content of hydrocarbon-containing moieties is indicated by a subscript designating the minimum and maximum number of carbon


- 33 -

atoms in the moiety, e.g., "C₁₋₆alkyl" refers to an alkyl group having one to six carbon atoms, inclusive.

In the structures herein, for a bond lacking a substituent, the substituent is methyl, for example,



When no substituent is indicated as attached to a carbon atom on a ring, it is understood that the carbon atom contains the appropriate number of hydrogen atoms. In addition, when no substituent is indicated as attached to a carbonyl group or a nitrogen atom, for example, the substituent is understood to be hydrogen, e.g.,



The abbreviation "Me" is methyl. The abbreviation CO and C(O) is carbonyl (C=O).

The notation N(R^x)₂, wherein x represents an alpha or numeric character, such as for example R^a, R^b, R⁴, R¹², and the like, is used to denote two R^x groups attached to a common nitrogen atom. When used in such notation, the R^x group can be the same

- 34 -

or different, and is selected from the group as defined by the R^{*} group.

The present invention also is directed to pharmaceutical compositions containing one or more compounds of structural formula (II) and (III), to use of the compounds and compositions containing the compounds in therapeutic treatment of a disease or disorder, and to methods of preparing the compounds and intermediates involved in the synthesis of the compounds of structural formula (II) and (III).

Compounds useful for the method of the present invention have demonstrated activity in inhibiting Chk1 *in vitro*. Compounds of the present invention have demonstrated selectivity for Chk1 as against other protein kinases including Cdc2, Chk2, Atr, DNA-PK, PKA, and CaM KII.

Compounds of the present invention can be used to potentiate the therapeutic effects of radiation and/or chemotherapeutics used in the treatment of cancers and other cell proliferation disorders in humans or animals. For example, compounds of the invention can be used to enhance treatment of tumors that are customarily treated with an antimetabolite, e.g., methotrexate or 5-fluorouracil (5-FU). The method of the present invention comprises administration of a Chk1 inhibitor compound in combination with a chemotherapeutic agent that can effect single- or double-strand DNA breaks or that can block DNA replication or cell proliferation.

Alternatively, the method of the present invention comprises administration of a Chk1 inhibitor compound in combination with therapies that include use of an antibody, e.g., herceptin, that has activity in inhibiting the proliferation of cancer cells.

Accordingly, cancers such as colorectal cancers,

- 35 -

head and neck cancers, pancreatic cancers, breast cancers, gastric cancers, bladder cancers, vulvar cancers, leukemias, lymphomas, melanomas, renal cell carcinomas, ovarian cancers, brain tumors, osteosarcomas, and lung carcinomas, are susceptible to enhanced treatment in combination with the Chk1 inhibitors of the invention.

Tumors or neoplasms include growths of tissue cells wherein multiplication of cells is uncontrolled and progressive. Some such growths are benign, but others are termed "malignant," and can lead to death of the organism. Malignant neoplasms, or "cancers," are distinguished from benign growths in that, in addition to exhibiting aggressive cellular proliferation, can invade surrounding tissues and metastasize. Moreover, malignant neoplasms are characterized by showing a greater loss of differentiation (greater "dedifferentiation") and organization relative to one another and surrounding tissues. This property is called "anaplasia."

Neoplasms treatable by the present invention also include solid tumors, i.e., carcinomas and sarcomas. Carcinomas include malignant neoplasms derived from epithelial cells which infiltrate (i.e., invade) surrounding tissues and give rise to metastases. Adenocarcinomas are carcinomas derived from glandular tissue, or from tissues that form recognizable glandular structures. Another broad category of cancers includes sarcomas, which are tumors whose cells are embedded in a fibrillar or homogeneous substance, like embryonic connective tissue. The invention also enables treatment of cancers of the myeloid or lymphoid systems, including leukemias, lymphomas, and other cancers that typically are not present as a tumor mass, but

are distributed in the vascular or lymphoreticular systems.

Chk1 activity is associated with various forms of cancer in, for example, adult and pediatric oncology, growth of solid tumors/malignancies, myxoid and round cell carcinoma, locally advanced tumors, metastatic cancer, human soft tissue sarcomas, including Ewing's sarcoma, cancer metastases, including lymphatic metastases, squamous cell carcinoma, particularly of the head and neck, esophageal squamous cell carcinoma, oral carcinoma, blood cell malignancies, including multiple myeloma, leukemias, including acute lymphocytic leukemia, acute nonlymphocytic leukemia, chronic lymphocytic leukemia, chronic myelocytic leukemia, and hairy cell leukemia, effusion lymphomas (body cavity based lymphomas), thymic lymphoma lung cancer (including small cell carcinoma, cutaneous T cell lymphoma, Hodgkin's lymphoma, non-Hodgkin's lymphoma, cancer of the adrenal cortex, ACTH-producing tumors, nonsmall cell cancers, breast cancer, including small cell carcinoma and ductal carcinoma), gastrointestinal cancers (including stomach cancer, colon cancer, colorectal cancer, and polyps associated with colorectal neoplasia), pancreatic cancer, liver cancer, urological cancers (including bladder cancer, such as primary superficial bladder tumors, invasive transitional cell carcinoma of the bladder, and muscle-invasive bladder cancer), prostate cancer, malignancies of the female genital tract (including ovarian carcinoma, primary peritoneal epithelial neoplasms, cervical carcinoma, uterine endometrial cancers, vaginal cancer, cancer of the vulva, uterine cancer and solid tumors in the ovarian follicle), malignancies of the male genital

- 37 -

tract (including testicular cancer and penile cancer), kidney cancer (including renal cell carcinoma, brain cancer (including intrinsic brain tumors, neuroblastoma, astrocytic brain tumors, gliomas, and metastatic tumor cell invasion in the central nervous system), bone cancers (including osteomas and osteosarcomas), skin cancers (including malignant melanoma, tumor progression of human skin keratinocytes, and squamous cell cancer), thyroid cancer, retinoblastoma, neuroblastoma, peritoneal effusion, malignant pleural effusion, mesothelioma, Wilms's tumors, gall bladder cancer, trophoblastic neoplasms, hemangiopericytoma, and Kaposi's sarcoma.

Compounds of the present invention also can potentiate the efficacy of drugs in the treatment of inflammatory diseases. Examples of diseases that can benefit from combination therapy with compounds suitable for the method of the present invention are rheumatoid arthritis, psoriasis, vitiligo, Wegener's granulomatosis, and systemic lupus erythematosus (SLE). Treatment of arthritis, Wegener's granulomatosis, and SLE often involves the use of immunosuppressive therapies, such as ionizing radiation, methotrexate, and cyclophosphamide. Such treatments typically induce, either directly or indirectly, DNA damage. Inhibition of Chk1 activity within the offending immune cells render the cells more sensitive to control by these standard treatments. Psoriasis and vitiligo commonly are treated with ultraviolet radiation (UV) in combination with psoralen. The present DNA damaging agents induce the killing effect of UV and psoralen, and increase the therapeutic index of this treatment regimen. In general, compounds useful in methods of the present invention potentiate control of inflammatory disease

- 38 -

cells when in combination with currently used immunosuppressive drugs.

The present invention includes all possible stereoisomers and geometric isomers of compounds of the present method and of structural formulae (I), (II), and (III). The present invention includes not only racemic compounds, but optically active isomers as well. When a compound of structural formulae (I), (II), or (III) is desired as a single enantiomer, it can be obtained either by resolution of the final product or by stereospecific synthesis from either isomerically pure starting material or use of a chiral auxiliary reagent, for example, see Z. Ma et al., *Tetrahedron: Asymmetry*, 8(6), pages 883-888 (1997). Resolution of the final product, an intermediate, or a starting material can be achieved by any suitable method known in the art. Additionally, in situations where tautomers of the compounds of structural formulae (I), (II), and (III) are possible, the present invention is intended to include all tautomeric forms of the compounds. As demonstrated hereafter, specific stereoisomers can exhibit an exceptional ability to inhibit Chk1 in combination with chemoradiotherapy with diminished adverse effects typically associated with chemotherapeutic or radiotherapeutic treatments.

Prodrugs of compositions of structural formulae (I), (II), and (III) also can be used as the compound and in the method of the present invention. It is well established that a prodrug approach, wherein a compound is derivatized into a form suitable for formulation and/or administration, and then is released as a drug *in vivo*, has been successfully employed to transiently (e.g., biore-

- 39 -

versibly) alter the physicochemical properties of the compound (see, H. Bundgaard, Ed., *Design of Prodrugs*, Elsevier, Amsterdam, (1985); R.B. Silverman, *The Organic Chemistry of Drug Design and Drug Action*, Academic Press, San Diego, chapter 8, (1992); K.M. Hillgren et al., *Med. Res. Rev.*, 15, 83 (1995)).

Compounds of the present invention can contain several functional groups. The introduced functional groups, if desired or necessary, then can be modified to provide a prodrug for dose of formulation and/or administration. Suitable prodrugs include, for example, acid derivatives, like amides, esters, and the like. It also is appreciated by those skilled in the art that N-oxides can be used as a prodrug.

As used herein, the term pharmaceutically acceptable salts refers compounds of structural formula (I), (II), and (III) which contain acidic moieties and form salts with suitable cations. Suitable pharmaceutically acceptable cations include alkali metal (e.g., sodium or potassium) and alkaline earth metal (e.g., calcium or magnesium) cations. The pharmaceutically acceptable salts of the compounds of structural formula (I), (II), and (III), which contain a basic center, are acid addition salts formed with pharmaceutically acceptable acids. Examples include the hydrochloride, hydrobromide, sulfate or bisulfate, phosphate or hydrogen phosphate, acetate, benzoate, succinate, fumarate, maleate, lactate, citrate, tartrate, gluconate, methanesulfonate, benzene sulphonate, and p-toluenesulphonate salts. In light of the foregoing, any reference to compounds of the present invention appearing herein is intended to include compounds of

- 40 -

structural formula (I), (II), and (III), as well as pharmaceutically acceptable salts and solvates thereof.

The compounds of the present invention can be therapeutically administered as the neat chemical, but it is preferable to administer compounds of structural formula (I), (II), and (III) as a pharmaceutical composition or formulation. Accordingly, the present invention further provides pharmaceutical formulations comprising a compound of structural formula (I), (II), and/or (III), or pharmaceutically acceptable salts thereof, together with one or more pharmaceutically acceptable carriers and, optionally, other therapeutic and/or prophylactic ingredients. The carriers are "acceptable" in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

Inhibition of the checkpoint kinase typically is measured using a dose-response assay in which a sensitive assay system is contacted with a compound of interest over a range of concentrations, including concentrations at which no or minimal effect is observed, through higher concentrations at which partial effect is observed, to saturating concentrations at which a maximum effect is observed. Theoretically, such assays of the dose-response effect of inhibitor compounds can be described as a sigmoidal curve expressing a degree of inhibition as a function of concentration. The curve also theoretically passes through a point at which the concentration is sufficient to reduce activity of the checkpoint enzyme to a level that is 50% that of the difference between minimal and maximal enzyme activity in the assay. This concentration is defined as

- 41 -

the Inhibitory Concentration (50%) or IC_{50} value. Determination of IC_{50} values preferably are made using conventional biochemical (acellular) assay techniques or cell-based assay techniques.

5 Comparisons of the efficacy of inhibitors often are provided with reference to comparative IC_{50} values, wherein a higher IC_{50} indicates that the test compound is less potent, and a lower IC_{50} indicates that the compound is more potent, than a reference
10 compound. Compounds useful in the method of the present invention demonstrate an IC_{50} value of at least 0.1 nM when measured using the dose-response assay. Preferred compounds demonstrate an IC_{50} value of less than 10 μ M. More preferred compounds demon-
15 strate an IC_{50} value of less than 500 nM. Still more preferred compounds of the present invention demon- strate an IC_{50} value of less than 250 nM, less than 100 nM, or less than 50 nM.

 Compounds and pharmaceutical compositions
20 suitable for use in the present invention include those wherein the active ingredient is administered in an effective amount to achieve its intended purpose. More specifically, a "therapeutically effective amount" means an amount effective to
25 inhibit development of, or to alleviate the existing symptoms of, the subject being treated. Determination of the effective amount is well within the capability of those skilled in the art, especially in light of the detailed disclosure provided herein.

30 A "therapeutically effective dose" refers to that amount of the compound that results in achieving the desired effect. Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cul-
35 tures or experimental animals, e.g., for determining

- 42 -

the LD₅₀ (the dose lethal to 50% of the population) and the ED₅₀ (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index, which is expressed as the ratio of LD₅₀ to ED₅₀. Compounds that exhibit high therapeutic indices (i.e., a toxic dose that is substantially higher than the effective dose) are preferred. The data obtained can be used in formulating a dosage range for use in humans. The dosage of such compounds preferably lies within a range of circulating concentrations that include the ED₅₀ with little or no toxicity. The dosage can vary within this range depending upon the dosage form employed, and the route of administration utilized.

The exact formulation, route of administration, and dosage is chosen by the individual physician in view of the patient's condition. Dosage amount and interval can be adjusted individually to provide plasma levels of the active compound that are sufficient to maintain desired therapeutic effects.

Pharmaceutical compositions of the invention can be formulated to include one or more cytokines, lymphokines, growth factors, or other hematopoietic factors which can reduce negative side effects that may arise from, or be associated with, administration of the pharmaceutical composition alone. Cytokines, lymphokines, growth factors, or other hematopoietic factors particularly useful in pharmaceutical compositions of the invention include, but are not limited to, M-CSF, GM-CSF, TNF, IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, IL-15, IL-16, IL-17, IL-18, IFN, TNF, G-CSF, Meg-CSF, GM-CSF,

- 43 -

thrombopoietin, stem cell factor, erythropoietin,
angiopoietins, including Ang-1, Ang-2, Ang-4, Ang-Y,
and/or the human angiopoietin-like polypeptide,
vascular endothelial growth factor (VEGF), angio-
5 genin, bone morphogenic protein-1 (BMP-1), BMP-2,
BMP-3, BMP-4, BMP-5, BMP-6, BMP-7, BMP-8, BMP-9,
BMP-10, BMP-11, BMP-12, BMP-13, BMP-14, BMP-15, BMP
receptor IA, BMP receptor IB, brain derived neuro-
trophic factor, ciliary neutrophic factor, ciliary
10 neutrophic factor receptor cytokine-induced neutro-
phil chemotactic factor 1, cytokine-induced neutro-
phil chemotactic factor 2, cytokine-induced neutro-
phil chemotactic factor 2, endothelial cell growth
factor, endothelin 1, epidermal growth factor,
15 epithelial-derived neutrophil attractant, fibroblast
growth factor (FGF) 4, FGF 5, FGF 6, FGF 7, FGF 8,
FGF 8b, FGF 8c, FGF 9, FGF 10, FGF acidic, FGF
basic, glial cell line-derived neutrophic factor
receptor 1, glial cell line-derived neutrophic
20 factor receptor 2, growth related protein, growth
related protein, growth related protein, growth
related protein, heparin binding epidermal growth
factor, hepatocyte growth factor, hepatocyte growth
factor receptor, insulin-like growth factor I,
25 insulin-like growth factor receptor, insulin-like
growth factor II, insulin-like growth factor binding
protein, keratinocyte growth factor, leukemia inhib-
itory factor, leukemia inhibitory factor receptor,
nerve growth factor nerve growth factor receptor,
30 neurotrophin-3, neurotrophin-4, placenta growth
factor, placenta growth factor 2, platelet-derived
endothelial cell growth factor, platelet derived
growth factor, platelet derived growth factor A
chain, platelet derived growth factor AA, platelet
35 derived growth factor AB, platelet derived growth

- 44 -

factor B chain, platelet derived growth factor BB, platelet derived growth factor receptor, platelet derived growth factor receptor, pre-B cell growth stimulating factor, stem cell factor, stem cell factor receptor, transforming growth factor (TGF), TGF, TGF 1, TGF 1.2, TGF 2, TGF 3, TGF 5, latent TGF 1, TGF, binding protein I, TGF binding protein II, TGF binding protein III, tumor necrosis factor receptor type I, tumor necrosis factor receptor type II, urokinase-type plasminogen activator receptor, vascular endothelial growth factor, and chimeric proteins and biologically or immunologically active fragments thereof.

The compounds useful according to the invention may be conjugated or linked to auxiliary moieties that promote any property of the compounds that may be beneficial in methods of therapeutic use. Such conjugates can enhance delivery of the compounds to a particular anatomical site or region of interest (e.g., a tumor), enable sustained therapeutic concentrations of the compounds in target cells, alter pharmacokinetic and pharmacodynamic properties of the compounds, and/or improve the therapeutic index or safety profile of the compounds. Suitable auxiliary moieties include, for example, amino acids, oligopeptides, or polypeptides, e.g., antibodies such as monoclonal antibodies and other engineered antibodies; and natural or synthetic ligands to receptors in target cells or tissues. Other suitable auxiliaries include fatty acid or lipid moieties, to promote biodistribution or uptake of the compound by target cells (see, e.g., Bradley et al., *Clin. Cancer Res.* (2001) 7:3229.

- 45 -

The therapeutic index of compositions comprising one or more compounds of the invention can be enhanced by conjugation of the compound(s) with antitumor antibodies as previously described (for example, Pietersz and McKinzie, *Immunol. Rev.* (1992) 129:57; Trail et al., *Science* (1993) 261:212; Rowlinson-Busza and Epenetos, *Curr. Opin. Oncol.* 1992; 4:1142). Tumor directed delivery of compounds of the invention would enhance the therapeutic benefit by minimizing potential nonspecific toxicities which can result from radiation treatment or chemotherapy. In another aspect, Chk1 inhibitors and radioisotopes or chemotherapeutic agents can be conjugated to the same antibody molecule. Alternatively, Chk1 inhibitor-conjugated tumor specific antibodies can be administered before, during, or after administration of chemotherapeutic-conjugated antitumor antibody or radioimmunotherapy.

Compounds of the present invention can enhance the therapeutic benefit of radiation and chemotherapy treatment, including induction chemotherapy, primary (neoadjuvant) chemotherapy, and both adjuvant radiation therapy and adjuvant chemotherapy. In addition, radiation and chemotherapy are frequently indicated as adjuvants to surgery in the treatment of cancer. The goal of radiation and chemotherapy in the adjuvant setting is to reduce the risk of recurrence and enhance disease-free survival when the primary tumor has been controlled. Chemotherapy is utilized as a treatment adjuvant for colon, lung, and breast cancer, frequently when the disease is metastatic. Adjuvant radiation therapy is indicated in several diseases including colon, lung, and breast cancers as described above. For example, radiation frequently is used both pre- and

- 46 -

post-surgery as components of the treatment strategy for rectal carcinoma. Compounds for the present invention are therefore particularly useful following surgery in the treatment of cancer in combination with radio- and/or chemotherapy.

A compound of the present invention also can radiosensitize a cell. The term "radiosensitize," as used herein, is defined as a molecule, preferably a low molecular weight molecule, administered to human or other animal in a therapeutically effective amount to increase the sensitivity of the cells to be radiosensitized to electromagnetic radiation and/or to promote the treatment of diseases that are treatable with electromagnetic radiation. Diseases that are treatable with electromagnetic radiation include neoplastic diseases, benign and malignant tumors, and cancerous cells.

Electromagnetic radiation treatment of other diseases not listed herein also is contemplated by the present invention. The terms "electromagnetic radiation" and "radiation" as used herein include, but are not limited to, radiation having the wavelength of 10^{-20} to 100 meters. Preferred embodiments of the present invention employ the electromagnetic radiation of: gamma-radiation (10^{-20} to 10^{-13} m), X-ray radiation (10^{-12} to 10^{-9} m), ultraviolet light (10 nm to 400 nm), visible light (400 nm to 700 nm), infrared radiation (700 nm to 1.0 mm), and microwave radiation (1 mm to 30 cm).

Many cancer treatment protocols currently employ radiosensitizers activated by electromagnetic radiation, e.g., X-rays. Examples of X-ray-activated radiosensitizers include, but are not limited to, the following: metronidazole, misonidazole,

- 47 -

desmethylnisonidazole, pimonidazole, etanidazole, nimorazole, mitomycin C, RSU 1069, SR 4233, EO9, RB 6145, nicotinamide, 5-bromodeoxyuridine (BUdR), 5-iododeoxyuridine (IUdR), bromodeoxycytidine, fluoro-deoxyuridine (FUdR), hydroxyurea, cisplatin, and therapeutically effective analogs and derivatives of the same.

Photodynamic therapy (PDT) of cancers employs visible light as the radiation activator of the sensitizing agent. Examples of photodynamic radiosensitizers include the following, but are not limited to: hematoporphyrin derivatives, PHOTO-FRIN®, benzoporphyrin derivatives, NPe6, tin etio-porphyrin (SnET2), pheoborbide-a, bacteriochloro-phyll-a, naphthalocyanines, phthalocyanines, zinc phthalocyanine, and therapeutically effective analogs and derivatives of the same.

Radiosensitizers can be administered in conjunction with a therapeutically effective amount of one or more compounds in addition to the Chk1 inhibitor, such compounds including, but not limited to, compounds that promote the incorporation of radiosensitizers to the target cells, compounds that control the flow of therapeutics, nutrients, and/or oxygen to the target cells, chemotherapeutic agents that act on the tumor with or without additional radiation, or other therapeutically effective compounds for treating cancer or other disease. Examples of additional therapeutic agents that can be used in conjunction with radiosensitizers include, but are not limited to, 5-fluorouracil (5-FU), leucovorin, oxygen, carbogen, red cell transfusions, perfluorocarbons (e.g., FLUOSOLW®-DA), 2,3-DPG, BW12C, calcium channel blockers, pentoxifylline, antiangiogenesis compounds, hydralazine, and L-BSO.

- 48 -

Chemotherapeutic agents that can be used include, but are not limited to, alkylating agents, antimetabolites, hormones and antagonists thereof, radioisotopes, antibodies, as well as natural products, and combinations thereof. For example, an inhibitor compound of the present invention can be administered with antibiotics, such as doxorubicin and other anthracycline analogs, nitrogen mustards, such as cyclophosphamide, pyrimidine analogs such as 5-fluorouracil, cisplatin, hydroxyurea, taxol and its natural and synthetic derivatives, and the like. As another example, in the case of mixed tumors, such as adenocarcinoma of the breast, where the tumors include gonadotropin-dependent and gonadotropin-independent cells, the compound can be administered in conjunction with leuprolide or goserelin (synthetic peptide analogs of LH-RH). Other anti-neoplastic protocols include the use of an inhibitor compound with another treatment modality, e.g., surgery or radiation, also referred to herein as "adjunct anti-neoplastic modalities." Examples of chemotherapeutic agents useful for the method of the present invention are listed in the following table.

<p>5</p> <p>10</p> <p>15</p> <p>20</p> <p>25</p> <p>30</p> <p>35</p> <p>40</p> <p>45</p> <p>50</p> <p>55</p> <p>60</p> <p>65</p>	<p><u>Alkylating agents</u></p> <p><u>Nitrogen mustards</u></p> <p>mechlorethamine</p> <p>cyclophosphamide</p> <p>ifosfamide</p> <p>melphalan</p> <p>chlorambucil</p> <p><u>Nitrosoureas</u></p> <p>carmustine (BCNU)</p> <p>lomustine (CCNU)</p> <p>semustine (methyl-CCNU)</p> <p><u>Ethylenimine/Methyl-</u></p> <p><u>melamine</u></p> <p>thiethylenemelamine</p> <p>(TEM)</p> <p>triethylene</p> <p>thiophosphoramide</p> <p>(thiotepa)</p> <p>hexamethylmelamine</p> <p>(HMM, altretamine)</p> <p><u>Alkyl sulfonates</u></p> <p>busulfan</p> <p><u>Triazines</u></p> <p>dacarbazine (DTIC)</p> <p><u>Antimetabolites</u></p> <p><u>Folic Acid analogs</u></p> <p>methotrexate</p> <p>trimetrexate</p> <p><u>Pyrimidine analogs</u></p> <p>5-fluorouracil</p> <p>fluorodeoxyuridine</p> <p>gemcitabine</p> <p>cytosine arabinoside</p> <p>(AraC, cytarabine)</p> <p>5-azacytidine</p> <p>2,2'-difluorodeoxy-</p> <p>cytidine</p> <p><u>Purine analogs</u></p> <p>6-mercaptopurine</p> <p>6-thioguanine</p> <p>azathioprine</p> <p>2'-deoxycoformycin</p> <p>(pentostatin)</p> <p>erythrohydroxynonyl-</p> <p>adenine (EHNA)</p> <p>fludarabine phosphate</p> <p>2-chlorodeoxyadenosine</p> <p>(cladribine, 2-CdA)</p> <p><u>Type I Topoisomerase</u></p> <p><u>Inhibitors</u></p> <p>camptothecin</p> <p>topotecan</p> <p>irinotecan</p> <p><u>Natural products</u></p> <p><u>Antimitotic drugs</u></p> <p>paclitaxel</p> <p>Vinca alkaloids</p> <p>vinblastine (VLB)</p> <p>vincristine</p> <p>vinorelbine</p> <p>Taxotere® (docetaxel)</p> <p>estramustine</p> <p>estramustine phosphate</p>	<p><u>Epipodophylotoxins</u></p> <p>etoposide</p> <p>teniposide</p> <p><u>Antibiotics</u></p> <p>actinomycin D</p> <p>daunomycin (rubido-</p> <p>mycin)</p> <p>doxorubicin (adria-</p> <p>mycin)</p> <p>mitoxantroneidarubicin</p> <p>bleomycin</p> <p>splicamycin</p> <p>(mithramycin)</p> <p>mitomycinC</p> <p>dactinomycin</p> <p><u>Enzymes</u></p> <p>L-asparaginase</p> <p><u>Biological response</u></p> <p><u>modifiers</u></p> <p>interferon-alpha</p> <p>IL-2</p> <p>G-CSF</p> <p>GM-CSF</p> <p><u>Differentiation Agents</u></p> <p>retinoic acid deriva-</p> <p>tives</p> <p><u>Radiosensitizers</u></p> <p>metronidazole</p> <p>misonidazole</p> <p>desmethylmisonidazole</p> <p>pimonidazole</p> <p>etanidazole</p> <p>nimorazole</p> <p>RSU 1069</p> <p>EO9</p> <p>RB 6145</p> <p>SR4233</p> <p>nicotinamide</p> <p>5-bromodeoxyuridine</p> <p>5-iododeoxyuridine</p> <p>bromodeoxycytidine</p> <p><u>Miscellaneous agents</u></p> <p><u>Platinum coordination</u></p> <p><u>complexes</u></p> <p>cisplatin</p> <p>carboplatin</p> <p><u>Anthracenedione</u></p> <p>mitoxantrone</p> <p><u>Substituted urea</u></p> <p>hydroxyurea</p> <p><u>Methylhydrazine deriva-</u></p> <p><u>tives</u></p> <p>N-methylhydrazine (MIH)</p> <p>procarbazine</p> <p><u>Adrenocortical suppres-</u></p> <p><u>sant</u></p> <p>mitotane (o,p'-DDD)</p> <p>ainoglutethimide</p> <p><u>Cytokines</u></p> <p>interferon (α, β, γ)</p> <p>interleukin-2</p>	<p><u>Hormones and antagonists</u></p> <p><u>Adrenocorticosteroids/</u></p> <p><u>antagonists</u></p> <p>prednisone and equiv-</p> <p>alents</p> <p>dexamethasone</p> <p>ainoglutethimide</p> <p><u>Progestins</u></p> <p>hydroxyprogesterone</p> <p>caproate</p> <p>medroxyprogesterone</p> <p>acetate</p> <p>megestrol acetate</p> <p><u>Estrogens</u></p> <p>diethylstilbestrol</p> <p>ethynyl estradiol/</p> <p>equivalents</p> <p><u>Anti estrogen</u></p> <p>tamoxifen</p> <p><u>Androgens</u></p> <p>testosterone propionate</p> <p>fluoxymesterone/equiv-</p> <p>alents</p> <p><u>Antiandrogens</u></p> <p>flutamide</p> <p>gonadotropin-releasing</p> <p>hormone analogs</p> <p>leuprolide</p> <p><u>Nonsteroidal</u></p> <p><u>antiandrogens</u></p> <p>flutamide</p> <p><u>Photosensitizers</u></p> <p>hematoporphyrin</p> <p>derivatives</p> <p>Photofrin®</p> <p>benzoporphyrin</p> <p>derivatives</p> <p>Npe6</p> <p>tin etioporphyrin (SnET2)</p> <p>pheoboride-a</p> <p>bacteriochlorophyll-a</p> <p>naphthalocyanines</p> <p>phthalocyanines</p> <p>zinc phthalocyanines</p>
--	---	--	--

Examples of chemotherapeutic agents that are particularly useful in conjunction with radio-

- 50 -

sensitizers include, for example, adriamycin, camptothecin, carboplatin, cisplatin, daunorubicin, doxorubicin, interferon (alpha, beta, gamma), interleukin 2, irinotecan, docetaxel, paclitaxel, topotecan, and therapeutically effective analogs and derivatives of the same.

As appreciated by persons skilled in the art, reference herein to treatment extends to prophylaxis, as well as to treatment of established diseases or symptoms. It is further appreciated that the amount of a compound of the invention required for use in treatment varies with the nature of the condition being treated, and with the age and the condition of the patient, and is ultimately determined by the attendant physician or veterinarian. In general, however, doses employed for adult human treatment typically are in the range of 0.001 mg/kg to about 100 mg/kg per day. The desired dose can be conveniently administered in a single dose, or as multiple doses administered at appropriate intervals, for example as two, three, four or more subdoses per day. In practice, the physician determines the actual dosing regimen most suitable for an individual patient, and the dosage varies with the age, weight, and response of the particular patient. The above dosages are exemplary of the average case, but there can be individual instances in which higher or lower dosages are merited, and such are within the scope of the present invention.

Formulations of the present invention can be administered in a standard manner for the treatment of the indicated diseases, such as orally, parenterally, transmucosally (e.g., sublingually or via buccal administration), topically, transdermally, rectally, via inhalation (e.g., nasal or deep

lung inhalation). Parenteral administration includes, but is not limited to intravenous, intra-arterial, intraperitoneal, subcutaneous, intramuscular, intrathecal, and intraarticular. Parenteral administration also can be accomplished using a high pressure technique, like POWDERJECT™.

For oral administration, including buccal administration, the composition can be in the form of tablets or lozenges formulated in conventional manner. For example, tablets and capsules for oral administration can contain conventional excipients such as binding agents (for example, syrup, acacia, gelatin, sorbitol, tragacanth, mucilage of starch, or polyvinylpyrrolidone), fillers (for example, lactose, sugar, microcrystalline cellulose, maize-starch, calcium phosphate, or sorbitol), lubricants (for example, magnesium stearate, stearic acid, talc, polyethylene glycol or silica), disintegrants (for example, potato starch or sodium starch glycolate), or wetting agents (for example, sodium lauryl sulfate). The tablets can be coated according to methods well known in the art.

Alternatively, the compounds of the present invention can be incorporated into oral liquid preparations such as aqueous or oily suspensions, solutions, emulsions, syrups, or elixirs, for example. Moreover, formulations containing these compounds can be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparations can contain conventional additives, for example suspending agents, such as sorbitol syrup, methyl cellulose, glucose/sugar syrup, gelatin, hydroxyethylcellulose, hydroxypropylmethylcellulose, carboxymethylcellulose, aluminum stearate gel, and hydrogenated edible

5

10

15

25

35

jection. Accordingly, the compounds of the invention can be formulated with suitable polymeric or hydrophobic materials (e.g., an emulsion in an acceptable oil), ion exchange resins, or as sparingly soluble derivatives (e.g., a sparingly soluble salt).

For veterinary use, a compound of formula (I), (II), or (III), or a nontoxic salt thereof, is administered as a suitably acceptable formulation in accordance with normal veterinary practice. The veterinarian can readily determine the dosing regimen and route of administration that is most appropriate for a particular animal.


Thus, the invention provides a pharmaceutical composition comprising a compound of the formula (I), (II), or (III), together with a pharmaceutically acceptable diluent or carrier therefor. Also provided is a process of preparing a pharmaceutical composition comprising a compound of formula (I), (II), or (III) comprising mixing a compound of formula (I), (II), or (III), together with a pharmaceutically acceptable diluent or carrier therefor.

Specific, nonlimiting examples of compounds of structural formula (I), (II), and (III) are provided below, the synthesis of which were performed in accordance with the procedures set forth below.

For ease of understanding, a compound having a particular structure is identified by the corresponding compound number provided in the following tables summarizing some of the compounds useful in the method. For example, the structure identified as Compound 1 is a compound of structural formula (IV), wherein R^{27} is hydrogen and R^{28} is $-C(O)NH(CH_2)_2(2-N\text{-methylpyrrolidyl})$.

- 54 -








Compounds suitable in the method include,
but are not limited to:











10









Compound d No.	R ²⁷	R ²⁸
1	H	
2	H	
3	H	
4	H	
5	H	
6	H	

15

Compound No.	R ²⁷	R ²⁸
7	H	
8	H	
9	H	
10		H
11		H
12	CH ₃	H
13		H
14	NH ₂	H
15		H









Compound No.	R ²⁷	R ²⁸
16		H
17		H
18		H
19	H	
20		H
21	Cl	
22		H
23	Cl	H
24	H	









- 57 -

Compound No.	R ²⁷	R ²⁸
25	H	
26		H
27	H	
28	H	
29	H	Cl
30		H
31		H
32	H	NH ₂
33	H	
34	H	









5

10







Compound No.	R ²⁷	R ²⁸
35	H	
36		H
37		H
38	H	H
39	H	
40	H	
41	H	
42		H
43	H	








Compound No.	R ²⁷	R ²⁸
44	H	
45		H
46	H	
47	H	
48		H
49		H
50		H
51		H








- 60 -









Compound No.	R ²⁷	R ²⁸
52	H	
53	H	
54		H
55		H
56		H
57	H	
58	H	
59		H

- 61 -


Compound No.	R ²⁷	R ²⁸
60	H	
61	H	
62		H
63		H
64		H
65	H	

Compound No.	R ²⁷	R ²⁸
66	H	
67	H	
68	H	
69	H	
70	H	
71	H	
72	H	


Compound No.	R ²⁷	R ²⁸
73	H	
74	H	
75	H	
76	H	
77		H
78		H
79		H

Compound No.	R ²⁷	R ²⁸
80		H
81		H
82		H
83		H
84		H
85		H
86		H
87		H





- 65 -

Compound No.	R ²⁷	R ²⁸
88		H

Hetero Ring Substitutions:











10







Compound No.	R ²⁹
89	
90	
91	
92	

15

- 66 -

Compound No.	R ²⁹
93	 A pyrimidine ring with a wavy line representing a substituent at the 2-position.
94	 An imidazole ring with a wavy line representing a substituent at the 2-position.
95	 A pyrimidin-4(1H)-one ring with a wavy line representing a substituent at the 2-position.
96	 A 1,2,4-triazin-3(1H)-one ring with a wavy line representing a substituent at the 2-position.
97	 A 1,3,5-triazine ring with a wavy line representing a substituent at the 2-position.
98	 A pyridine ring with a wavy line representing a substituent at the 2-position.
99	 A 1,2,4-triazole ring with a wavy line representing a substituent at the 2-position.
100	 A pyrimidine ring with bromine atoms at the 2 and 6 positions and a wavy line representing a substituent at the 4-position.

- 67 -




Compound No.	R ²⁹
101	
102	
103	
104	
105	
106	

Thiourea Compounds:




10





15















Compound No.	R ³⁰	R ³¹
112		Cl
113		H
114		Cl











5 Miscellaneous Class:















10











Compound No.	R ³²	Ar ¹	Ar ²
115	-N(H) -		
116	-N(H) -		













Compound No.	R ³²	Ar ¹	Ar ²
117	-N(H) -		
118	-N(CH ₃) -		
119	-N(H) -		
120	-N(H) -		
121	-N(H) -		
122	-N(H) -		
123	-N(H) -		













Compound No.	R ³²	Ar ¹	Ar ²
131	-N(H)-		
132	null		
133	-N(H)-		
134	-N(H)-		
135	-O-		



- 73 -

Compound No.	R ³²	Ar ¹	Ar ²
136	-N(H)-		
137	-N(H)-		
138	-N(H)-		
139	-N(H)-		
140	-O-		
141	-O-		

Compound No.	R ³²	Ar ¹	Ar ²
142	-N(H) -		
143	-O-		
144	-N(H) -		
145	-N(H) -		
146	-N(H) -		

Compound No.	R ³²	Ar ¹	Ar ²
147	-N(H)-		
148	-N(H)-		
149	-O-		
150	-N(H)-		
151	-N(H)-		
152	-N(H)-		

Compound No.	R ³²	Ar ¹	Ar ²
159	-N(H)-		
160	-N(H)-		
161	-N(H)-		
162	-N(H)-		
163	-N(H)-		
164	-N(H)-		

Compound No.	R ¹²	Ar ¹	Ar ²
165	-N(H)-		

Some preferred compounds include Compound Nos. 2, 4, 6, 12, 72, 76, 83, 84, 88, 89, and 90.

Generally, compounds of structural formulae (I), (II), and (III), including those of formulae (IV), (V), (VI), and (VII), can be prepared according to the following synthetic scheme. In the scheme described below, it is understood in the art that protecting groups can be employed where necessary in accordance with general principles of synthetic chemistry. These protecting groups are removed in the final steps of the synthesis under basic, acidic, or hydrogenolytic conditions which are readily apparent to those skilled in the art. By employing appropriate manipulation and protection of any chemical functionalities, synthesis of compounds of structural formulae (I), (II), and (III) not specifically set forth herein can be accomplished by methods analogous to the schemes set forth below.


Unless otherwise noted, all starting materials were obtained from commercial suppliers and used without further purification. All reactions and chromatography fractions were analyzed by thin-layer chromatography on 250-mm silica gel plates, visualized with UV (ultraviolet) light and I₂

- 79 -

(iodine) stain. Flash column chromatography was carried out using Biotage 40M silica gel (230-400 mesh). Products and intermediates were purified by flash chromatography or reverse-phase HPLC.

As illustrated below, the compounds of general structural formulae (I) and (II) can be prepared by the following general synthetic schemes.

General Scheme 1



In general, an aryl amine represented by the formula Ar-NH_2 is reacted with about 0.75-1.25 molar equivalent of 4-nitrophenyl chloroformate. The reaction preferably is performed under an inert atmosphere, for example, nitrogen (N_2), and typically is maintained at low temperature (about 0°C). The resulting product is treated with about 0.75-1.25 molar equivalent of a heteroaryl amine represented by the formula HetAr-NH_2 , preferably under an inert atmosphere at room temperature (about 25°C), to afford a crude aryl pyrazine-disubstituted urea compound.

5

10

15

The reaction scheme illustrates the synthesis of 2-((2-methoxy-4-((R)-amino)benzoyl)amino)pyridine through five steps:

- Step (1):** Esterification of 2-amino-4-methoxybenzoic acid with methyl ester using TMSCHN_2 to form methyl 2-amino-4-methoxybenzoate.
- Step (2):** Coupling of methyl 2-amino-4-methoxybenzoate with 4-nitrobenzoyl chloride ($\text{O}_2\text{N}-\text{C}_6\text{H}_4-\text{OCOCl}$) to form the dimeric intermediate methyl 2-((4-nitrobenzoyl)amino)-4-methoxybenzoate.
- Step (3):** Coupling of the dimeric intermediate with 2-aminopyridine to form the pyridine-containing intermediate methyl 2-((2-((4-nitrobenzoyl)amino)-4-methoxybenzoyl)amino)pyridine-4-carboxylate.
- Step (4):** Hydrolysis of the methyl ester group to form the carboxylic acid intermediate 2-((2-((4-nitrobenzoyl)amino)-4-methoxybenzoyl)amino)pyridine-4-carboxylic acid.
- Step (5):** Coupling of the carboxylic acid intermediate with an R-NH_2 group to form the final product, 2-((2-methoxy-4-((R)-amino)benzoyl)amino)pyridine.

vacuo to provide the desired ester as an off-white solid (94% yield).

Step (2): p-Nitrophenyl Carbamate Procedure

5

10

15

20

To a stirred, cooled (about 0°C) solution of methyl-3-amino-4-methoxy benzoate (5.0 g; 27.6 mmol) in dry dichloromethane (175 mL) was added pyridine (2.34 mL; 29 mmol) followed by 4-nitrophenyl chloroformate (5.8 g; 29 mmol) under a nitrogen (N₂) atmosphere. After stirring for 8 hours, the reaction was washed with 2N aqueous hydrochloric acid (2 X 200 mL), saturated aqueous sodium bicarbonate (2 X 200 mL), and brine (200 mL), then dried (MgSO₄), and filtered. The filtered solution was diluted with ethyl acetate and hexanes (about 800 mL) until a precipitate formed. The solid was collected on a Buchner funnel with suction, and air dried to provide the desired carbamate as a white solid (70% yield).

Step (3): Carbamate Coupling Procedure

25

30

To a stirred solution of 4-methoxy-3-(4-nitro-phenoxy-carbonylamino)-benzoic acid methyl ester (30 g; 8.7 mmol) in dry N-methyl pyrrolidine (50 mL) was added the amino pyrazine (0.84 g; 8.8 mmol) under a N₂ atmosphere at room temperature. The reaction mixture was heated to 80°C for 6 hours, then allowed to cool to room temperature. Dilution with ethyl acetate (200 mL) and water (200 mL) provided the desired urea as a white solid (54% yield).

Step (4): Lithium Hydroxide Hydrolysis Procedure

To a stirred solution of 4-methoxy-4-(3-pyrazin-2-yl-ureido)-benzoic acid methyl ester (1.0 g; 3.3 mmol) in methanol (35 mL) was added aqueous lithium hydroxide (5 mL of a 2N solution; 10 mmol) at room temperature. The reaction was heated to 67°C for 15 hours, then allowed to cool to room temperature. The reaction then was diluted with water (100 mL), and washed with ethyl acetate (2 X 100 mL). The pH of the aqueous layer was adjusted to pH 5.2 with 2N aqueous hydrochloric acid, and the resulting precipitate was collected on a Buchner funnel with suction and air-dried to provide the desired acid as a white solid.





Step (5): HBTU Coupling Procedure




To a stirred solution of the acid (30 mg; 0.11 mmol) in dry N-methyl pyrrolidinone (2 mL) was added O-benzotrazol-1-yl-N,N, N',N'-tetramethyluronium hexafluorophosphate (HBTU; 45 mg; 0.12 mmol), 4-(2-aminioethyl)-morpholine (15.7 L; 0.12 mmol) and diisopropyl ethyl amine (34 L, 0.2 mmol) at room temperature under a nitrogen atmosphere. The resulting solution was stirred 5 hours, then diluted with ethyl acetate (30 mL) and 10% aqueous sodium carbonate (30 mL). After stirring vigorously at room temperature for 15 minutes, the precipitate was collected on a Buchner funnel with suction and air-dried to provide the desired amide as a white solid (59% yield).

The following compounds were prepared using the general procedure described accompanying


General Scheme 2, but substituting the R group below for the R group shown in General Scheme 2:

5

Compound No.	R group	Characterization
22		¹ H NMR (400 MHz, D ₆ DMSO) δ 10.15 (br s, 1H), 8.90 (s, 1H), 8.69 (d, 1H), 8.53 (t, 1H), 8.27 (t, 1H), 8.18 (d, 1H), 7.53 (dd, 1H), 3.96 (br s, 2H), 3.93 (s, 3H), 3.67-3.53 (m, 6H), 3.29 (t, 2H), 3.10 (t, 2H). LRMS (esi, positive) m/e 401.2 (M+1).
11		¹ H NMR (400 MHz, D ₆ DMSO) δ 10.14 (br s, 1H), 8.91 (s, 1H), 8.66 (d, 1H), 8.27 (dd, 1H), 8.18 (d, 1H), 7.51 (dd, 1H), 7.03 (d, 1H), 3.91 (s, 3H), 3.53 (br s, 1H), 3.33 (d, 2H), 3.21 (br s, 1H), 3.02 (br s, 1H), 2.78 (d, 2H), 2.48 (s, 3H), 2.31 (br s, 1H), 2.13 (br s, 1H), 1.99-1.82 (m, 2H). LRMS (esi, positive) m/e 399.2 (M+1).
25		¹ H NMR (400 MHz, D ₆ DMSO) δ 10.20 (br s, 1H), 8.87 (s, 1H), 8.31-8.26 (m, 2H), 8.21 (br s, 1H), 7.52-7.44 (m, 2H), 3.95 (s, 3H), 3.72 (t, 2H), 3.56 (t, 3H).
28		¹ H NMR (400 MHz, D ₆ DMSO) δ 10.22 (br s, 1H), 8.90 (br s, 1H), 8.64 (t, 1H), 8.31-8.28 (m, 2H), 8.19 (d, 1H), 7.51-7.47 (m, 2H), 3.97 (br s, 2H), 3.93 (s, 3H), 3.69-3.52 (m, 6H), 3.30 (t, 2H), 3.11 (t, 2H). LRMS (esi, positive) m/e 401.1 (M+1).

Compound No.	R group	Characterization
1		¹ H NMR (400 MHz, D ₆ DMSO) δ 10.20 (br s, 1H), 8.88 (br s, 1H), 8.40 (t, 1H), 8.32 (t, 1H), 8.23 (dd, 2H), 7.47 (d, 1H), 7.43 (dd, 1H), 3.94 (s, 3H), 2.91 (c, 2H), 2.19 (s, 3H), 2.05-1.98 (m, 2H), 1.93-1.80 (m, 2H), 1.60 (c, 1H), 1.43 (c, 1H). LRMS (esi, positive) m/e 399.0 (M+1).
35		¹ H NMR (400 MHz, D ₆ DMSO) δ 10.21 (br s, 1H), 8.88 (br s, 1H), 8.48 (br s, 1H), 8.32 (t, 1H), 8.26 (s, 1H), 8.23 (d, 1H), 7.50 (br s, 1H), 7.46 (dd, 1H), 3.95 (s, 3H), 3.36 (c, 2H), 3.10 (t, 2H), 2.83 (s, 3H). LRMS (esi, positive) m/e 409.0 (M+1).
44		¹ H NMR (400 MHz, D ₆ DMSO) δ 11.47 (br s, 1H), 8.48 (br s, 1H), 8.41 (s, 1H), 8.35 (d, 1H), 8.25-8.23 (M, 2H), 7.06-7.02 (m, 2H), 3.98 (s, 3H), 3.38 (br s, 2H), 3.04 (br s, 3H), 1.21 (br s, 3H). LRMS (esi, positive) m/e 330.0 (M+1).




General Scheme 3





Isocyanate Procedure:

To a stirred solution of 2-methoxy-5-methyl-phenylisocyanate (43 mL; 0.3 mmol) in dry dichloroethane (0.4 mL) was added 2-aminoquinoxaline (43.5 mg; 0.3 mmol) in a reaction vial under a nitrogen atmosphere. The vial was capped and heated to 80°C overnight (14 hours). The reaction mixture then was filtered, and the residue washed with dichloromethane to provide the desired urea as a white solid (91% yield).

The following compounds were prepared using the procedure described accompanying General Scheme 3, but substituting the Ar group in the table below for the Ar group in General Scheme 3:


Compound No.	Ar group	Characterization
90		¹ H NMR (400 MHz, D ₆ DMSO) d 11.65 (br s, 1H), 10.58 (br s, 1H), 10.58 (br s, 1H), 8.80 (s, 1H), 8.15 (d, 1H), 7.98 (d, 1H), 7.84-7.80 (m, 2H), 7.65 (c, 1H), 6.98 (d, 1H), 6.98 (d, 1H), 6.84 (dd, 1H), 3.99 (s, 3H), 2.25 (s, 3H). LRMS (esi, positive) m/e 309.1 (M+1).
98		¹ H NMR (400 MHz, D ₆ DMSO) d 9.61 (s, 1H), 8.06 (d, 1H), 7.35 (d, 1H), 6.88 (dd, 1H), 8.87 (d, 1H), 5.42 (d, 1H), 3.89 (s, 3H), 2.33 (s, 3H). LRMS (esi, positive) m/e 259.0 (M+1).
97		¹ H NMR (400 MHz, D ₆ DMSO) d 11.29 (s, 1H), 10.64 (s, 1H), 9.05 (d, 1H), 8.70 (d, 1H), 8.05 (d, 1H), 6.92 (d, 1H), 6.81 (c, 1H), 3.85 (s, 3H), 2.24 (s, 3H). LRMS (esi, positive) m/e 259.0 (M+1).

Compound No.	Ar group	Characterization
91		¹ H NMR (400 MHz, D ₆ CDCl ₃) δ: 11.12 (br s, 1H), 8.61 (br s, 1H), 8.28 (s, 1H), 8.24 (s, 1H), 8.14 (d, 1H), 8.04-8.01 (m, 2H), 7.55-7.50 (m, 3H), 6.86 (dd, 1H), 6.76 (d, 1H), 3.52 (s, 3H), 2.35 (s, 3H). LRMS (esi, positive) m/e 335.2 (M+1).
12		¹ H NMR (300 Mhz, d ₆ -DMSO) δ: 10.10 (s, 1H), 10.00 (br s, 1H), 8.90 (s, 1H), 8.32 (s, 1H), 8.23 (s, 1H), 8.04 (s, 1H), 7.93 (d, 1H), 6.81 (d, 1H), 3.87 (s, 3H), 2.28 (s, 3H). ¹³ C NMR (75 Mhz, d ₆ -DMSO) δ: 151.5, 149.3, 146.0, 140.9, 137.2, 135.2, 129.2, 127.7, 122.7, 119.5, 110.8, 55.9, 20.5

EXAMPLES

Example 1

Preparation of Compound 115




1-[2-(1,1-difluoromethoxy)-phenyl]-3-pyrazin-2-yl-urea

2-(Difluoromethoxy)phenylisocyanate (1.0 g, 5.4 mmol) and aminopyrazine (0.51g, 5.4 mmol) were reacted for 6 hours in refluxing dimethoxyethane (20 mL). The reaction mixture was cooled to room temperature to precipitate the product, which was collected by filtration, washed with ethyl

acetate, and dried *in vacuo* (765 mg, 50%). ^1H NMR (300 Mhz, d_6 -DMSO) δ : 10.49 (br s, 1H), 10.26 (s, 1H), 8.83 (s, 1 H), 8.35-8.24 (m, 3H), 7.53-7.00 (m, 4H).

Example 2

Preparation of Compound 165




1-(2-methylsulfanylphenyl)-3-pyrazin-2-yl urea

2-(Methylthiophenyl)-isocyanate (1.0 g, 6.1 mmol) and aminopyrazine (0.58 g, 6.1 mmol) were reacted for 16 hours in refluxing dimethoxyethane (40 mL). The product precipitated from the cooled reaction mixture and was collected by filtration, washed with dimethoxyethane, and dried *in vacuo* (715 mg, 45%). ^1H NMR (300 MHz, d_6 -DMSO) δ : 10.35 (br s, 1H), 10.29 (s, 1H), 8.84 (s, 1H), 8.33 (s, 1H), 8.27 (s, 1H), 8.09 (d, 1H), 7.45 (d, 1H), 7.29, (t, 1H), 7.10 (t, 1H), 2.43 (s, 3H). ^{13}C NMR (75 Mhz, d_6 -DMSO) δ : 151.8, 149.2, 140.5, 137.5, 137.3, 135.2, 130.1, 127.1, 126.9, 123.7, 121.4, 16.5.

Example 3

Preparation of Compound 159




1-(2-methoxy-5-nitrophenyl)-3-pyrazin-2-yl-urea

A mixture of 2-methoxy-5-nitrophenyl isocyanate (5.0 g, 25 mmol) and aminopyrazine (2.5 g, 26 mmol) in tetrahydrofuran (THF, 250 mL) was stirred at reflux for 24 hours. The product was precipitated from the cooled reaction mixture and was collected by filtration, washed with ethyl acetate, and dried in vacuo (4.3 g, 57%). ¹H NMR (300 MHz, d₆-DMSO) (mixture of rotamers) δ: 10.38 (br, s, 1H), 10.27 (s, 1H), 9.39, 8.88 (2 singlets, 1H), 9.10(d, 1H), 8.33 (s, 1H), 8.26 (d, 1H), 7.98-8.25 (m, 1H), 7.97-7.84 (m, 1H), 4.05, 4.03 (2 singlets, 77:28 ratio, 3H).

- 89 -

Example 4

Preparation of Compound 14



1-(5-amino-2-methoxyphenyl)-3-pyrazin-2-yl-urea

A solution of (2-methoxy-5-nitrophenyl)-3-pyrazin-2-yl-urea (Compound 159, Example 3) (16.9 g, 55 mmol) in dimethylformamide (DMF, 320 mL) was shaken under H₂ in the presence of palladium on carbon (Pd/C) catalyst (1.6 g, 10% Pd) at 80°C for 12 h. A second portion of catalyst was added (1.6 g) and shaking was continued for an additional 8 h at the same temperature. The solution was filtered through a pad of celite using an additional 200 mL of DMF. The filtrate was concentrated *in vacuo* and the residue was triturated with methanol (100 mL). The solid was collected, stirred in boiling methanol, and solids present (1.8 g) were filtered off and discarded. The filtrate was cooled at 4°C overnight. Solids (1.4 g) were removed by filtration and the filtrate was concentrated *in vacuo* to a tan solid (2.6 g). The crude solid product was triturated with THF (200 mL), collected by filtration, and dried *in vacuo* to afford the product as a tan solid (1.85 g, 13%). ¹H NMR (300 Mhz, d₆ DMSO) δ: 10.10 (s, 1H), 9.94 (br s, 1H), 8.89 (s, 1H), 8.32 (s, 1H), 8.22 (s, 1H), 7.58 (s, 1H), 7.75 (d, 1H), 6.21 (d, 1 H), 4.70 (s, 2H), 3.76 (s, 3H). ¹³C

- 90 -


NMR (75 Mhz, d_6 -DMSO) δ : 151.4, 149.4, 142.6, 140.9, 139.8, 137.2, 135.2, 128.7, 112.8, 107.7, 106.0, 56.8.

5

Example 5

Preparation of Compound 48

10



15

N-[4-methoxy-3-(3-pyrazin-2-yl-ureido)phenyl]-succinamic acid

20


25

30

A solution of 1-(5-amino-2-methoxyphenyl)-3-pyrazin-2-yl-urea (Compound 14, Example 4) (260 mg, 1 mmol) and succinic anhydride (131 mg, 1.3 mmol) in dry pyridine (10 mL) was stirred 16 h at room temperature. The resulting solid was collected by filtration and triturated with chloroform, and dried *in vacuo* to afford the off-white product (175 mg, 50%). ^1H NMR (300 Mhz, d_6 -DMSO) δ : 10.15 (s, 1H), 10.05 (s, 1H), 9.87 (s, 1H), 8.90 (s, 1H), 8.33 (s, 2H), 8.24 (s, 1H), 7.42 (dd, J = 8.8, 2.2 Hz, 1H), 6.96 (d, J = 8.8 Hz, 1H), 3.87 (s, 3H), 2.54 (br s, 4H).

Example 6

Preparation of Compound 36




(S)-1-(2,2,2-trifluoroethanoyl)pyrrolidine-2-carboxylic acid [4-methoxy-3-(3-pyrazin-2-ylureido)phenyl]-amide

A solution of 1-(5-amino-methoxyphenyl)-3-pyrazin-2-yl-urea (Compound 14, Example 4) (105 mg, 0.4 mmol) in dry pyridine (2 mL) at 0°C was treated with a solution of N-trifluoroacetyl-(S)-prolyl chloride (0.1 M in dichloromethane, 4.5 mL, 0.45 mmol) and stirred 2 h at room temperature. The reaction was quenched with 1 N HCl (50 mL) and extracted with ethyl acetate (3 X 50 mL). The combined organic layers were washed with 1 N HCl (2 X 20 mL), water (20 mL), brine (20 mL), dried over sodium sulfate, and concentrated *in vacuo* to a beige solid (60 mg). Recrystallization from acetonitrile yielded the final solid product (30 mg, 17%). ¹H NMR (300 Mhz, *d*₆-DMSO) δ: (10.16-10.06, m, 3H), 8.90 (s, 1H), 8.36-8.30 (m, 2H), 8.25 (d, J = 2.6 Hz, 1H), 7.42 (dd, J = 8.8, 2.6 Hz, 1H), 6.98 (d, J = 8.8 Hz, 1H), 4.57 (dd, J = 8.5, 4.4 Hz, 1 H), 3.88 (s, 3H), 3.73 (t, J = 6.5 Hz, 1H), 2.27-2.21 (m,

1H), 2.06-1.90 (m, 3H). LRMS (ESI, positive) m/e 453.1 (M + 1).

Example 7

Preparation of Compound 16




(S)-pyrrolidine-2-carboxylic acid [4-methoxy-3-(3-pyrazin-2-yl-ureido)-phenyl]-amide

A suspension of (S)-1-(2,2,2-trifluoroethanoyl)-pyrrolidine-2-carboxylic acid [4-methoxy-3-(3-pyrazin-2-yl-ureido)-phenyl]-amide (Compound 36, Example 6) (22 mg, 0.05 mmol) in a mixture of methanol (MeOH, 5 mL) and water (about 0.25 mL) was treated with KOH (100 mg, large excess). Within 10 minutes, all ingredients were in solution. The reaction mixture was treated with water (20 mL) and extracted with ethyl acetate (2 x 20 mL). The organic layers were combined and washed with water (10 mL) and brine (10 mL), dried with sodium sulfate, and concentrated to a tan solid (13 mg, 75%). ¹H NMR (300 MHz, d₆-DMSO) δ: 10.13 (s, 1H), 10.03 (s, 1H), 9.85 (s, 1H), 8.90 (s, 1H), 8.38-8.28 (m, 2H), 8.25 (d, J = 2.6 Hz, 1H), 7.41 (dd, J = 8.8, 2.6 Hz), 6.98 (d, J = 8.8 Hz, 1H), 3.88 (s, 3H, 3.69 (dd, J = 8.7, 5.6 Hz, 1H), 2.94-2.87 (m, 2H), 2.10-1.97 (m, 1H), 1.81-1.60 (m, 3H). LRMS (ESI, positive) m/e 357.1 (M + 1).

Example 8


Preparation of Compound 42



N-[4-methoxy-3-(3-pyrazin-2-yl-ureido)phenyl]-methanesulfonamide

A solution of 1-(5-amino-2-methoxyphenyl)-3-pyrazin-2-yl-urea (Compound 14, Example 4) (260 mg, 1 mmol) in dry pyridine (15 mL) was treated with methanesulfonyl chloride (0.08 mL, 1 mmol) and stirred 16 h at room temperature. The reaction mixture was concentrated *in vacuo* and the solid residue was triturated with ethanol, collected by filtration, and dried *in vacuo* to afford the product (205 mg, 61%). ¹H NMR (300 Mhz, *d*₆-DMSO) δ: 10.16 (s, 1H), 10.07 (s, 1H), 9.40 (s, 1H), 8.92 (s, 1H), 8.35 (s, 1H), 8.27 (s, 1H), 8.19 (d, *J* = 2.2 Hz, 1H), 7.04 (d, *J* = 8.8 Hz, 1H), 6.91 (dd, *J* = 8.7, 2.4 Hz, 1H), 3.91 (s, 3H), 2.91 (s, 3H).


Example 10
Preparation of Compound 32



1-(4-amino-2-methoxyphenyl)-3-pyrazin-2-yl-urea

A solution of (2-methoxy-4-nitrophenyl)-3-pyrazin-2-yl-urea (Compound 65, Example 9) (7.9 g, 27 mmol) in DMF (300 mL) was shaken under H₂ in the presence of Pd/C catalyst (1.6 g, 10% Pd) at 110°C for 4 h. The mixture was filtered through a pad of celite using an additional 200 mL of DMF. The filtrate was concentrated in vacuo and the residue was recrystallized from ethanol (with a hot filtration step) to yield the light gray product (2.9 g, 41%). ¹H NMR (300 MHz, d₆-DMSO) δ: 9.83 (s, 1H), 9.50 (s, 1H), 8.86 (s, 1H), 8.28 (s, 1H), 8.19 (d, J = 2.5 Hz, 1H), 7.64 (d, J = 8.5 Hz, 1H), 6.31 (d, J = 2.0 Hz, 1H), 6.13 (dd, J = 8.5, 2.0 Hz, 1H), 4.92 (s, 2H), 3.79 (s, 3H). ¹³C NMR (75 Mhz, d₆-DMSO) δ: 151.6, 150.0, 149.6, 145.2, 140.9, 136.9, 135.1, 121.6, 116.8, 105.5, 98.0, 55.4.


Example 11
Preparation of Compound 3



C-dimethylamino-N-[3-methoxy-4-(3-pyrazin-2-yl-ureido)phenyl]-acetamide

A solution of N,N-dimethylglycine (124 mg, 1.2 mmol) and triethylamine (0.33 mL, 2.4 mmol) in dry acetonitrile (5 mL) at 0°C was treated dropwise with isobutyl chloroformate (0.16 mL, 1.2 mmol) and stirred 15 min. This mixture was treated dropwise with a solution of 1-(4-amino-2-methoxyphenyl)-3-pyrazin-2-yl-urea (Compound 32, Example 10) (100 mg, 0.4 mmol) in dimethyl sulfoxide (DMSO, 1 mL). The reaction mixture was stirred at room temperature for 3 h, quenched with water (20 mL), and extracted with ethyl acetate (2 x 15 mL). The combined organic layers were washed with water (10 mL) and brine (10 mL), dried over sodium sulfate, and concentrated *in vacuo*. The residue was dissolved in DMSO (1 mL) and purified by HPLC (YMC 20 x 50 mm C18 CombiPrep column, 20 mL/min, 2-50% CH₃CN/water in 6 min, all solvents contained 0.05% trifluoroacetic acid (TFA), 0.35 mL injections, detector at 254 nm, detector path length 0.2 mm). Fractions containing the product were concentrated *in vacuo* to afford the product as the trifluoroacetate (TFA) salt (24 mg, 17%).


Example 12
Preparation of Compound 8



3-chloro-N-[3-methoxy-4-(3-pyrazin-2-yl-ureido)phenyl]-propionamide

A solution of 1-(4-amino-2-methoxyphenyl)-3-pyrazin-2-yl-urea (Compound 32, Example 10) (259 mg, 1 mmol) in pyridine (3 mL) at 0°C was treated with chloroacetyl chloride (0.29 mL, 3 mmol). The suspension was warmed at 80°C until most solids dissolved, the reaction mixture was cooled to room temperature and the product was precipitated with ether (10 mL). This crude product was used without purification for further reactions, but a portion (30 mg) was purified by HPLC (Luna 10 x 250 mm C18 column, 4.7 mL/min, 2-80% CH₃CN/water in 15 min, all solvents contained 0.05% TFA, 0.25 mL injections, detector at 254 nm, detector path length 0.3 mm). Fractions containing the product were concentrated *in vacuo* to afford the product. ¹H NMR (300 Mhz, d₆-DMSO) δ: 10.00 (s, 2H), 9.94 (s, 1H), 8.87 (s, 1H), 8.32 (dd, J = 2.5, 1.5 Hz, 1H), 8.23 (d, J = 2.6 Hz, 1H), 8.05 (d, J = 8.7 Hz, 1H), 7.49 (d, J = 2.1 Hz, 1H), 7.07 (dd, J = 8.7, 2.1 Hz, 1H), 3.90-3.80 (m, 5H), 2.80 (t, J = 6.2 Hz, 2H). LRMS (ESI, positive)-m/e 350, 352 (M + 1).

Example 13
Preparation of Compound 4




(cyclohexyl-methyl-amino)-N-[3-methoxy-4-(3-pyrazin-2-yl-ureido)phenyl]-propionamide

A mixture of 3-chloro-N-[3-methoxy-4-(3-pyrazin-2-yl-ureido)phenyl]-propionamide (Compound 8, Example 12) and N-cyclohexyl-methylamine (0.5 mL, large excess) was warmed at 80°C for 1 h and cooled to room temperature. Crude product was precipitated from ether (10 mL), collected by filtration, and dissolved in DMSO (0.5 mL). Aliquots (about 0.25 mL) were purified by HPLC (Luna 10 x 250 mm C18 column, 4.7 mL/min, 2-80% CH₃CN/water in 15 min, all solvents contained 0.05% trifluoroacetic acid, detector at 254 nm, detector path length 0.3 mm). Fractions containing the product were concentrated *in vacuo* to afford the product as the TFA salt (4.7 mg, 11%). ¹H NMR (300 Mhz, d₆-DMSO) δ: 10.18 (s, 1H), 10.07 (s, 1H), 9.98 (s, 1H), 9.04 (br s, 1H), 8.88 (s, 1H), 8.33 (dd, J = 2.6, 1.5 Hz, 1H), 8.24 (d, J = 2.6 Hz, 1H), 8.08 (d, J = 8.7 Hz, 1H), 7.43 (d, J = 2.1 Hz, 1H), 7.09 (dd, J = 8.7, 2.1 Hz, 1H), 3.90-3.83 (m, 1H), 3.30-3.16 (m, 2H), 2.81 (t, J = 6.7 Hz, 1H), 2.74 (d, J = 5.0 Hz, 2H), 2.03-1.89 (m, 2H), 1.89-1.75 (m, 2H), 1.70-1.53 (m, 1H), 1.46-1.10 (m, 5H). LRMS (ESU, positive) m/e 427.2 (M + 1).

Example 14

Preparation of Compound 2

5



10

3-cyclopentylamino-N-[3-methoxy-4-(3-pyrazin-2-yl-ureido)phenyl]-propionamide


15

20

25

A mixture of 3-chloro-N-[3-methoxy-4-(3-pyrazin-2-yl-ureido)phenyl]-propionamide (Compound 8, Example 12) and cyclopentylamine (0.5 mL, large excess) was warmed at 80°C for 1 h and cooled to room temperature. The product was precipitated from ether (10 mL), collected by filtration, washed with ether, and dried *in vacuo* (24 mg, 60%). ¹H NMR (300 Mhz, d₆-DMSO) δ: 10.14 (s, 1H), 10.03 (s, 1H), 9.93 (s, 1H), 8.87 (d, J = 1.1Hz, 1H), 8.31 (dd, J = 2.6, 1.5 Hz, 1H), 8.23 (d, J = 2.7 Hz, 1H), 8.03(d, J = 8.7 Hz, 1H), 7.47(d, J = 2.1 Hz, 1H), 7.04(dd, J = 8.7, 2.1 Hz, 1H), 3.87 (s, 3H), 3.05 (quintet, J = 6.3 Hz, 1H), 2.80 (t, J = 6.6 Hz, 2H), 2.44 (t, J = 6.6 Hz, 2H), 1.80-1.67 (m, 2H), 1.65-1.56 (m, 2H), 1.53-1.42 (m, 2H), 1.37-1.27 (m, 2H). LRMS (ESI, positive) m/e 399.1 (M + 1).

Compound 166:



3-Methoxy-4-(3-pyrazin-2-yl-ureido)-benzoic acid

Step 1: Methyl-3-amino-4-methoxy benzoate. To a cooled (about 0 °C), stirred solution of 4-amino-3-methoxybenzoic acid (5.0 g, 30 mmol) in dry methanol (150 mL) was added trimethylsilyldiazomethane (60 mL of 2M solution in hexanes, 120 mmol) slowly over 1 hour. After stirring for 4 hours, the reaction was concentrated at reduced pressure, dissolved in ethyl acetate (200 mL), washed with 10% aqueous sodium carbonate and brine, then dried (MgSO₄), filtered, and concentrated in vacuo to provide the desired ester as an off-white solid (94% yield).


Step 2: 4-Methoxy-3-(4-nitro-phenoxy-carbonylamino)-benzoic acid methyl ester. To a stirred, cooled (about 0 °C) solution of methyl-3-amino-4-methoxy benzoate (5.0 g, 27.6 mmol) in dry dichloromethane (90 mL) was added pyridine (2.34 mL, 29 mmol) followed by 4-nitrophenyl chloroformate (5.8 g, 29 mmol) under a nitrogen atmosphere. After stirring for 1 hour, the reaction was diluted to 200 mL with dichloromethane and washed with 2N aqueous hydrochloric acid (2 x 200 mL), saturated aqueous sodium bicarbonate (2 x 200 mL), and brine (200 mL), then dried (MgSO₄), and filtered. The filtered solution was concentrated to a white solid corresponding to the desired carbamate (98% yield).

Step 3: 3-Methoxy-4-(3-pyrazin-2-yl-ureido)-benzoic acid methyl ester. To a stirred solution of 4-methoxy-3-(4-nitro-phenoxy-carbonylamino)-benzoic acid methyl ester (10.64 g, 30.7 mmol) in dry N-methyl pyrrolidinone (31 mL) at room temperature under nitrogen was added aminopyrazine (2.92 g, 30.7 mmol) and the reaction was warmed to 85°C. After 6 hours, the reaction was cooled to room temperature and triturated with ethyl acetate (200 mL). The precipitate formed was filtered off, rinsed with ethyl acetate and dried to give the urea as a tan solid (66 % yield).

Step 4: 3-Methoxy-4-(3-pyrazin-2-yl-ureido)-benzoic acid. To a stirred suspension of 3-methoxy-4-(3-pyrazin-2-yl-ureido)-benzoic acid methyl ester (6.07 g, 20 mmol) in 200 mL 3:1 MeOH:H₂O at room temperature under nitrogen was added lithium hydroxide monohydrate (8.4 g, 200 mmol) and the reaction heated to 65 °C overnight. The reaction was then cooled to room temperature and most of the methanol removed by rotary evaporation. The remaining suspension was neutralized to pH about 4 with concentrated HCl. The formed precipitate was isolated by filtration and rinsing with H₂O and then drying under high vacuum to give the desired acid as a white solid (5.34 g, 93%).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 8.89 (br s, 1H), 8.34 (s, 1H), 8.24 (s, 1H), 8.16 (d, 1H), 7.56 (s, 1H), 7.50 (d, 1H), 3.91 (s, 3H)

Compound 167:




N-Butyl-3-methoxy-4-(3-pyrazin-2-yl-ureido)-benzamide

To a stirred suspension of Compound 1xx (32 mg, 0.11 mmol) in 1 mL of NMP at room temperature in a capped reaction vial was added HBTU (0.4 M in NMP, 300 μ L, 0.12 mmol) and the suspension stirred for 15 minutes. N-Butyl amine (0.4 M in NMP, 300 μ L, 0.12 mmol) was then added followed by DIEA (38 μ L, 0.22 mmol). The reaction was stirred at room temperature overnight and was then diluted with EtOAc (20 mL) and 10% Na_2CO_3 (20 mL) and stirred rapidly for 5 minutes. A precipitate formed which was isolated by filtration and rinsing with H_2O and EtOAc. After air drying, the amide was isolated as an off-white solid (12.2 mg, 32 %).

$^1\text{H-NMR}$ (400 MHz, d_6 -DMSO) δ 8.92 (br s, 1H), 8.37 (br s, 2H), 8.23 (d, 1H), 8.21 (s, 1H), 7.53 (s, 1H), 7.47 (d, 1H), 3.97 (s, 3H), 3.23 (q, 2H), 1.52 (m, 2H), 1.35 (m, 2H), 0.92 (t, 3H)

LRMS (apci, positive) m/e 344.1 (M+1)

Compound 168:




N-Benzyl-3-methoxy-4-(3-pyrazin-2-yl-ureido)-benzamide

Prepared according to the procedure of Compound 167 using benzyl amine (39% yield).

$^1\text{H-NMR}$ (400 MHz, d_6 -DMSO) δ 8.96 (t, 1H), 8.91 (s, 1H), 8.56 (s, 1H), 8.50 (s, 1H), 8.44 (m, 1H), 7.58 (s, 1H), 7.56 (d, 1H), 7.35 (m, 4H), 7.23 (m, 1H), 4.46 (d, 2H), 3.97, (s, 3H)

LRMS (apci, positive) m/e 378.1 (M+1)

Compound 169:




3-Methoxy-N-phenethyl-4-(3-pyrazin-2-yl-ureido)-benzamide

Prepared according to the procedure of Compound 167 using phenethyl amine (49% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 8.92 (br s, 1H), 8.51 (t, 1H), 8.36 (br s, 1H), 8.25 (s, 1H), 8.22 (d, 1H), 7.52 (s, 1H), 7.45 (d, 1H), 7.36-7.20 (m, 5H), 3.98 (s, 3H), 3.46 (m, 2H), 2.84 (dd, 2H)

LRMS (apci, positive) m/e 392.1 (M+1)

Compound 170:




3-Methoxy-N-(3-phenyl-propyl)-4-(3-pyrazin-2-yl-ureido)-benzamide

Prepared according to the procedure of Compound 167 using phenpropyl amine (71% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 8.92 (br s, 1H), 8.40 (t, 1H), 8.36 (br s, 1H), 8.25 (d, 1H), 8.23 (s, 1H), 7.52 (s, 1H), 7.46 (d, 1H), 7.32-7.18 (m, 5H), 3.97 (s, 3H), 3.25 (m, 2H), 2.61 (m, 2H), 1.82 (m, 2H)

LRMS (apci, positive) m/e 406.1 (M+1)

Compound 171:




N-(2-Benzenesulfonyl-ethyl)-3-methoxy-4-(3-pyrazin-2-yl-ureido)-benzamide

Prepared according to the procedure of Compound 167 using 2-benzenesulfonyl-ethylamine (57% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 8.90 (br s, 1H), 8.43 (br m, 1H), 8.35 (br s, 1H), 8.25 (s, 1H), 8.22 (d, 1H), 7.96 (s, 1H), 7.93 (s, 1H), 7.72 (m, 1H), 7.63 (m, 2H), 7.38 (s, 1H), 7.33 (d, 1H), 3.95 (s, 3H), 3.59 (m, 2H), 3.55 (m, 2H)

LRMS (apci, positive) m/e 456.0 (M+1)

Compound 172:




N-(4-Iodo-benzyl)-3-methoxy-4-(3-pyrazin-2-yl-ureido)-benzamide

Prepared according to the procedure of Compound 167 using 4-iodo benzyl amine (66% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 8.97 (t, 1H), 8.90 (s, 1H), 8.36 (s, 1H), 8.26 (d, 1H), 8.24 (s, 1H), 7.69 (d, 2H), 7.56 (m, 2H), 7.16 (d, 2H), 4.41 (d, 2H), 3.97 (s, 3H)

LRMS (apci, positive) *m/e* 504.0 (M+1)

Compound 173:




3-Methoxy-4-(3-pyrazin-2-yl-ureido)-N-(2-pyridin-2-yl-ethyl)-benzamide

Prepared according to the procedure of Compound 167 using 2-pyridin-2-yl-ethylamine (57% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 8.90 (br s, 1H), 8.51 (br m, 2H), 8.36 (s, 1H), 8.22 (m, 2H), 7.71 (t, 1H), 7.48 (s, 1H), 7.43 (d, 1H), 7.26 (d, 1H), 7.21 (m, 1H), 3.97 (s, 3H), 3.60 (m, 2H), 3.00 (dd, 2H)

LRMS (esi, positive) *m/e* 393.3 (M+1)

Compound 174:




3-Methoxy-4-(3-pyrazin-2-yl-ureido)-N-(2-pyridin-4-yl-ethyl)-benzamide

Prepared according to the procedure of Compound 167 using 2-pyridin-4-yl-ethylamine (45% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 8.93 (s, 1H), 8.63 (d, 2H), 8.51 (t, 1H), 8.37 (s, 1H), 8.27 (s, 1H), 8.24 (d, 1H), 7.60 (d, 2H), 7.43 (m, 2H), 3.97 (s, 3H), 3.58 (m, 2H), 3.01 (m, 2H)

LRMS (esi, positive) *m/e* 393.1 (M+1)

Compound 175:




N-(1H-Benzoimidazol-2-ylmethyl)-3-methoxy-4-(3-pyrazin-2-yl-ureido)-benzamide

Prepared according to the procedure of Compound 167 using C-(1H-benzoimidazol-2-yl)-methylamine (53% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 8.90 (s, 1H), 8.35 (s, 1H), 8.32 (d, 1H), 8.22 (s, 1H), 7.61 (m, 3H), 7.47 (m, 2H), 7.12 (m, 2H), 4.66 (s, 2H), 3.98 (s, 3H)

LRMS (esi, positive) *m/e* 418.2 (M+1)

Compound 176:




N-[2-(1H-Indol-3-yl)-ethyl]-3-methoxy-4-(3-pyrazin-2-yl-ureido)-benzamide

Prepared according to the procedure of Compound 167 using tryptamine (74% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 8.92 (s, 1H), 8.55 (br t, 1H), 8.26 (s, 1H), 8.24 (d, 1H), 8.23 (s, 1H), 7.60 (d, 1H), 7.53 (s, 1H), 7.50 (d, 1H), 7.26 (d, 1H), 7.19 (s, 1H), 7.05 (dd, 1H), 6.98 (dd, 1H), 3.97 (s, 3H), 3.56 (m, 2H), 2.96 (m, 2H)

LRMS (esi, positive) *m/e* 431.2 (M+1)

Compound 177:




3-Methoxy-N-[3-(methyl-phenyl-amino)-propyl]-4-(3-pyrazin-2-yl-ureido)-benzamide

Prepared according to the procedure of Compound 167 using N1-methyl-N1-phenyl-propane-1,3-diamine (68% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 8.89 (s, 1H), 8.41 (br s, 1H), 8.36 (s, 1H), 8.23 (d, 1H), 8.22 (s, 1H), 7.52 (s, 1H), 7.49 (d, 1H), 7.16 (m, 2H), 6.70 (d, 2H), 6.59 (dd, 1H), 3.96 (s, 3H), 3.38 (m, 2H), 3.30 (m, 2H), 2.87 (s, 3H), 1.77 (m, 2H)

LRMS (esi, positive) *m/e* 435.2 (M+1)

Compound 178:




N-(1-Benzyl-pyrrolidin-3-yl)-3-methoxy-4-(3-pyrazin-2-yl-ureido)-benzamide

Prepared according to the procedure of Compound 167 using 3-amino-1-benzyl pyrrolidine (62% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 8.88 (br s, 1H), 8.36 (br m, 2H), 8.24 (d, 1H), 8.21 (s, 1H), 7.51 (s, 1H), 7.50 (d, 1H), 7.32 (m, 4H), 7.22 (m, 1H), 4.39 (br m, 1H), 3.96 (s, 3H), 3.59 (s, 2H), 2.79 (m, 1H), 2.62 (m, 1H), 2.40 (m, 1H), 2.16 (m, 1H), 1.81 (m, 2H)

LRMS (esi, positive) *m/e* 447.2 (M+1)

Compound 179:




N-(3-(*R*)-1-Benzyl-pyrrolidin-3-yl)-3-methoxy-4-(3-pyrazin-2-yl-ureido)-benzamide

Prepared according to the procedure of Compound 167 using 3-(*R*)-amino-1-benzyl pyrrolidine (57% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 8.83 (br s, 1H), 8.36-8.24 (m, 3H), 8.15 (m, 1H), 7.48 (m, 2H), 7.32 (m, 4H), 7.22 (m, 1H), 4.37 (m, 1H), 3.96 (s, 3H), 3.59 (s, 2H), 2.78 (m, 1H), 2.63 (m, 1H), 2.41 (m, 1H), 2.17 (m, 1H), 1.80 (m, 2H)

LRMS (esi, positive) *m/e* 447.1 (M+1)

Compound 180:




N-(3-(*S*)-1-Benzyl-pyrrolidin-3-yl)-3-methoxy-4-(3-pyrazin-2-yl-ureido)-benzamide

Prepared according to the procedure of Compound 167 using 3-(*S*)-amino-1-benzyl pyrrolidine (57% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 8.85 (s, 1H), 8.34 (br s, 1H), 8.30 (s, 1H), 8.25 (d, 1H), 8.19 (s, 1H), 7.50 (m, 2H), 7.32 (m, 4H), 7.22 (m, 1H), 4.39 (m, 1H), 3.97 (s, 3H), 3.59 (s, 2H), 2.79 (m, 1H), 2.62 (m, 1H), 2.41 (m, 1H), 2.16 (m, 1H), 1.81 (m, 2H)

LRMS (esi, positive) m/e 447.1 (M+1)

Compound 181:




N-(2-Dimethylamino-ethyl)-3-methoxy-N-methyl-4-(3-pyrazin-2-yl-ureido)-benzamide

Prepared according to the procedure of Compound 167 using N,N,N'-trimethyl-ethane-1,2-diamine (57% yield).

¹H-NMR (400 MHz, D₂O) δ 8.17 (s, 1H), 8.05 (s, 1H), 7.94 (s, 1H), 7.76 (d, 1H), 6.92 (m, 2H), 3.79 (m, 2H), 3.75 (s, 3H), 3.36 (m, 2H), 2.97 (s, 3H), 2.88 (s, 6H)

LRMS (esi, positive) m/e 373.2 (M+1)

Compound 182:




3-Methoxy-N-(3-methylamino-propyl)-4-(3-pyrazin-2-yl-ureido)-benzamide

Prepared according to the procedure of Compound 167 using N1-methyl-propane-1,3-diamine (25% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 8.93 (s, 1H), 8.38-8.25 (m, 4H), 7.59 (d, 1H), 7.52 (m, 1H), 3.98 (s, 3H), 3.92 (m, 2H), 2.92, (m, 2H), 2.50 (s, 3H), 1.82 (m, 2H)

LRMS (esi, positive) m/e 359.1 (M+1)

Compound 183:




N-(3-Dimethylamino-propyl)-3-methoxy-4-(3-pyrazin-2-yl-ureido)-benzamide

Prepared according to the procedure of Compound 167 using N,N-dimethyl propyldiamine (81% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 8.93 (s, 1H), 8.56 (t, 1H), 8.37 (s, 1H), 8.25 (d, 1H), 8.23 (s, 1H), 7.52 (s, 1H), 7.50 (d, 1H), 4.10 (m, 2H), 3.97 (s, 3H), 3.35 (s, 6H), 3.05 (m, 2H), 1.84 (m, 2H)

LRMS (esi, positive) m/e 373.1 (M+1)

Compound 184:




N-(3-Dimethylamino-propyl)-3-methoxy-N-methyl-4-(3-pyrazin-2-yl-ureido)-benzamide

Prepared according to the procedure of Compound 167 using N,N,N'-trimethyl propyldiamine (88% yield).

¹H-NMR (400 MHz, CDCl₃/CD₃OD) δ 8.60 (s, 1H), 8.33 (d, 1H), 8.23 (s, 1H), 8.19 (s, 1H), 7.01 (m, 2H), 3.99 (s, 3H), 3.83 (s, 3H), 3.59 (m, 2H), 2.78 (m, 2H), 2.59 (s, 6H), 2.22 (m, 2H)

LRMS (esi, positive) m/e 387.1 (M+1)

Compound 185:




3-Methoxy-N-(3-morpholin-4-yl-propyl)-4-(3-pyrazin-2-yl-ureido)-benzamide

Prepared according to the procedure of Compound 167 using 3-morpholin-4-yl-propylamine (53% yield).

¹H-NMR (400 MHz, d₆-DMSO) δ 8.92 (s, 1H), 8.57 (t, 1H), 8.36 (s, 1H), 8.28 (d, 1H), 8.24 (s, 1H), 7.52 (s, 1H), 7.50 (d, 1H), 3.96 (s, 3H), 3.61 (m, 2H), 3.42 (m, 2H), 3.32 (m, 4H), 3.10 (m, 4H), 1.90 (m, 2H)

LRMS (esi, positive) m/e 415.1 (M+1)

Compound 186:




3-Methoxy-N-[3-(4-methyl-piperazin-1-yl)-propyl]-4-(3-pyrazin-2-yl-ureido)-benzamide

Prepared according to the procedure of Compound 167 using 3-(4-methyl-piperazin-1-yl)-propylamine (63% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 8.92 (s, 1H), 8.41 (m, 1H), 8.36 (s, 1H), 8.24 (m, 2H), 7.52 (s, 1H), 7.49 (d, 1H), 3.97 (s, 3H), 3.31 (m, 11H), 2.70 (m, 2H), 2.41 (m, 2H), 1.72 (m, 2H)

LRMS (esi, positive) *m/e* 428.1 (M+1)

Compound 187:




{2-[3-Methoxy-4-(3-pyrazin-2-yl-ureido)-benzoylamino]-ethyl}-trimethyl-ammonium chloride

Prepared according to the procedure of Compound 167 using 2-N,N,N-trimethylammonium ethylamine(46% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 8.92 (s, 1H), 8.77 (m, 1H), 8.36 (s, 1H), 8.30 (d, 1H), 8.24 (s, 1H), 7.53 (s, 1H), 7.51 (d, 1H), 3.97 (s, 3H), 3.70 (m, 2H), 3.50 (m, 2H), 3.15 (s, 9H)

LRMS (esi, positive) *m/e* 373.1 (M+)

Compound 188:



4-Methoxy-3-(3-pyrazin-2-yl-ureido)-benzoic acid


Step 1: 4-Methoxy-3-(4-nitro-phenoxy-carbonylamino)-benzoic acid methyl ester. To a stirred, cooled (0°C) solution of methyl-3-amino-4-methoxy benzoate (5.0 g; 27.6 mmol) in methylene chloride (100 mL) was added pyridine (2.34 mL; 29 mmol) followed by 4 nitrophenyl chloroformate (5.8 g; 29 mmol). After stirring for 8 hours, the reaction was diluted with methylene chloride (100 mL), washed with 1N hydrochloric acid (2 x 125 mL), 10% aqueous sodium carbonate (2 x 125 mL), brine (1 x 125 mL), then dried (MgSO₄), and filtered. The filtered material was concentrated under reduced pressure. The residue was taken up in ethyl acetate (100 mL) followed by hexanes (700 mL). A precipitate formed which was filtered to yield an off white solid (80% yield).

Step 2: 4-Methoxy-3-(3-pyrazin-2-yl-ureido)-benzoic acid methyl ester. To a stirred solution of the carbamate piece (1.0 g; 2.9 mmol) in N-methyl pyrrolidinone (5 mL) was added amino pyrazine (285 mg; 3.0 mmol). The reaction was heated to 85 °C and stirred for 12 hours. The reaction was allowed to cool to room temperature, then diluted with ethyl acetate (50 mL) and water (50 mL). A precipitate formed which was filtered and dried under reduced pressure to yield an off white solid (55% yield).

Step 3: 4-Methoxy-3-(3-pyrazin-2-yl-ureido)-benzoic acid. To a stirred solution of 4-methoxy-3-(3-pyrazin-2-yl-ureido)-benzoic acid methyl ester (1.0 g; 3.3 mmol) in methanol (25 mL) was added

lithium hydroxide (5 mL of a 2M aqueous solution). The reaction was heated to 60 °C and stirred for 12 hours. The reaction was allowed to cool to room temperature and the pH was adjusted to 5.5 with hydrochloric acid (1N). A precipitate formed which was filtered and dried under reduced pressure to yield an off white solid (58% yield).

Compound 189:




N-Butyl-4-methoxy-3-(3-pyrazin-2-yl-ureido)-benzamide

To a stirred solution of 4-Methoxy-3-(3-pyrazin-2-yl-ureido)-benzoic acid (32 mg; 0.11 mmol) in N-methyl pyrrolidinone (1 mL) was added O-benzotrazol-1-yl-N,N,N',N'-tetramethyl-uronium hexafluorophosphate (HBTU; 45 mg; 0.12 mmol), butyl amine (12 µL; 0.12 mmol), and diisopropylethylamine (35 µL; 0.20 mmol). The reaction was stirred at room temperature for 12 hours. The reaction was diluted with ethyl acetate (20 mL) and 10% aqueous sodium carbonate (20 mL) and stirred for 5 minutes. A precipitate formed which was filtered and dried under reduced pressure to yield an off white solid (49% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 8.90 (s, 1H), 8.63 (s, 1H), 8.35 (s, 1H), 8.23 (m, 1H), 8.21 (s, 1H), 7.52 (d, 1H), 7.06 (d, 1H), 3.95 (s, 3H), 3.22 (m, 2H), 1.50 (m, 2H), 1.33 (m, 2H), 0.90 (t, 3H)

LRMS (apci, positive) *m/e* 344.1 (M+1)

Compound 190:




N-Benzyl-4-methoxy-3-(3-pyrazin-2-yl-ureido)-benzamide

Prepared according to the procedure of Compound 189 using benzyl amine (70% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 8.92 (s, 1H), 8.90 (br s, 1H), 8.73 (s, 1H), 8.33 (s, 1H), 8.23 (s, 1H), 7.60 (d, 1H), 7.33 (m, 4H), 7.23 (m, 1H), 7.12 (d, 1H), 4.44 (s, 2H), 3.96 (s, 3H)

LRMS (apci positive) *m/e* 378.1 (M+1)

Compound 191:




4-Methoxy-N-phenethyl-3-(3-pyrazin-2-yl-ureido)-benzamide

Prepared according to the procedure of Compound 189 using phenethyl amine (68% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 8.92 (s, 1H), 8.64 (s, 1H), 8.40 (m, 1H), 8.35 (s, 1H), 8.23 (s, 1H), 7.52 (d, 1H), 7.33-7.18 (m, 5H), 7.08 (d, 1H), 3.96 (s, 3H), 3.44 (m, 2H), 2.82 (m, 2H)

LRMS (apci positive) *m/e* 392.1 (M+1)

Compound 192:




4-Methoxy-N-(3-phenyl-propyl)-3-(3-pyrazin-2-yl-ureido)-benzamide

Prepared according to the procedure of Compound 189 using phenpropyl amine (65% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 8.92 (s, 1H), 8.64 (s, 1H), 8.36 (s, 1H), 8.23 (s, 1H), 7.56 (d, 1H), 7.32-7.22 (m, 5H), 7.18 (m, 1H), 7.10 (d, 1H), 3.97 (s, 3H), 3.24 (m, 2H), 2.62 (dd, 2H), 1.82 (m, 2H)

LRMS (apci positive) *m/e* 406.1 (M+1)

Compound 193:




N-(2-Benzenesulfonyl-ethyl)-4-methoxy-3-(3-pyrazin-2-yl-ureido)-benzamide

Prepared according to the procedure of Compound 189 using 2-benzenesulfonyl-ethylamine (42% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 8.90 (s, 1H), 8.58 (s, 1H), 8.38 (br s, 1H), 8.32 (s, 1H), 8.23 (s, 1H), 7.94 (d, 2H), 7.74 (m, 1H), 7.65 (d, 2H), 7.38 (d, 1H), 7.05 (d, 1H), 3.95 (s, 3H), 3.57 (m, 2H), 3.51 (m, 2H)

LRMS (apci positive) *m/e* 456.0 (M+1)

Compound 194:




4-Methoxy-3-(3-pyrazin-2-yl-ureido)-N-(2-pyridin-2-yl-ethyl)-benzamide

Prepared according to the procedure of Compound 189 using 2-pyridin-2-yl-ethylamine (16% yield)

¹H-NMR (400 MHz, *d*₆-DMSO) δ 8.90 (s, 1H), 8.64 (s, 1H), 8.52 (d, 1H), 8.41 (m, 1H), 8.32 (s, 1H), 8.21 (s, 1H), 7.72 (m, 1H), 7.50 (d, 1H), 7.28 (d, 1H), 7.22 (m, 1H), 7.08 (d, 1H), 3.96 (s, 3H), 3.59 (m, 2H), 2.98 (m, 2H)

LRMS (esi positive) *m/e* 415.2 (M+1)

Compound 195:




4-Methoxy-3-(3-pyrazin-2-yl-ureido)-N-(2-pyridin-4-yl-ethyl)-benzamide

Prepared according to the procedure of Compound 189 using 2-pyridin-4-yl-ethylamine (41% yield)

¹H-NMR (400 MHz, *d*₆-DMSO) δ 8.91 (s, 1H), 8.63 (s, 1H), 8.46 (d, 2H), 8.40 (m, 1H), 8.35 (s, 1H), 8.24 (s, 1H), 7.47 (d, 1H), 7.26 (d, 2H), 7.08 (d, 1H), 3.96 (s, 3H), 3.50 (m, 2H), 2.84 (m, 2H)

LRMS (esi positive) *m/e* 415.2 (M+1)

Compound 196:




N-(1H-Benzoimidazol-2-ylmethyl)-4-methoxy-3-(3-pyrazin-2-yl-ureido)-benzamide

Prepared according to the procedure of Compound 189 using C-(1H-benzoimidazol-2-yl)-methylamine (26% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 8.99 (t, 1H), 8.91 (s, 1H), 8.77 (s, 1H), 8.36 (s, 1H), 8.23 (s, 1H), 7.66 (d, 1H), 7.56 (d, 1H), 7.44 (d, 1H), 7.15 (m, 3H), 4.65 (d, 2H), 3.97 (s, 3H)

LRMS (esi positive) m/e 418.1 (M+1)

Compound 197:




N-[2-(1H-Indol-3-yl)-ethyl]-4-methoxy-3-(3-pyrazin-2-yl-ureido)-benzamide

Prepared according to the procedure of Compound 189 using tryptamine (51% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 8.92 (s, 1H), 8.69 (s, 1H), 8.43 (t, 1H), 8.33 (s, 1H), 8.24 (s, 1H), 7.60 (d, 1H), 7.56 (d, 1H), 7.35 (d, 1H), 7.17 (s, 1H), 7.11 (d, 1H), 7.07 (m, 1H), 6.97 (m, 1H), 3.96 (s, 3H), 3.54 (m, 2H), 2.95 (m, 2H).

LRMS (esi positive) m/e 431.1 (M+1)

Compound 198:




4-Methoxy-N-[3-(methyl-phenyl-amino)-propyl]-3-(3-pyrazin-2-yl-ureido)-benzamide

Prepared according to the procedure of Compound 189 using N-methyl-N-phenyl propyldiamine (81% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 8.92 (s, 1H), 8.64 (s, 1H), 8.37 (m, 1H), 8.35 (s, 1H), 8.23 (s, 1H), 7.55 (d, 1H), 7.14 (m, 3H), 6.70 (d, 2H), 6.58 (t, 1H), 3.96 (s, 3H), 3.37 (m, 2H), 3.25 (m, 2H), 2.86 (s, 3H), 1.77 (m, 2H).

LRMS (esi positive) m/e 435.2 (M+1)

Compound 199:




N-(1-Benzyl-pyrrolidin-3-yl)-4-methoxy-3-(3-pyrazin-2-yl-ureido)-benzamide

Prepared according to the procedure of Compound 189 using 3-amino-1-benzyl pyrrolidine (48% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 8.90 (s, 1H), 8.62 (s, 1H), 8.36 (s, 1H), 8.30 (d, 1H), 8.22 (s, 1H), 7.56 (d, 1H), 7.32 (m, 4H), 7.22 (m, 1H), 7.15 (d, 1H), 4.37 (m, 1H), 3.96 (s, 3H), 3.59 (s, 2H), 2.79 (m, 1H), 2.60 (m, 1H), 2.39 (m, 1H), 2.14 (m, 1H), 1.80 (m, 2H).

LRMS (esi positive) *m/e* 447.2 (M+1)

Compound 200:




N-(2-Dimethylamino-ethyl)-4-methoxy-N-methyl-3-(3-pyrazin-2-yl-ureido)-benzamide

Prepared according to the procedure of Compound 189 using N,N,N'-trimethyl ethyldiamine (93% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 9.39 (br s, 1H), 8.91 (s, 1H), 8.35 (s, 1H), 8.33 (d, 1H), 8.23 (s, 1H), 7.16-7.09 (m, 2H), 3.96 (s, 3H), 3.76 (m, 2H), 3.37 (m, 2H), 2.99 (s, 3H), 2.85 (br s, 6H).

LRMS (esi, positive) *m/e* 373.2 (M+1)

Compound 201:




4-Methoxy-N-(3-methylamino-propyl)-3-(3-pyrazin-2-yl-ureido)-benzamide

Prepared according to the procedure of Compound 189 using N-methyl propyldiamine (15% yield).

¹H-NMR (400 MHz, D₂O) δ 7.98 (m, 2H), 7.91 (s, 1H), 7.82 (s, 1H), 7.17 (d, 1H), 6.73 (d, 1H), 3.73 (s, 3H), 3.29 (m, 2H), 2.98 (m, 2H), 2.61 (s, 3H), 1.88 (m, 2H).

LRMS (esi, positive) *m/e* 359.2 (M+1)

Compound 202:




N-(3-Dimethylamino-propyl)-4-methoxy-3-(3-pyrazin-2-yl-ureido)-benzamide

Prepared according to the procedure of Compound 189 using N,N-dimethyl propyldiamine (51% yield).

¹H-NMR (400 MHz, D₂O) δ 7.98 (s, 1H), 7.96 (s, 2H), 7.81 (s, 1H), 7.16 (d, 1H), 6.72 (d, 1H), 3.73 (s, 3H), 3.29 (m, 2H), 3.09 (m, 2H), 2.80 (s, 6H), 1.93 (m, 2H).

LRMS (esi, positive) m/e 373.2 (M+1)

Compound 203:




N-(3-Dimethylamino-propyl)-4-methoxy-N-methyl-3-(3-pyrazin-2-yl-ureido)-benzamide

Prepared according to the procedure of Compound 189 using N,N,N'-trimethyl propyldiamine (60% yield).

¹H-NMR (400 MHz, D₂O) δ 8.37 (s, 1H), 8.19 (s, 1H), 8.04 (s, 1H), 7.80 (d, 1H), 7.13 (m, 1H), 7.01 (m, 1H), 3.82 (s, 3H), 3.51 (m, 2H), 3.11 (m, 2H), 2.96 (s, 3H), 2.80 (s, 6H), 2.01 (m, 2H).

LRMS (esi, positive) m/e 387.1 (M+1)

Compound 204:




4-Methoxy-N-[3-(4-methyl-piperazin-1-yl)-propyl]-3-(3-pyrazin-2-yl-ureido)-benzamide

Prepared according to the procedure of Compound 189 using 3-(4-methyl-piperazin-1-yl)-propylamine (57% yield).

¹H-NMR (400 MHz, D₂O) δ 8.18 (s, 1H), 8.06 (s, 1H), 8.04 (s, 1H), 7.96 (s, 1H), 7.32 (d, 1H), 6.85 (d, 1H), 3.79 (s, 3H), 3.48 (br s, 8H), 3.36 (m, 2H), 3.17 (m, 2H), 2.83 (s, 3H), 1.96 (m, 2H).

LRMS (esi, positive) m/e 428.2 (M+1)

Compound 205:




{2-[4-Methoxy-3-(3-pyrazin-2-yl-ureido)-benzoylamino]-ethyl}-trimethyl-ammonium chloride

Prepared according to the procedure of Compound 189 using 2-trimethylammonium ethyl amine (63% yield).

¹H-NMR (400 MHz, D₂O) δ 8.17 (s, 1H), 8.05 (s, 1H), 8.03 (s, 1H), 7.93 (s, 1H), 7.32 (d, 1H), 6.84 (d, 1H), 3.80 (s, 3H), 3.76 (m, 2H), 3.44 (m, 2H), 3.11 (s, 9H).

LRMS (esi, positive) m/e 373.0 (M⁺)

Compound 206:




4-Methoxy-N-(3-morpholin-4-yl-propyl)-3-(3-pyrazin-2-yl-ureido)-benzamide

Prepared according to the procedure of Compound 189 using 3-morpholin-4-yl-propylamine (69% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 8.91 (s, 1H), 8.63 (s, 1H), 8.36 (s, 1H), 8.35 (m, 1H), 8.23 (s, 1H), 7.54 (d, 1H), 7.10 (d, 1H), 3.96 (s, 3H), 3.57 (m, 4H), 3.26 (m, 2H), 2.34 (m, 4H), 2.32 (m, 2H), 1.66 (m, 2H).

LRMS (esi, positive) m/e 415.2 (M+1)

Compound 207:



3-Methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzoic acid

Step 1: (5-Methyl-pyrazin-2-yl)-carbamic acid tert-butyl ester. To a stirred solution of 5-methyl pyrazine carboxylic acid (13.8 g, 100 mmol) in 300 mL of toluene at room temperature under nitrogen was added triethyl amine (14 mL, 100 mmol) followed by diphenyl phosphoryl azide (21.6 mL, 100 mmol). After 30 min. at room temperature, 2-methyl-2-propanol (19 mL, 200 mmol) was added and the solution immersed in a 90 °C oil bath. After 2 hours, the reaction was cooled to RT, diluted to 600


mL with EtOAc, and washed 3 x 60 mL with 10% Na₂CO₃ and 1 x 600 mL with saturated NaCl. The organics were dried (MgSO₄), filtered and concentrated to a yellow solid (17.5 g, 83%). ¹H-NMR (400 MHz, CDCl₃) δ 9.16 (s, 1H), 8.05 (s, 1H), 7.56 (br s, 1H), 2.50 (s, 3H), 1.55 (s, 9H).

Step 2: 5-Methyl-2-aminopyrazine. To a stirred solution of (5-methyl-pyrazin-2-yl)-carbamic acid tert-butyl ester (2.1 g, 10 mmol) in 30 mL CH₂Cl₂ at 0 °C under nitrogen was added trifluoroacetic acid (30 mL). The solution was allowed to warm to RT overnight. The solution was rotary evaporated to remove TFA and the residue was redissolved in 200 mL CH₂Cl₂ and stirred with 100 mL 10% Na₂CO₃. The organics were isolated and the aqueous solution extracted 3 x 100 mL with CH₂Cl₂. The organics were combined, dried (MgSO₄), filtered and concentrated to an orange solid (1 g, 92%). ¹H-NMR (400 MHz, CDCl₃) δ 8.46 (s, 1H), 7.70 (s, 1H), 2.49 (s, 3H).

Step 3: 3-Methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzoic acid methyl ester. To a stirred solution of 3-methoxy-4-(4-nitro-phenoxy-carbonylamino)-benzoic acid methyl ester (11.7 g, 33.8 mmol) in 34 mL NMP at room temperature under nitrogen was added 5-methyl-2-aminopyrazine (3.69 g, 33.8 mmol) and the reaction was immersed in an 85 °C oil bath. After 6 hours the reaction was allowed to cool to room temperature and a precipitate formed. EtOAc (200 mL) was added and the precipitate was isolated by filtration (4.7 g, 44%). ¹H-NMR (400 MHz, d₆-DMSO) δ 8.79 (br s, 1H), 8.36 (d, 1H), 8.23 (s, 1H), 7.60 (d, 1H), 7.52 (s, 1H), 3.98 (s, 3H), 3.81 (s, 3H), 2.42 (s, 3H).

Step 4: 3-Methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzoic acid. To a stirred suspension of 3-methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzoic acid methyl ester (7.15 g, 22.6 mmol) in 3:1 MeOH:H₂O (226 mL) at RT under N₂ was added lithium hydroxide monohydrate (9.5 g, 226 mmol) as a solid and the mixture heated to 65 °C. After reaching temperature, the suspension gradually became a bright yellow solution. After about 4 hours a precipitate formed but the reaction was continued overnight. After cooling to RT, MeOH was removed by rotovap and the aqueous suspension diluted with 100 mL H₂O and neutralized to pH=5 with concentrated HCl. As pH=5 was approached, the suspension turned from yellow to white. The suspension was then filtered through paper on a large ceramic funnel. The filtration went very slowly, taking several hours. The filter cake was washed twice with H₂O. When most of the H₂O was removed, the residue was dried under high vacuum in a dessicator overnight to give the free acid as a white solid (6 g, 88%). ¹H-NMR (400 MHz, d₆-DMSO) δ 8.79 (br s, 1H), 8.36 (d, 1H), 8.22 (s, 1H), 7.57 (d, 1H), 7.51 (s, 1H), 3.97 (s, 3H), 2.42 (s, 3H).

Compound 208:



3-Methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-N-(2-pyridin-2-yl-ethyl)-benzamide


To a stirred solution of 3-methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzoic acid (30 mg, 0.1 mmol) in 1 mL NMP at RT in a capped reaction vial was added HBTU (42 mg, 0.11 mmol). The

suspension was stirred for 15 minutes and then treated with 2-ethylaminopyridine (13.2 μ L, 0.11 mmol) followed by Hunigs Base (35 μ L, 0.2 mmol). After stirring overnight, NMP was removed by bulb to bulb transfer at 70 °C under high vacuum and the residue stirred with a mixture of CH_2Cl_2 (10 mL) and 10% Na_2CO_3 (10 mL) until complete dissolution occurred. The organics were isolated, dried (MgSO_4), filtered and concentrated. The residue was triturated with EtOAc to produce a solid which was isolated by filtration (71% yield).

$^1\text{H-NMR}$ (400 MHz, d_6 -DMSO) δ 8.90 (s, 1H), 8.58 (d, 1H), 8.50 (t, 1H), 8.23 (d, 1H), 8.21 (s, 1H), 7.82 (t, 1H), 7.48-7.31 (m, 4H), 3.97 (s, 3H), 3.62 (m, 2H), 3.04 (m, 2H), 2.42 (s, 3H).

LRMS (esi, positive) m/e 407.1 (M+1)

Compound 209:




N-(1-Benzyl-piperidin-4-yl)-3-methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzamide

Prepared according to the method of Compound 208 using 4-amino-1-benzyl piperidine except the crude product was purified by chromatography on a Biotage 12S column eluting with 92.5/7.5 $\text{CH}_2\text{Cl}_2/\text{MeOH}$. (61% yield).

$^1\text{H-NMR}$ (400 MHz, $\text{CDCl}_3/\text{CD}_3\text{OD}$) δ 8.44 (br s, 1H), 8.32 (d, 1H), 8.09 (s, 1H), 7.47 (s, 1H), 7.38-7.28 (m, 6H), 7.09 (d, 1H), 4.00 (m, 1H), 3.99 (s, 3H), 3.60 (s, 2H), 2.98 (m, 2H), 2.52 (s, 3H), 2.25 (m, 2H), 2.00 (m, 2H), 1.64 (m, 2H)

LRMS (esi, positive) m/e 475.2 (M+1)

Compound 210:




N-(3-Dimethylamino-propyl)-3-methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzamide

Prepared according to the method of Compound 208 using N,N-dimethyl propyldiamine (70% yield).

$^1\text{H-NMR}$ (400 MHz, d_6 -DMSO) δ 8.90 (br s, 1H), 8.55 (t, 1H), 8.26 (d, 1H), 8.22 (s, 1H), 7.52 (s, 1H), 7.50 (d, 1H), 3.98 (s, 3H), 3.32 (m, 2H), 3.07 (m, 2H), 2.77 (s, 6H), 2.41 (s, 3H), 1.89 (m, 2H).

LRMS (esi, positive) m/e 387.1 (M+1)

Compound 211:




3-Methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-N-(3-morpholin-4-yl-propyl)-benzamide

Prepared according to the method of Compound 208 using 3-morpholin-4-yl-propylamine (79% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 8.90 (br s, 1H), 8.57 (t, 1H), 8.25 (d, 1H), 8.21 (s, 1H), 7.51 (s, 1H), 7.50 (d, 1H), 3.97 (s, 3H), 3.62 (m, 2H), 3.31 (m, 8H), 3.12 (m, 2H), 2.41 (s, 3H), 1.92 (m, 2H)

LRMS (esi, positive) *m/e* 429.1 (M+1)

Compound 212:




N-(2-Dimethylamino-2-phenyl-ethyl)-3-methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzamide

Prepared according to the method of Compound 208 using N-(2-dimethylamino)-2-phenyl ethyl amine except the crude product was purified by chromatography on a Biotage 12S column eluting with 92.5/7.5 CH₂Cl₂/MeOH. (12% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 10.05 (s, 1H), 8.88 (s, 1H), 8.20 (m, 3H), 7.41-7.19 (m, 7H), 3.96 (s, 3H), 3.78 (m, 1H), 3.70 (m, 1H), 3.57 (m, 1H), 3.38 (m, 1H), 2.50 (s, 6H), 2.40 (s, 3H)

LRMS (esi, positive) *m/e* 448.9 (M+1)

Compound 213:




N-(2-Dimethylamino-1-phenyl-ethyl)-3-methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzamide

Prepared according to the method of Compound 208 using N-(2-dimethylamino)-1-phenyl ethyl amine except the crude product was purified by chromatography on a Biotage 12S column eluting with 92.5/7.5 CH₂Cl₂/MeOH. (27% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 11.61 (br s, 1H), 9.71 (br s, 1H), 8.42 (d, 1H), 8.36 (s, 1H), 8.09 (s, 1H), 7.52-7.21 (m, 7H), 5.05 (br s, 1H), 3.93 (s, 3H), 2.80 (m, 1H), 2.52 (s, 3H), 2.37 (s, 6H), 1.79 (br s, 2H)

LRMS (esi, positive) m/e 449.0 (M+1)

Compound 214:




N-(1-Aza-bicyclo[2.2.2]oct-3-yl)-3-methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzamide

Prepared according to the method of Compound 208 using 1-aza-bicyclo[2.2.2]oct-3-ylamine (27% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 10.17 (br s, 1H), 8.90 (br s, 1H), 8.24 (d, 1H), 8.23 (s, 1H), 8.17 (d, 1H), 7.53 (d, 1H), 7.50 (s, 1H), 3.98 (s, 3H), 3.95 (m, 1H), 3.10 (m, 1H), 2.90 (m, 1H), 2.64 (m, 4H), 2.42 (s, 3H), 1.87 (m, 1H), 1.80 (m, 1H), 1.59 (m, 2H), 1.31 (m, 1H).

LRMS (esi, positive) m/e 411.0 (M+1)

Compound 215:



3-Methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-N-(3-*R*-1-pyridin-2-ylmethyl-pyrrolidin-3-yl)-benzamide

Step 1: (3-*R*-1-Pyridin-2-ylmethyl-pyrrolidin-3-yl)-carbamic acid tert-butyl ester. To a stirred solution of (R)-*boc*-3-aminopyrrolidine (372.5 mg, 2 mmol) in dichloroethane (6 mL) at room temperature under nitrogen was added pyridine-2-carboxaldehyde (190 μL, 2 mmol) followed by sodium triacetoxyborohydride (593 mg, 2.8 mmol). The reaction was stirred at room temperature overnight and was then quenched by addition of saturated NaHCO₃ (6 mL) with stirring for 15 minutes. The reaction was then partitioned between CH₂Cl₂ (25 mL) and 10% Na₂CO₃ (25 mL). The organics were isolated, dried (MgSO₄), filtered and concentrated to the pure product (526 mg, 95%).

Step 2: 3-*R*-1-Pyridin-2-ylmethyl-pyrrolidin-3-ylamine dihydrochloride. A stirred solution of (3-*R*-1-pyridin-2-ylmethyl-pyrrolidin-3-yl)-carbamic acid tert-butyl ester (277 mg, 1 mmol) in 10 mL 4N HCl in dioxane at room temperature in a capped flask was reacted overnight. The reaction was concentrated by rotary evaporation and high vacuum to give the di HCl salt (250 mg, quantitative).


Step 3: Prepared according to the procedure of Compound 208 except 3-*R*-1-pyridin-2-ylmethyl-pyrrolidin-3-ylamine dihydrochloride salt was mixed with excess DIEA (70 μL, 0.4 mmol) in 500 μL

NMP to form a solution which was added to the acid/HBTU mixture. The crude product was purified by chromatography on a Biotage 12S column eluting with 9/1 CH₂Cl₂/MeOH (60% yield).

¹H-NMR (400 MHz, CDCl₃) δ 8.57 (s, 1H), 8.51 (d, 1H), 8.39 (d, 1H), 8.17 (s, 1H), 8.09 (s, 1H), 7.92 (s, 1H), 7.64 (d, 1H), 7.50 (s, 1H), 7.30 (d, 1H), 6.70 (d, 1H), 4.69 (m, 1H), 4.00 (s, 3H), 3.66 (dd, 2H), 2.98 (m, 1H), 2.80 (m, 1H), 2.70 (m, 1H), 2.53 (s, 3H), 2.39 (m, 3H), 1.76 (m, 1H).

LRMS (esi, positive) m/e 462.3 (M+1)

Compound 216:




3-Methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-N-(3-*R*-1-methyl-pyrrolidin-3-yl)-benzamide

Prepared according to the procedure of Compound 215 using 3-(*R*)-amino-1-methyl pyrrolidine (29% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 8.78 (br s, 1H), 8.36 (d, 1H), 8.22 (d, 1H), 8.20 (s, 1H), 7.52 (s, 1H), 7.50 (d, 1H), 4.39 (m, 1H), 3.96 (s, 3H), 2.64 (m, 1H), 2.61 (m, 1H), 2.43, (s, 3H), 2.39 (m, 2H), 2.24 (s, 3H), 2.17 (m, 1H), 1.76 (m, 1H).

LRMS (esi, positive) m/e 385.3 (M+1)

Compound 217:




N-(3-*R*-1-Benzyl-pyrrolidin-3-yl)-3-methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzamide

Prepared according to the method of Compound 215 using 3-(*R*)-amino-1-benzyl pyrrolidine except the crude product was purified by chromatography on a Biotage 12S column eluting with 92.5/7.5 CH₂Cl₂/MeOH. (54% yield).

¹H-NMR (400 MHz, CDCl₃) δ 9.09 (br s, 1H), 8.43 (d, 1H), 8.09 (s, 1H), 8.02 (s, 1H), 7.57 (m, 1H), 7.53 (s, 1H), 7.46 (d, 1H), 7.33-7.22 (m, 4H), 4.79 (m, 1H), 4.00 (s, 3H), 3.78 (dd, 2H), 3.13 (m, 2H), 2.76 (m, 1H), 2.55 (s, 3H), 2.44 (m, 1H), 1.79 (m, 1H).

LRMS (esi, positive) m/e 461.1 (M+1)

Compound 218:




3-Methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-N-(3-R-1-pyridin-4-ylmethyl-pyrrolidin-3-yl)-benzamide

Prepared according to the procedure of Compound 215 using 1-(R)-Pyridin-4-ylmethyl-pyrrolidin-3-ylamine (60% yield).

¹H-NMR (400 MHz, CDCl₃) δ 8.56 (d, 2H), 8.41 (d, 1H), 8.21 (d, 1H), 8.20 (s, 1H), 8.08 (s, 1H), 7.52 (s, 1H), 7.32 (d, 1H), 7.25 (d, 2H), 6.71 (d, 1H), 4.73 (m, 1H), 4.00 (s, 3H), 3.66 (dd, 2H), 2.97 (m, 1H), 2.82 (m, 1H), 2.71 (m, 1H), 2.54 (s, 3H), 2.39 (m, 2H), 1.78 (m, 1H).

LRMS (esi, positive) m/e 462.3 (M+1)

Compound 219:




3-Methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-N-(3-R-1-thiophen-2-ylmethyl-pyrrolidin-3-yl)-benzamide

Prepared according to the procedure of Compound 215 using (R)-1-thiophen-2-ylmethyl-pyrrolidin-3-ylamine (60% yield).

¹H-NMR (400 MHz, CDCl₃) δ 8.96 (br s, 1H), 8.43 (d, 1H), 8.09 (d, 2H), 7.57 (s, 1H), 7.50 (d, 1H), 7.10 (d, 1H), 6.91 (m, 2H), 4.80 (m, 1H), 4.00 (s, 3H), 3.96 (dd, 2H), 3.17 (m, 1H), 3.08 (m, 1H), 2.71 (m, 1H), 2.54 (s, 3H), 2.42 (m, 2H), 1.80 (m, 1H).

LRMS (esi, positive) m/e 467.2 (M+1)

Compound 220:




N-(3-*R*-1-Cyclohexylmethyl-pyrrolidin-3-yl)-3-methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzamide

Prepared according to the procedure of Compound 215 using 1-cyclohexylmethyl-pyrrolidin-3-*R*-ylamine (71% yield).

¹H-NMR (400 MHz, CDCl₃) δ 8.89 (br s, 1H), 8.32 (d, 1H), 8.23 (d, 1H), 8.21 (s, 1H), 7.51 (s, 1H), 7.49 (d, 1H), 4.38 (m, 1H), 3.96 (s, 3H), 2.76 (m, 1H), 2.56 (m, 1H), 2.42 (s, 3H), 2.38 (m, 1H), 2.25-2.06 (m, 3H), 1.77 (m, 3H), 1.61 (m, 3H), 1.40 (m, 1H), 1.19 (m, 3H), 0.83 (m, 2H)

LRMS (esi, positive) m/e 467.3 (M+1)

Compound 221:




3-Methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-N-(3-*R*-1-methyl-pyrrolidin-3-yl)-benzamide

Prepared according to the procedure of Compound 215 using 1-methyl-pyrrolidin-3-*R*-ylamine (51% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 8.78 (br s, 1H), 8.35 (d, 1H), 8.22 (d, 1H), 8.21 (s, 1H), 7.52 (s, 1H), 7.49 (d, 1H), 4.39 (m, 1H), 3.96 (s, 3H), 2.66 (m, 1H), 2.61 (m, 1H), 2.43 (s, 3H), 2.39 (m, 2H), 2.24 (s, 3H), 2.17 (m, 1H), 1.75 (m, 1H)

LRMS (esi, positive) m/e 385.4 (M+1)

Compound 222:




N-(3-*S*-1-Benzyl-pyrrolidin-3-yl)-3-methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzamide

Prepared according to the method of Compound 215 using 1-Benzyl-pyrrolidin-3-*S*-ylamine except the crude product was purified by chromatography on a Biotage 12S column eluting with 92.5/7.5 CH₂Cl₂/MeOH. (54% yield).

¹H-NMR (400 MHz, CDCl₃) δ 8.81 (br s, 1H), 8.43 (d, 1H), 8.06 (d, 2H), 7.53 (s, 1H), 7.42 (d, 1H), 7.35-7.20 (m, 5H), 4.78 (m, 1H), 3.99 (s, 3H), 3.78 (dd, 2H), 3.15 (m, 1H), 3.04 (m, 1H), 2.77 (m, 1H), 2.54 (s, 3H), 2.44 (m, 1H), 1.79 (m, 2H).

LRMS (esi, positive) m/e 461.1 (M+1)

Compound 223:




3-Methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-N-(3-S-1-pyridin-2-ylmethyl-pyrrolidin-3-yl)-benzamide

Prepared according to the procedure of Compound 215 using 1-pyridin-2-ylmethyl-pyrrolidin-3-S-ylamine (60% yield).

¹H-NMR (400 MHz, CDCl₃) δ 8.56 (d, 1H), 8.38 (s, 1H), 8.32 (d, 1H), 8.06 (s, 1H), 7.68 (t, 1H), 7.51 (s, 1H), 7.40 (d, 1H), 7.35 (d, 1H), 7.21 (m, 1H), 4.72 (m, 1H), 4.00 (s, 3H), 3.79 (dd, 2H), 3.40 (m, 1H), 3.03 (m, 1H), 2.86 (m, 1H), 2.64 (m, 1H), 2.53 (s, 3H), 2.35 (m, 1H), 1.78 (m, 2H).

LRMS (esi, positive) m/e 462.1 (M+1)

Compound 224:




3-Methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-N-(3-S-1-pyridin-3-ylmethyl-pyrrolidin-3-yl)-benzamide

Prepared according to the procedure of Compound 215 using 1-pyridin-3-ylmethyl-pyrrolidin-3-S-ylamine (60% yield).

¹H-NMR (400 MHz, CDCl₃) δ 8.56 (s, 1H), 8.51 (m, 1H), 8.39 (d, 1H), 8.18 (s, 1H), 8.08 (s, 2H), 7.64 (d, 1H), 7.50 (s, 1H), 7.33 (d, 1H), 6.77 (d, 1H), 4.72 (m, 1H), 4.00 (s, 3H), 3.66 (dd, 2H), 2.97 (m, 1H), 2.82 (m, 1H), 2.71 (m, 1H), 2.54 (s, 3H), 2.40 (m, 2H), 1.76 (m, 2H).

LRMS (esi, positive) m/e 462.3 (M+1)

Compound 225:




3-Methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-N-(3-*S*-1-pyridin-4-ylmethyl-pyrrolidin-3-yl)-benzamide

Prepared according to the procedure of Compound 215 using 1-pyridin-4-ylmethyl-pyrrolidin-3-*S*-ylamine (60% yield).

¹H-NMR (400 MHz, CDCl₃) δ 8.56 (d, 2H), 8.41 (d, 1H), 8.24 (s, 1H), 8.20 (s, 1H), 8.08 (s, 1H), 7.53 (s, 1H), 7.32 (d, 1H), 7.24 (d, 2H), 6.75 (d, 1H), 4.72 (m, 1H), 4.00 (s, 3H), 3.67 (dd, 2H), 2.98 (m, 1H), 2.82 (m, 1H), 2.71 (m, 1H), 2.54 (s, 3H), 2.40 (m, 2H), 1.79 (m, 2H).

LRMS (esi, positive) m/e 462.3 (M+1)

Compound 226:




3-Methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-N-(3-*S*-1-thiophen-2-ylmethyl-pyrrolidin-3-yl)-benzamide

Prepared according to the procedure of Compound 215 using 1-thiophen-2-ylmethyl-pyrrolidin-3-*S*-ylamine (55% yield).

¹H-NMR (400 MHz, CDCl₃) δ 8.71 (br s, 1H), 8.42 (d, 1H), 8.11 (d, 2H), 7.55 (s, 1H), 7.44 (d, 1H), 7.39 (d, 1H), 7.20 (d, 1H), 6.91 (m, 2H), 4.78 (m, 1H), 4.00 (s, 3H), 3.94 (dd, 2H), 3.17 (m, 1H), 3.05 (m, 1H), 2.69 (m, 1H), 2.53 (s, 3H), 2.42 (m, 2H), 1.80 (m, 2H).

LRMS (esi, positive) m/e 467.2 (M+1)

Compound 227:




N-(3-*S*-1-Cyclohexylmethyl-pyrrolidin-3-yl)-3-methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzamide

Prepared according to the procedure of Compound 215 using 1-cyclohexylmethyl-pyrrolidin-3-*S*-ylamine (60% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 8.78 (s, 1H), 8.31 (d, 1H), 8.23 (d, 1H), 8.20 (s, 1H), 7.51 (s, 1H), 7.48 (d, 1H), 4.37 (m, 1H), 3.96 (s, 3H), 2.76 (m, 1H), 2.56 (m, 1H), 2.42 (s, 3H), 2.38 (m, 1H), 2.25-2.06 (m, 3H), 1.77 (m, 3H), 1.61 (m, 3H), 1.40 (m, 1H), 1.19 (m, 3H), 0.83 (m, 2H).

LRMS (esi, positive) m/e 467.3 (M+1)

Compound 228:



N-(3-S-1-Benzyl-pyrrolidin-3-yl)-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-3-trifluoromethoxy-benzamide

Step 1: 4-Amino-3-trifluoromethoxy-benzoic acid methyl ester. To a stirred solution of 4-amino-3-trifluoromethoxy-benzoic acid (1.2 g, 5.4 mmol) in 16 mL of 4:1 THF:MeOH at 0 °C was added TMS-diazomethane (2M solution in hexane, 6 mL, 12 mmol) dropwise and conversion was monitored by TLC in 2/3 EtOAc/hexane. When complete, the reaction was concentrated to a white solid corresponding to the methyl ester (1.27 g, quantitative).

Step 2: 4-(4-Nitro-phenoxy-carbonylamino)-3-trifluoromethoxy-benzoic acid methyl ester. To a stirred solution of 4-amino-3-trifluoromethoxy-benzoic acid methyl ester (1.38 gm, 5.9 mmol) in 18 mL CH₂Cl₂ at 0 °C under nitrogen was added pyridine (521 µL, 6.4 mmol) followed by p-nitrophenylchloroformate (1.18 gm, 5.9 mmol). After 4 hours at 0 °C the reaction was diluted to 60 mL with CH₂Cl₂ and washed 2 x 60 mL with 2N HCl, 1 x 60 mL with H₂O and 1 x 60 mL with saturated NaCl. The organics were dried (MgSO₄), filtered and concentrated to a white solid corresponding to the carbamate (2.1 gm, 89 %).

Step 3: 4-[3-(5-Methyl-pyrazin-2-yl)-ureido]-3-trifluoromethoxy-benzoic acid methyl ester. To a stirred solution of 4-(4-nitro-phenoxy-carbonylamino)-3-trifluoromethoxy-benzoic acid methyl ester (400 mg, 1 mmol) in 1 mL of NMP in a capped reaction vial at room temperature was added 2-amino-5-methyl-pyrazine (109 mg, 1 mmol) and the solution heated to 90 °C for 6 hours. After cooling to room temperature, the solution was diluted to 30 mL with EtOAc and washed 4 x 30 mL with 10% NaHCO₃ to remove the phenol by-product and 1 x 30 mL with saturated NaCl. The organics were dried (MgSO₄), filtered and concentrated. Trituration and filtration with EtOAc gave a beige solid corresponding to the urea ester (136 mg, 37%).


Step 4: 4-[3-(5-Methyl-pyrazin-2-yl)-ureido]-3-trifluoromethoxy-benzoic acid. To a stirred suspension of 4-[3-(5-methyl-pyrazin-2-yl)-ureido]-3-trifluoromethoxy-benzoic acid methyl ester (136 mg, 0.37 mmol) in 4 mL 3:1 MeOH:H₂O under nitrogen was added lithium hydroxide monohydrate (154 mg, 3.7 mmol), and the reaction heated to 65 °C. The reaction turned yellow/green and eventually became a solution. After stirring overnight, the reaction was cooled to RT and partially concentrated by rotovap to remove most of the MeOH. The residual suspension was neutralized with 2N HCl until pH=6. The reaction formed a flocculent precipitate which was filtered off with H₂O and dried overnight under high vacuum to give the acid as a white solid (102 mg, 78 %).

Step 5: To a stirred suspension of 4-[3-(5-methyl-pyrazin-2-yl)-ureido]-3-trifluoromethoxy-benzoic acid (102 mg, 0.29 mmol) in 2.9 mL NMP in a capped reaction vial at room temperature was added HBTU (109 mg, 0.29 mmol). After 15 minutes 3-*S*-1-Benzyl-pyrrolidin-3-ylamine dihydrochloride (prepared analogously to 1-Pyridin-2-ylmethyl-pyrrolidin-3-ylamine used in the synthesis of Compound 215) (73 mg, 0.29 mmol) was added followed by DIEA (200 μ L, 1.2 mmol). The reaction was stirred overnight and NMP was then removed under high vacuum at 70 $^{\circ}$ C. The residue was partitioned between 30 mL CH_2Cl_2 and 30 mL 10% Na_2CO_3 . The organics were isolated and washed 1 x 30 mL with saturated NaCl, dried (MgSO_4), filtered and concentrated to a yellow foam corresponding to the desired amide (103 mg, 70%).

^1H -NMR (400 MHz, $\text{CDCl}_3/\text{CD}_3\text{OD}$) δ 8.56 (d, 1H), 8.37 (br s, 1H), 8.08 (s, 1H), 7.84 (s, 1H), 7.66 (d, 1H), 7.40-7.28 (m, 6H), 4.73 (m, 1H), 3.68 (dd, 2H), 2.77 (dd, 2H), 2.53 (s, 3H), 2.43 (m, 1H), 2.37 (dd, 2H), 1.78 (m, 1H)

LRMS (esi, positive) m/e 514.9 ($M+1$)

Compound 229:



N-(1-Benzyl-piperidin-4-ylmethyl)-3-methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzamide


Step 1: *N*-benzylisonipecotamide. To a suspension of isonipecotamide (12.3 g, 96 mmol) in 200 mL of dichloromethane, was added, benzaldehyde (10.6 g, 100 mmol) and sodium triacetoxyborohydride (29.7 gm, 140 mmol) and the mixture was stirred at room temperature for 5 days. The thick white mixture was diluted with 100 mL water and extracted with EtOAc (2 x 20 mL). The aqueous phase was basified with 1*N* NaOH to pH > 12. The resulting white precipitate was collected by suction filtration. The white solid was subsequently taken up in 50 mL of EtOAc, and was washed with 20 mL of brine, then dried (MgSO_4), filtered and concentrated to give 10.84 g (50 %) of the desired product. ^1H -NMR (400 MHz, CDCl_3) δ 7.85 (m, 5H), 5.45 (s, 1H), 5.34 (s, 1H), 3.5 (s, 2H), 2.93 (d, J = 10.96 Hz, 2H), 2.16 (t, J = 12.13 Hz, 1H), 2.01 (t, J = 11.74 Hz, 2H), 1.87 (d, J = 12.52 Hz, 2H), 1.76 (q, J = 12.52 Hz, 2H).

Step 2: 4-Aminomethyl-1-benzyl piperidine. To a solution of *N*-benzylisonipecotamide (7.34g, 34 mmol) in 60 mL of anhydrous THF, was added, LiAlH_4 (1.9 g, 51 mmol) and the mixture was stirred at room temperature for 10 minutes followed by heating to reflux for 3 h. The reaction was quenched by addition of 100 mL sat'd. sodium potassium tartrate, and was extracted with EtOAc (3 x 50 mL). The combined extracts were washed with 20 mL water and 20 mL brine then dried over MgSO_4 , filtered and concentrated to give the desired product. ^1H -NMR (400 MHz, CDCl_3) δ 7.13 (m, 5H), 3.42 (s, 2H), 2.91 (m, 2H), 2.59 (m, 2H), 1.97 (m, 2H), 1.62 (m, 2H), 2.21 (m, 5H).

Step 3: *N*-(1-Benzyl-piperidin-4-ylmethyl)-3-methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzamide. Prepared from 4-aminomethyl-1-benzyl piperidine, according to the procedure of Compound 208:

¹H-NMR (400 MHz, *d*₆-DMSO) δ 10.1 (s, 2H), 8.81 (s, 1H), 8.48 (s, 1H), 8.26 (m, 2H), 7.52 (m, 7H), 3.97 (s, 5H), 3.33 (s, 3H), 3.18 (s, 3H), 2.43 (s, 3H), 1.75 (s, 3H), 1.44 (s, 2H). MS APCI-pos, M+1 = 489.1.

Compound 230:



N-[3-*S*-1-(4-Fluoro-benzyl)-pyrrolidin-3-yl]-3-methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzamide


Step 1: (3*S*)-3-(tert-Butoxycarbonylamino)-1-(4-fluoro-benzyl)-pyrrolidine. To a solution of (3*S*)-(+)-3-(tert-butoxycarbonylamino)pyrrolidine (410 mg, 2.20 mmol), in 22 mL EtOH, was added, (288 μL, 2.31 mmol) of 4-fluorobenzyl bromide, and (788 mg, 2.42 mmol) of finely powdered cesium carbonate. The stirred reaction mixture was heated at 80 °C, under nitrogen for 3h, after which time, TLC indicated the reaction was complete. The reaction was then concentrated *in vacuo* to about 5 mL, and was then diluted with 30 mL of EtOAc, and washed with 20 mL of 5% NH₄OH. The aqueous fraction was extracted with diethyl ether (3 x 20 mL). The combined organics were washed with 20 mL of brine, dried (MgSO₄), filtered and concentrated. The crude white solid was triturated with 1:1 ether-hexane to give 501 mg (77%) of the desired product as a white solid. ¹H-NMR (400 MHz, CDCl₃) δ 7.2-7.3 (m, 2H), 6.9-7.1 (m, 2H), 4.85 (br. s, 1H, exchanges), 4.18 (br. s, 1H), 3.55 (s, 2H), 2.65 (br. m, 1H), 2.58 (m, 1H), 2.50 (m, 1H), 2.2-2.4 (m, 2H), 1.5-1.7, (m, 1H), 1.4 (s, 9H).

Step 2: (3*S*)-3-Amino-1-(4-fluoro-benzyl)-pyrrolidine. A solution of (3*S*)-3-(tert-butoxycarbonylamino)-1-(4-fluoro-benzyl)-pyrrolidine (400 mg, 1.36 mmol) was stirred in 15 mL of formic acid at room temperature. After 3h, the clear colorless solution was concentrated to dryness and the residue was taken-up in 20 mL of EtOAc and washed with 10 mL of 5% NH₄OH, followed by 10 mL of brine. The solution was then dried over MgSO₄, filtered and concentrated to give 240 mg (91%) of product, as a yellow oil. ¹H-NMR (400 MHz, CDCl₃) δ 7.28 (m, 2H), 6.99 (m, 2H), 3.56 (d, *J* = 6.2 Hz, 2H), 3.47-3.53 (m, 1H), 2.67-2.71 (m, 2H), 2.42-2.48 (m, 1H), 2.28-2.31 (m, 1H), 2.21-2.28 (m, 1H), 1.59 (s, 2H, NH₂), 1.43-1.52 (m, 1H).

Step 3: *N*-[(3*S*)-1-(4-Fluoro-benzyl)-pyrrolidin-3-yl]-3-methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzamide. Prepared from (3*S*)-3-amino-1-(4-fluoro-benzyl)-pyrrolidine, according to the procedure

of Compound 208. ¹H-NMR (400 MHz, CDCl₃) δ 10.8 (s, 2H), 8.8 (s, 1H), 8.38 (s, 1H), 8.27 (s, 1H), 8.24 (s, 2H), 7.51 (s, 1H), 7.49 (s, 1H), 7.37 (s, 2H), 7.15 (m, 2H), 4.4 (br.s, 1H), 3.98 (s, 3H), 3.6 (br.s, 2H), 3.06 (br.s, 1H), 2.65 (br.s, 1H), 2.81 (br.s, 1H), 2.43 (s, 3H), 2.16 (s, 1H), 1.83 (s, 1H). MS apci-pos M+1 = 479.2.

Compound 231:



3-Methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzoic acid 3-S-1-benzyl-pyrrolidin-3-yl ester

Step 1: 3-Methoxy-4-nitro-benzoyl chloride. Thionyl chloride 3.7 mL, (50 mmol) was added drop wise, at room temperature, to a stirred solution of 3-methoxy-4-nitro-benzoic acid (1.0 g, 5.07 mmol) in 15 mL of dioxane under a nitrogen atmosphere. After the addition was complete, the reaction was allowed to stir at room temperature for 1 hr. The reaction flask was then fitted with a distillation head, and excess thionyl chloride and about ½ of the solvent was removed by distillation in a 120 °C oil bath. The remaining solvent was removed by rotary evaporation, and the residue was taken up in 20 mL toluene and again about ½ of the solvent was removed by distillation. The remaining solution was then cooled to 0 °C, and the white solid which precipitated, was collected by suction filtration, and rinsed with 1-1 Et₂O-hexane. ¹H-NMR (400 MHz, CDCl₃) δ 7.87 (d, *J* = 7.83 Hz, 1H), 7.82 (d, *J* = 7.83 Hz, 1H), 7.77 (s, 1H), 4.04 (s, 3H).

Step 2: (3S)-1-Benzyl-pyrrolidin-3-ol. A solution of (3S)-3-hydroxypyrrolidine (2.0 g, 25 mmol) in 50 mL CH₂Cl₂, under a nitrogen atmosphere, was cooled to 0 °C and benzaldehyde (2.92 g, 27.5 mmol) was added, followed by 1g of powdered 4A molecular sieves. Sodium triacetoxyborohydride (7.4 gm, 35 mmol) was added in several portions over 30 min, and the reaction was allowed to stir at room temperature for 18h. The reaction was again cooled to 0 °C and quenched by addition of 10 mL methanol, followed by 5 mL of 1N HCl. The molecular sieve solids were removed by filtration through a glass fiber paper, and the filtrate was extracted with 20 mL diethyl ether. The organic phase was discarded, and the aqueous phase was first basified by addition of conc. ammonium hydroxide, and then extracted with EtOAc (3 x 20 mL). The combined extracts were washed with 20 mL brine, dried (MgSO₄), filtered, and concentrated *in vacuo* to furnish 3.2 g (73%) of clear yellow oil, which required no further purification. ¹H-NMR (400 MHz, CDCl₃) δ 7.2-7.4 (m, 5H), 4.33 (m, 1H), 3.63, (s, 2H), 2.83-3.89 (m, 1H), 2.67 (d, *J* = 9.9 Hz, 1H), 2.53-2.55 (m, 1H), 2.23-2.34 (m, 1H), 2.14-2.20 (m, 2H), 1.70-1.77 (m, 1H).


Step 3: 3-Methoxy-4-nitro-benzoic acid (3S)-1-benzyl-pyrrolidin-3-yl ester. A solution of 3-methoxy-4-nitro-benzoyl chloride (608 mg, 2.82 mmol) in 10 mL of CH₂Cl₂ was added to stirred solution of 500

mg (2.82 mmol) of (3S)-1-benzyl-pyrrolidin-3-ol and 1 mL of pyridine in 15 mL CH₂Cl₂ at room temperature. After stirring for 18 h, the reaction mixture was diluted with 20 mL of saturated aqueous NaHCO₃ and extracted with CH₂Cl₂ (2 x 10 mL). The combined extracts were washed with 20 mL of brine, dried over MgSO₄, filtered, and concentrated *in vacuo* to furnish 780 mg (71%) of the desired ester as a yellow oil, which solidified after drying under high vacuum. The resulting solid was further purified by trituration with 1:1 ether-hexane. ¹H-NMR (400 MHz, CDCl₃) δ 7.83 (d, *J* = 7.83 Hz, 1H), 7.74 (s, 1H), 7.68 (d, *J* = 7.83 Hz, 1H), 7.2-7.4 (m, 5H), 5.44 (br.m, 1H), 4.02 (s, 2H), 3.69 (dd, *J* = 30 Hz, *J* = 13 Hz, 1H), 2.8-3.0 (m, 2H), 2.5-2.6 (m, 1H), 2.3-2.45 (m, 1H), 2.0-2.1 (m, 1H)

Step 4: 4-Amino-3-methoxy-benzoic acid (3S)-1-benzyl-pyrrolidin-3-yl ester. The aniline was prepared by nickel boride reduction of 3-methoxy-4-nitro-benzoic acid (3S)-1-benzyl-pyrrolidin-3-yl ester analogously to the preparation of 4-[(Benzyl-methyl-amino)-methyl]-2-methoxy-phenylamine used in the synthesis of Compound 340. ¹H-NMR (400 MHz, CDCl₃) δ 7.54 (d, *J* = 8.21 Hz, 1H), 7.44 (s, 1H), 7.2-7.4 (m, 5H) 6.65 (d, *J* = 8.21 Hz, 1H), 5.38 (br.m, 1H), 4.22 (s, 1H), 3.90 (s, 3H), 3.69 (q, *J* = 24.63 Hz, 1H), 3.0 (m, 1H), 2.7-2.9 (m, 2H), 2.5-2.6 (m, 1H), 2.3-2.4 (m, 1H), 1.9-2.1 (m, 1H). MS apci-pos M+1=327.2.

Step 5: 3-Methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzoic acid (3S)-1-benzyl-pyrrolidin-3-yl ester. Prepared from 4-amino-3-methoxy-benzoic acid (3S)-1-benzyl-pyrrolidin-3-yl ester according to the procedure of Compound 208: ¹H-NMR (400 MHz, CDCl₃) δ 10.15 (s, 2H), 8.8 (s, 1H), 8.35 (d, *J* = 8.61 Hz, 1H), 8.25 (s, 1H), 7.58 (d, *J* = 8.61 Hz, 1H), 7.5 (s, 1H), 7.30- 7.35 (m, 4H), 7.26 -7.24 (m, 1H), 5.28 (s, 1H), 3.99 (s, 3H), 3.62 (s, 2H), 2.82 (m, 1H), 2.74 (m, 1H), 2.67 (m, 1H), 2.44 (s, 3H), 2.28 (m, 1H), 1.91 (m, 1H). MS apci-pos M+1 = 462.2.

Compound 232:



N-(3-S-1-Benzyl-pyrrolidin-3-yl)-3-methoxy-4-[3-(5-trifluoromethyl-pyrazin-2-yl)-ureido]-benzamide

Step 1: 3-Methoxy-4-[3-(5-trifluoromethyl-pyrazin-2-yl)-ureido]-benzoic acid methyl ester. To a stirred solution of 5-trifluoromethylaminopyrazine (326 mg, 2 mmol) at room temperature in NMP (2 mL) in a capped reaction vial was added 4-methoxy-3-(4-nitrophenylcarbonylamino)-benzoic acid methyl ester (692 mg, 2 mmol) and the solution was heated to 85° C for 6 hours. The reaction was cooled to RT and triturated with EtOAc (5 mL) and the tan solid that formed was isolated by filtration and rinsing with EtOAc (213 mg, 28%).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 10.78 (s, 1H), 10.16 (br s, 1H), 9.07 (s, 1H), 8.84 (s, 1H), 8.37 (d, 1H), 7.63 (d, 1H), 7.56 (s, 1H), 4.01 (s, 3H), 3.83 (s, 3H)


Step 2: 3-Methoxy-4-[3-(5-trifluoromethyl-pyrazin-2-yl)-ureido]-benzoic acid. To a stirred solution of 3-methoxy-4-[3-(5-trifluoromethyl-pyrazin-2-yl)-ureido]-benzoic acid methyl ester (213 mg, 0.575 mmol) in 5.75 mL 3:1 MeOH:H₂O at room temperature under nitrogen was added lithium hydroxide monohydrate (240 mg, 5.8 mmol) and the reaction warmed to 65 °C. After reaching temperature, the suspension gradually became a bright yellow solution. After about 4 hours a precipitate formed but the reaction was continued overnight. After cooling to RT, MeOH was removed by rotovap and the aqueous suspension neutralized to pH=5 with concentrated HCl. As pH=5 was approached, the suspension turned from yellow to white. The suspension was then filtered through paper on a ceramic funnel. When most of the H₂O was removed the residue was dried under high vacuum in a dessicator overnight (133 mg, 55%).

Step 3: To a stirred solution of 3-methoxy-4-[3-(5-trifluoromethyl-pyrazin-2-yl)-ureido]-benzoic acid (35 mg, 0.1 mmol) in 0.5 mL NMP in a capped reaction vial at RT was added HBTU (41 mg, 0.11 mmol) and the activation stirred for 15 min. 1-Benzyl-pyrrolidin-3-ylamine dihydrochloride (prepared analogously to 1-Pyridin-2-ylmethyl-pyrrolidin-3-ylamine used in the synthesis of Compound 215) (crude 0.1 mmol) was then added as a solution in 0.5 mL NMP with DIEA (69 µL, 0.4 mmol). After stirring overnight at RT, NMP was removed by distillation under high vacuum at 70° C. The residue was dissolved in 25 mL CH₂Cl₂ with a small amount of MeOH and washed 2 x 25 mL with 10% Na₂CO₃. The organics were dried (MgSO₄), filtered and concentrated to a residue that was chromatographed on a biotage 12S column with 5/95 MeOH/CH₂Cl₂. This material was concentrated to dryness and triturated/filtered with Et₂O to give pure product as a white solid (9.5 mg, 19%).

¹H-NMR (400 MHz, CDCl₃) δ 11.53 (s, 1H), 9.86 (br s, 1H), 8.57 (s, 1H), 8.40 (d, 1H), 8.12 (s, 1H), 7.56 (s, 1H), 7.46 (m, 2H), 7.25 (m, 4H), 4.79 (m, 1H), 4.01 (s, 3H), 3.73 (dd, 2H), 3.22 (m, 2H), 2.76 (m, 1H), 2.47 (m, 2H), 1.76 (m, 1H)

LRMS (esi, positive) m/e 515.1 (M+1)

Compound 233:



5-Methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-N-(2-pyridin-2-yl-ethyl)-2-trifluoromethyl-benzamide

Step 1: Mixture of 5-Fluoro-4-nitro-2-trifluoromethyl-benzoic acid and 5-Fluoro-3-nitro-2-trifluoromethyl-benzoic acid. To a stirred solution of 3-fluoro-6-trifluoromethylbenzoic acid (3.58 gm, 17.2 mmol) in 17 mL concentrated H₂SO₄ at 0° C was carefully added 70% HNO₃ (17 mL) dropwise. The reaction was heated to 100° C overnight and was then cooled to room temperature and poured into 250 mL of H₂O/ice with stirring. EtOAc (250 mL) was added and the mixture stirred. The layers were separated and the organics washed 1 x 250 mL with H₂O and 1 x 250 mL with saturated NaCl. The

organics were dried (MgSO_4), filtered and concentrated to an oil which solidified upon standing. The solid was a 1:1 mixture of regioisomers and was carried on crude.

Step 2: Mixture of 5-Fluoro-4-nitro-2-trifluoromethyl-benzoic acid methyl ester and 5-Fluoro-3-nitro-2-trifluoromethyl-benzoic acid methyl ester. To a stirred solution of the nitro acids (crude, 17.2 mmol) in 60 mL of 4:1 THF:MeOH at 0 °C was added 2M TMS-diazomethane in hexane dropwise until a yellow color persisted. After 30 minutes the reaction was concentrated to a crude oil and used directly in the next step.

Step 3: Mixture of 5-Methoxy-4-nitro-2-trifluoromethyl-benzoic acid methyl ester and 5-Methoxy-3-nitro-2-trifluoromethyl-benzoic acid methyl ester. To a stirred solution of the fluoro nitro esters (crude, 17.2 mmol) in 22 mL MeOH at room temperature was added 150 mL of 0.5 M sodium methoxide in MeOH. The red solution was stirred for 30 minutes and then partitioned between CH_2Cl_2 (500 mL) and H_2O (500 mL). The organics were washed 2 x 500 mL with H_2O , dried (MgSO_4), filtered and concentrated to a white solid.


Step 4: 5-Methoxy-4-amino-2-trifluoromethyl-benzoic acid methyl ester. To a stirred solution of the methoxy nitro esters (crude, 17.2 mmol) in 172 mL MeOH at room temperature purged with nitrogen was carefully added 10% Pd on C (1 gm). The suspension was put through a vacuum/purge cycle three times with hydrogen gas and then held under 1 atmosphere of hydrogen. After stirring overnight the suspension was filtered through GF/F filter paper and concentrated to a clear oil. This material was loaded directly onto a Biotage 40M column with CH_2Cl_2 and eluted first with 9/1 hexane/EtOAc to elute the undesired hi Rf regioisomer and then with 85/15 hexane/EtOAc to elute the desired lower Rf regioisomer. After concentration, the lower Rf isomer solidified to a clear solid (1.75 gm, 41%). ^1H -NMR (400 MHz, CDCl_3) δ 7.33 (s, 1H), 6.98 (s, 1H), 3.92 (s, 3H), 3.85 (s, 3H)

Step 5: 5-Methoxy-4-(4-nitro-phenoxy-carbonylamino)-2-trifluoromethyl-benzoic acid methyl ester. To a stirred solution of 5-methoxy-4-amino-2-trifluoromethyl-benzoic acid methyl ester (552 mg, 2.22 mmol) in 6.6 mL CH_2Cl_2 at 0° C under nitrogen was added pyridine (180 μL , 2.22 mmol) followed by p-nitrophenyl chloroformate (448 mg, 2.22 mmol) as a solid. After stirring for 1 hour, the reaction was diluted to 30 mL with CH_2Cl_2 and washed 2 x 30 mL with 2N HCl and 1 x 30 mL with H_2O . The organics were dried (MgSO_4), filtered and concentrated to the p-nitrophenyl carbamate which was isolated as a white foam (878 mg, 96%). ^1H -NMR (400 MHz, CDCl_3) δ 8.57 (br s, 1H), 8.30 (d, 2H), 7.77 (br s, 1H), 7.40 (d, 2H), 7.37 (s, 1H), 4.03 (s, 3H), 3.94 (s, 3H)

Step 6: 5-Methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-2-trifluoromethyl-benzoic acid methyl ester. To a stirred solution of 5-methoxy-4-(4-nitro-phenoxy-carbonylamino)-2-trifluoromethyl-benzoic acid methyl ester (878 mg, 2.1 mmol) in 4.2 mL NMP at room temperature under nitrogen was added 2-amino-5-methylpyrazine (232 mg, 2.1 mmol) and the reaction heated to 85 °C. After six hours the reaction was cooled to room temperature and a precipitate formed. The precipitate was triturated with EtOAc (25 mL) and the urea isolated by filtration as a tan solid (470 mg, 58%).

LRMS (esi, positive) m/e 475.1 (M+1)

Compound 234:



3-(3-Dimethylamino-propoxy)-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzoic acid

Step 1: 3-(3-Dimethylamino-propoxy)-4-nitro-benzoic acid methyl ester. To a stirred, cooled (about 0°C) solution of 3-hydroxy-4-nitro benzoic acid methyl ester. (394 mg, 2.0 mmol), triphenyl phosphine (525 mg, 2.0 mmol), and 3-(dimethylamino)-propanol (237 µL, 2.0 mmol) in dry tetrahydrofuran (5 mL) was added diisopropyl azodicarboxylate (394 µL, 2.0 mmol in 1 mL of tetrahydrofuran). After stirring for 12 hours, the reaction was diluted with hydrochloric acid (30 mL of a 1M solution) and extracted with ethyl acetate (2x 50mL). The aqueous solution was basified with 10% aqueous sodium carbonate (50 mL) and the product was extracted with ethyl acetate (3 x 100 mL). The ethyl acetate


was washed with brine (1 x 30 mL), then dried (MgSO₄), and filtered. The filtered solution was concentrated under reduced pressure to provide the desired crude product (86 % yield). ¹H-NMR (400 MHz, CDCl₃) δ 7.81 (d, 1H), 7.78 (s, 1H), 7.65 (d, 1H), 4.23 (t, 2H), 3.91 (s, 3H), 2.44 (t, 2H), 2.22 (s, 6H), 2.05 (m, 2H).

Step 2: 4-Amino-3-(3-dimethylamino-propoxy)-benzoic acid methyl ester. To a stirred, cooled (about 0°C) solution of 3-(3-dimethylamino-propoxy)-4-nitro-benzoic acid methyl ester (282 mg, 1.0 mmol) in methanol (2 mL) and saturated aqueous ammonium chloride (1mL) was added zinc dust (2.0 mmol). After stirring for 12 hours, the reaction was diluted with 30 mL of ethyl acetate and washed with 30 mL of 10% aqueous sodium carbonate (2 x 30 mL), brine (1 x 30 mL), then dried (MgSO₄), and filtered. The filtered solution was concentrated under reduced pressure to yield the desired product as a tan solid (88% yield). ¹H-NMR (400 MHz, CDCl₃) δ 7.52 (d, 1H), 7.48 (s, 1H), 7.25 (s, 1H), 6.65 (d, 1H), 4.29 (br s, 1H), 4.09 (t, 2H), 3.85 (s, 3H), 2.49 (t, 2H), 2.25 (s, 6H), 2.00 (m, 2H).

Step 3: 3-(3-Dimethylamino-propoxy)-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzoic acid methyl ester. To a stirred, cooled (about 0°C) solution of 4-amino-3-(3-dimethylamino-propoxy)-benzoic acid methyl ester (252 mg, 1.0 mmol) in toluene (3.0 mL) was added triethylamine (139 µL, 1.0 mmol) and triphosgene (98 mg, 0.33 mmol). After stirring for 30 minutes, 5-methyl-2-amino pyrazine (109 mg, 1.0 mmol) was added and the reaction was heated to 65 degrees C. The reaction was allowed to cool to room temperature, then diluted with ethyl acetate (50 mL) and water (50 mL). A precipitate formed which was filtered and dried under reduced pressure to yield the desired material as a white solid (40% yield). ¹H-NMR (400 MHz, d₆-DMSO) δ 8.62 (br s, 1H), 8.41 (d, 1H), 8.19 (s, 1H), 7.59 (d, 1H), 7.49 (s, 1H), 4.15 (t, 2H), 3.81 (s, 3H), 2.42 (m, 5H), 2.18 (s, 6H), 2.01 (m, 2H).

Step 4: 3-(3-Dimethylamino-propoxy)-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzoic acid. To a stirred solution of 3-(3-dimethylamino-propoxy)-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzoic acid methyl ester (1.0 gm; 3.3 mmol) in methanol (25 mL) was added lithium hydroxide (5 mL of a 2M aqueous solution). The reaction was heated to 60 degrees and stirred for 12 hours. The reaction was allowed to cool to room temperature and the pH was adjusted to 5.5 with 1N hydrochloric acid. A precipitate formed which was filtered and dried under reduced pressure to yield the desired product as a tan solid (52% yield). ¹H-NMR (400 MHz, d₆-DMSO) δ 8.62 (br s, 1H), 8.39 (d, 1H), 8.21 (s, 1H), 7.59 (d, 1H), 7.43 (s, 1H), 4.15 (t, 2H), 2.42 (m, 5H), 2.18 (s, 6H), 2.01 (m, 2H).

Compound 235:



4-[3-(5-Methyl-pyrazin-2-yl)-ureido]-3-(pyridin-3-ylmethoxy)-benzoic acid


Step 1: 4-Nitro-3-(pyridin-3-ylmethoxy)-benzoic acid methyl ester. To a stirred, cooled (about 0°C) solution of 3-hydroxy-4-nitro benzoic acid methyl ester (394 mg, 2.0 mmol), triphenyl phosphine (525 mg, 2.0 mmol), and 3-pyridylcarbinol (194 μ L, 2.0 mmol) in dry tetrahydrofuran (5 mL) was added diisopropyl azodicarboxylate (394 μ L, 2.0 mmol in 1 mL of tetrahydrofuran). After stirring for 12 hours, the reaction was diluted with hydrochloric acid (30 mL of a 1M solution) and extracted with ethyl acetate (2 x 50 mL). The aqueous solution was basified with 10% aqueous sodium carbonate (50 mL) and the product was extracted with ethyl acetate (3 x 100 mL). The ethyl acetate was washed with brine (1 x 30 mL), then dried (MgSO₄), and filtered. The filtered solution was concentrated under reduced pressure to provide the desired crude product as an off white solid (76 % yield). ¹H-NMR (400 MHz, CDCl₃) δ 8.71 (s, 1H), 8.62 (d, 1H), 7.88 (m, 3H), 7.79 (d, 1H), 7.39 (m, 1H), 5.31 (s, 2H), 3.98 (s, 3H).

Step 2: 4-Amino-3-(pyridin-3-ylmethoxy)-benzoic acid methyl ester. To a stirred, cooled (about 0°C) solution of 4-nitro-3-(pyridin-3-ylmethoxy)-benzoic acid methyl ester (288 mg, 1.0 mmol) in methanol (2 mL) and saturated aqueous ammonium chloride (1 mL) was added zinc dust (131 mg, 2.0 mmol). After stirring for 12 hours, the reaction was diluted with 30 mL of ethyl acetate and washed with 30 mL of 10% aqueous sodium carbonate (2 x 30 mL), brine (1 x 30 mL), then dried (MgSO₄), and filtered. The filtered solution was concentrated under reduced pressure to yield the desired product as a yellow solid (97% yield). ¹H-NMR (400 MHz, CDCl₃) δ 8.78 (s, 1H), 8.61 (d, 1H), 7.79 (d, 1H), 7.59 (m, 2H), 7.39 (m, 1H), 6.71 (d, 1H), 5.18 (s, 2H), 4.28 (br s, 2H), 3.85 (s, 3H).

Step 3: 4-[3-(5-Methyl-pyrazin-2-yl)-ureido]-3-(pyridin-3-ylmethoxy)-benzoic acid methyl ester. To a stirred, cooled (about 0°C) solution of 4-amino-3-(pyridin-3-ylmethoxy)-benzoic acid methyl ester (258 mg, 1.0 mmol) in toluene (3.0 mL) was added triethylamine (139 μ L, 1.0 mmol) and triphosgene (98 mg, 0.33 mmol). After stirring for 30 minutes, 5-methyl-2-amino pyrazine (109 mg, 1.0 mmol) was added and the reaction was heated to 65 degrees C. The reaction was allowed to cool to room temperature, then diluted with ethyl acetate (50 mL) and water (50 mL). A precipitate formed which was filtered and dried under reduced pressure to yield the desired material as a white solid (47% yield). ¹H-NMR (400 MHz, d₆-DMSO) δ 10.29 (s, 1H), 8.79 (s, 1H), 8.68 (d, 1H), 8.59 (br s, 1H), 8.48 (d, 1H), 7.70 (s, 1H), 7.62 (d, 1H), 7.51 (m, 1H), 5.32 (s, 2H), 3.88 (s, 3H), 2.32 (s, 3H).

Step 4: 4-[3-(5-Methyl-pyrazin-2-yl)-ureido]-3-(pyridin-3-ylmethoxy)-benzoic acid. To a stirred solution of 4-[3-(5-methyl-pyrazin-2-yl)-ureido]-3-(pyridin-3-ylmethoxy)-benzoic acid methyl ester (1.3 gm; 3.3 mmol) in methanol (25 mL) was added lithium hydroxide (5 mL of a 2M aqueous solution). The reaction was heated to 60 degrees and stirred for 12 hours. The reaction was allowed to cool to room temperature and the pH was adjusted to 5.5 with 1N hydrochloric acid. A precipitate formed which was filtered and dried under reduced pressure to yield the desired product as a tan solid (90% yield). ¹H-NMR (400 MHz, d₆-DMSO) δ 10.29 (s, 1H), 8.79 (s, 1H), 8.68 (d, 1H), 8.59 (br s, 1H), 8.48 (d, 1H), 7.70 (s, 1H), 7.62 (d, 1H), 7.51 (m, 1H), 5.32 (s, 2H), 2.32 (s, 3H).

Compound 236:




3-(3-Dimethylamino-propoxy)-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzamide

Prepared according to the procedure of Compound 208 using ammonia (33% yield)

¹H-NMR (400 MHz, *d*₆-DMSO) δ 10.18 (s, 1H), 8.56 (br s, 1H), 8.20 (d, 1H), 8.11 (s, 1H), 7.43 (d, 1H), 7.40 (s, 1H), 4.03 (m, 2H), 2.40 (m, 2H), 2.33 (s, 3H), 2.05 (s, 6H), 1.91 (m, 2H)

LRLCMS (esi, positive) *m/e* 374.2 (M+1)

Compound 237:




3-(3-Dimethylamino-propoxy)-N,N-dimethyl-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzamide

Prepared according to the procedure of Compound 208 using dimethylamine (97% yield)

¹H-NMR (400 MHz, CDCl₃) δ 11.30 (br s, 1H), 8.39 (d, 1H), 8.29 (s, 1H), 8.20 (s, 1H), 7.78 (br s, 1H), 7.07 (s, 1H), 7.04 (d, 1H), 4.16 (t, 2H), 2.55 (s, 3H), 2.53 (m, 2H), 2.26 (s, 6H), 2.10 (m, 2H)

LRMS (esi, positive) *m/e* 401.1 (M+1)

Compound 238:




N-Benzyl-3-(3-dimethylamino-propoxy)-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzamide

Prepared according to the procedure of Compound 208 using benzylamine (65% yield)

¹H-NMR (400 MHz, CDCl₃) δ 8.43 (d, 1H), 8.26 (s, 1H), 8.20 (s, 1H), 7.91 (br s, 1H), 7.56 (s, 1H), 7.38 (m, 4H), 7.31 (m, 1H), 6.41 (m, 1H), 4.64 (d, 2H), 4.20 (t, 2H), 2.53 (s, 3H), 2.52 (m, 2H), 2.24 (s, 6H), 2.14 (m, 2H)

LRMS (esi, positive) m/e 463.2 (M+1)

Compound 239:




N-Benzyl-3-(3-dimethylamino-propoxy)-N-methyl-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzamide

Prepared according to the procedure of Compound 208 using N-methyl benzylamine (68% yield).

¹H-NMR (400 MHz, CDCl₃) δ 8.37 (br s, 1H), 8.25 (s, 1H), 8.18 (s, 1H), 7.37 (m, 4H), 7.31 (m, 1H), 7.18 (m, 2H), 4.67 (br m, 2H), 4.07 (br m, 2H), 2.99 (br s, 3H), 2.53 (s, 3H), 2.50 (br m, 2H), 2.24 (s, 6H), 2.06 (br m, 2H)

LRMS (esi, positive) m/e 477.2 (M+1)

Compound 240:




3-(3-Dimethylamino-propoxy)-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-N-(2-morpholin-4-yl-ethyl)-benzamide

Prepared according to the procedure of Compound 208 using 2-morpholin-4-yl-ethylamine (69% yield).

¹H-NMR (400 MHz, CDCl₃) δ 8.44 (d, 1H), 8.31 (s, 1H), 8.21 (s, 1H), 7.90 (br s, 1H), 7.55 (s, 1H), 7.24 (d, 1H), 6.76 (m, 1H), 4.20 (t, 2H), 3.76 (m, 4H), 3.56 (m, 2H), 2.61 (m, 2H), 2.53 (m, 7H), 2.24 (s, 6H), 2.13 (m, 2H)

LRMS (esi, positive) m/e 486.2 (M+1)

Compound 241:




3-(3-Dimethylamino-propoxy)-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-N-[2-(1-methyl-pyrrolidin-2-yl)-ethyl]-benzamide

Prepared according to the procedure of Compound 208 using 2-(1-methyl-pyrrolidin-2-ylamine (68% yield).

¹H-NMR (400 MHz, CDCl₃) δ 11.35 (br s, 1H), 8.42 (s, 1H), 8.40 (s, 1H), 8.24 (s, 1H), 8.20 (s, 1H), 7.57 (s, 1H), 7.46 (br s, 1H), 7.20 (m, 1H), 4.19 (m, 2H), 3.76 (m, 1H), 3.44 (m, 1H), 3.15 (m, 1H), 2.54 (s, 3H), 2.42 (m, 1H), 2.37 (s, 3H), 2.25 (s, 6H), 2.22 (m, 1H), 2.13 (m, 2H), 1.90 (m, 2H), 1.77 (m, 2H)

LRMS (esi, positive) m/e 484.3 (M+1)

Compound 242:




N-(2-Dimethylamino-ethyl)-3-(3-dimethylamino-propoxy)-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzamide

Prepared according to the procedure of Compound 208 using 2-dimethylamino-ethylamine (37% yield).

¹H-NMR (400 MHz, CDCl₃) δ 11.38 (br s, 1H), 8.42 (d, 1H), 8.30 (s, 1H), 8.20 (s, 1H), 7.96 (br s, 1H), 7.53 (s, 1H), 7.27 (d, 1H), 6.79 (m, 1H), 4.20 (m, 2H), 3.53 (m, 2H), 2.52 (m, 2H), 2.52 (s, 3H), 2.26 (m, 2H), 2.24 (s, 6H), 2.22 (s, 6H), 2.12 (m, 2H)

LRMS (esi, positive) m/e 444.2 (M+1)

Compound 243:




N-(3-S-1-Benzyl-pyrrolidin-3-yl)-3-(3-dimethylamino-propoxy)-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzamide

Prepared according to the procedure of Compound 215 using 1-benzyl-pyrrolidin-3-S-ylamine (20% yield)

¹H-NMR (400 MHz, CDCl₃) δ 8.61 (br s, 1H), 8.45 (d, 1H), 8.20 (s, 1H), 8.16 (m, 1H), 7.51 (s, 1H), 7.39 (d, 1H), 7.31 (m, 4H), 7.16 (m, 1H), 4.72 (m, 1H), 4.18 (m, 2H), 3.72 (dd, 2H), 3.04 (m, 1H), 2.96 (m, 1H), 2.69 (m, 1H), 2.53 (m, 2H), 2.51 (s, 3H), 2.40 (m, 2H), 2.25 (s, 6H), 2.11 (m, 2H), 1.76 (m, 1H)

LRMS (esi, positive) m/e 532.2 (M+1)

Compound 244:




4-[3-(5-Methyl-pyrazin-2-yl)-ureido]-3-(pyridin-3-ylmethoxy)-benzamide

Prepared according to the procedure of Compound 208 using ammonia (99% yield).

¹H-NMR (400 MHz, d₆-DMSO) δ 10.27 (s, 1H), 8.82 (s, 1H), 8.66 (d, 1H), 8.60 (br s, 1H), 8.29 (d, 1H), 8.03 (m, 1H), 8.00 (s, 1H), 7.76 (s, 1H), 7.56 (d, 1H), 7.50 (m, 1H), 7.37 (br m, 1H), 7.26 (br s, 1H), 5.32 (s, 2H), 2.32 (s, 3H)

LRMS (apci, positive) m/e 379.1 (M+1)

Compound 245:




N-Methyl-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-3-(pyridin-3-ylmethoxy)-benzamide

Prepared according to the procedure of Compound 208 using methylamine (99% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 10.25 (s, 1H), 8.80 (s, 1H), 8.60 (m, 3H), 8.26 (d, 1H), 8.03 (d, 1H), 7.78 (s, 1H), 7.52 (m, 2H), 7.37 (m, 1H), 5.34 (s, 2H), 2.79 (d, 3H), 2.36 (s, 3H)

LRMS (apci, positive) *m/e* 393.2 (M+1)

Compound 246:




N,N-Dimethyl-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-3-(pyridin-3-ylmethoxy)-benzamide

Prepared according to the procedure of Compound 208 using dimethylamine (88% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 10.16 (s, 1H), 8.80 (s, 1H), 8.66 (d, 1H), 8.58 (s, 1H), 8.28 (d, 1H), 7.96 (d, 1H), 7.50 (m, 1H), 7.37 (br s, 1H), 7.25 (s, 1H), 7.03 (d, 1H), 5.30 (s, 2H), 2.95 (s, 6H), 2.33 (s, 3H)

LRMS (apci, positive) *m/e* 407.4 (M+1)

Compound 247:




N-Benzyl-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-3-(pyridin-3-ylmethoxy)-benzamide

Prepared according to the procedure of Compound 208 using benzylamine (41% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 10.23 (s, 1H), 9.05 (t, 1H), 8.82 (s, 1H), 8.67 (d, 1H), 8.58 (br s, 1H), 8.33 (d, 1H), 8.00 (d, 1H), 7.78 (s, 1H), 7.59 (d, 1H), 7.50 (m, 1H), 7.33 (m, 4H), 7.25 (m, 1H), 5.32 (s, 2H), 4.50 (d, 2H), 2.32 (s, 3H)

LRMS (apci, positive) *m/e* 469.1 (M+1)

Compound 248:




N-Benzyl-N-methyl-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-3-(pyridin-3-ylmethoxy)-benzamide

Prepared according to the procedure of Compound 208 using N-methyl benzylamine (71% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 10.16 (s, 1H), 8.76 (br s, 1H), 8.64 (d, 1H), 8.57 (m, 1H), 8.28 (br m, 1H), 7.95 (br m, 1H), 7.48 (m, 1H), 7.40-7.23 (br m, 7H), 7.06 (m, 1H), 5.23 (br m, 2H), 4.62 (br m, 2H), 2.88 (s, 3H), 2.33 (s, 3H)

LRMS (apci, positive) *m/e* 483.3 (M+1)

Compound 249:




4-[3-(5-Methyl-pyrazin-2-yl)-ureido]-N-(2-morpholin-4-yl-ethyl)-3-(pyridin-3-ylmethoxy)-benzamide

Prepared according to the procedure of Compound 208 using 2-morpholin-4-yl-ethylamine (99% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 10.27 (s, 1H), 8.81 (s, 1H), 8.66 (d, 1H), 8.61 (br s, 1H), 8.52 (t, 1H), 8.29 (d, 1H), 8.03 (d, 1H), 7.76 (s, 1H), 7.52 (m, 2H), 7.36 (m, 1H), 5.33 (s, 2H), 3.56 (m, 4H), 3.40 (m, 2H), 2.48 (m, 2H), 2.43 (m, 4H), 2.34 (s, 3H)

LRMS (apci, positive) *m/e* 492.4 (M+1)

Compound 250:




4-[3-(5-Methyl-pyrazin-2-yl)-ureido]-N-[2-(1-methyl-pyrrolidin-2-yl)-ethyl]-3-(pyridin-3-ylmethoxy)-benzamide

Prepared according to the procedure of Compound 208 using 2-(1-methyl-pyrrolidin-2-yl)-ethylamine (99% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 10.21 (s, 1H), 8.81 (s, 1H), 8.64 (d, 1H), 8.58 (br s, 1H), 8.49 (m, 1H), 8.33 (d, 1H), 8.00 (d, 1H), 7.70 (s, 1H), 7.52 (m, 2H), 7.34 (m, 1H), 5.33 (s, 2H), 3.28 (m, 2H), 2.95 (m, 1H), 2.36 (s, 3H), 2.21 (s, 3H), 2.04 (m, 2H), 1.90 (m, 2H), 1.62 (m, 2H), 1.44 (m, 2H)

LRMS (apci, positive) *m/e* 490.3 (M+1)

Compound 251:




N-(2-Dimethylamino-ethyl)-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-3-(pyridin-3-ylmethoxy)-benzamide

Prepared according to the procedure of Compound 208 using 2-dimethylamino-ethylamine (99% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 10.24 (s, 1H), 8.82 (s, 1H), 8.66 (d, 1H), 8.60 (br s, 1H), 8.43 (t, 1H), 8.30 (d, 1H), 8.02 (d, 1H), 7.74 (s, 1H), 7.52 (m, 2H), 7.35 (br m, 1H), 5.32 (s, 2H), 3.36 (m, 2H), 2.39 (m, 2H), 2.33 (s, 3H), 2.18 (s, 6H)

LRMS (apci, positive) *m/e* 450.2 (M+1)

Compound 252:




N-(3-S-1-Benzyl-pyrrolidin-3-yl)-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-3-(pyridin-3-ylmethoxy)-benzamide

Prepared according to the procedure of Compound 215 using 1-benzyl-pyrrolidin-3-S-ylamine (99% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 10.26 (s, 1H), 8.82 (s, 1H), 8.66 (d, 1H), 8.60 (br s, 1H), 8.47 (d, 1H), 8.28 (d, 1H), 8.02 (d, 1H), 7.74 (s, 1H), 7.55 (d, 1H), 7.51 (m, 1H), 7.32 (m, 4H), 7.24 (m, 1H), 5.31 (s, 2H), 4.40 (m, 1H), 3.60 (s, 2H), 2.80 (m, 1H), 2.64 (m, 1H), 2.45 (m, 2H), 2.34 (s, 3H), 2.15 (m, 1H), 1.85 (m, 1H)

LRMS (apci, positive) *m/e* 538.2 (M+1)

Compound 253:



1-[2-(3-Dimethylamino-propoxy)-5-methyl-phenyl]-3-pyrazin-2-yl-urea

Step 1: (2-Hydroxy-5-methyl-phenyl)-carbamic acid *tert*-butyl ester. To a stirred solution of 2-amino-4-methyl-phenol (6.15 gm; 50 mmol) in dioxane (100 mL) was added carbonic acid di-*tert*-butyl ester (9.8 gm; 45 mmol) followed by sodium bicarbonate (12.6 gm; 150 mmol in 75 mL of H₂O). After stirring for 8 hours, the reaction was diluted with 100 mL of ethyl acetate and washed with 1N aqueous hydrochloric acid (2 x 100 mL), saturated aqueous sodium bicarbonate (1 x 100 mL), and brine (100 mL), then dried (MgSO₄), and filtered. The filtered solution was concentrated under reduced pressure to provide the desired (2-Hydroxy-5-methyl-phenyl)-carbamic acid *tert*-butyl ester as a brown solid (95% yield).

Step 2: [2-(3-Dimethylamino-propoxy)-5-methyl-phenyl]-carbamic acid *tert*-butyl ester. To a stirred, cooled (about 0°C) solution of (2-hydroxy-5-methyl-phenyl)-carbamic acid *tert*-butyl ester (447mg, 2.0 mmol), triphenyl phosphine (525 mg, 2.0 mmol), and 3-(dimethylamino)-1-propanol (237 μL, 2.0

mmol) in dry tetrahydrofuran (5 mL) was added diisopropyl azodicarboxylate (394 μ L, 2.0 mmol) in 1 mL of tetrahydrofuran). After stirring for 12 hours, the reaction was diluted with 30 mL of ethyl acetate and washed with 30 mL of 10% aqueous sodium carbonate (2 x 30 mL), brine (1 x 30 mL), then dried (MgSO_4), and filtered. The filtered solution was concentrated under reduced pressure to provide the desired crude product.


Step 3: 2-(3-Dimethylamino-propoxy)-5-methyl-phenylamine. To a stirred solution of [2-(3-dimethylamino-propoxy)-5-methyl-phenyl]-carbamic acid tert-butyl ester (617 mg, 2.0 mmol) in 5 mL of dioxane was added hydrochloric acid (2 mL; 4N in dioxane). After stirring for 12 hours, the reaction was diluted with 20 mL of 1N hydrochloric acid and washed with ethyl acetate (2 x 30 mL). The aqueous layer was basified with 10% aqueous sodium carbonate (50 mL) and extracted with ethyl acetate (3 x 50 mL). The ethyl acetate was washed with brine (1 x 30 mL), then dried (MgSO_4), and filtered. The filtered solution was concentrated under reduced to yield the corresponding aniline.

Step 4: 1-[2-(3-Dimethylamino-propoxy)-5-methyl-phenyl]-3-pyrazin-2-yl-urea. To a stirred, cooled (about 0°C) solution of 2-(3-dimethylamino-propoxy)-5-methyl-phenylamine (208 mg, 1.0 mmol) in toluene (3.0 mL) was added triethylamine (140 μ L, 1.0 mmol) and triphosgene (98 mg, 0.33 mmol). After stirring for 30 minutes, amino pyrazine (95 mg, 1.0 mmol) was added and the reaction was heated to 65 degrees C. After stirring for 4 hours, the reaction was cooled to room temperature and stirred for 8 hours. The precipitate that formed was filtered, rinsed with toluene (2 x 1 mL), and dried under reduced pressure (35% yield).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 8.45 (s, 1H), 8.39 (br s, 1H), 8.22 (s, 1H), 8.21 (d, 1H), 8.19 (d, 1H), 4.09 (t, 2H), 2.55 (t, 2H), 2.36 (s, 3H), 2.26 (s, 6H), 2.05 (m, 2H).

LRMS (esi, positive) m/e 330.10 (M+1)

Compound 254:




1-[2-(2-Dimethylamino-ethoxy)-5-methyl-phenyl]-3-pyrazin-2-yl-urea

Prepared according to the method of Compound 253 using N,N-dimethyl ethanolamine (36% yield).

$^1\text{H-NMR}$ (400 MHz, d_6 -DMSO) δ 8.79 (br s, 1H), 8.25 (s, 1H), 8.22 (s, 1H), 8.05 (s, 1H), 6.95 (d, 1H), 6.80 (d, 1H), 4.15 (t, 2H), 2.55 (s, 3H), 2.31 (t, 2H), 2.26 (s, 6H).

LRMS (esi, positive) m/e 316.21 (M+1)

Compound 255:




1-[5-Methyl-2-(pyridin-3-ylmethoxy)-phenyl]-3-pyrazin-2-yl-urea

Prepared according to the method of Compound 253 using 3-hydroxymethyl pyridine (10% yield).

¹H-NMR (400 MHz, CDCl₃) δ 8.82 (s, 1H), 8.65 (d, 1H), 8.35 (s, 1H), 8.30 (s, 1H), 8.21 (s, 2H), 8.05 (s, 1H), 7.81 (m, 2H), 7.35 (m, 1H), 6.9 (dd, 2H), 5.15 (s, 2H), 2.39 (s, 3H).

LRMS (esi, positive) m/e 336.21 (M+1)

Compound 256:




1-[5-Methyl-2-[3-(2-oxo-pyrrolidin-1-yl)-propoxy]-phenyl]-3-pyrazin-2-yl-urea

Prepared according to the method of Compound 253 using 3-(2-oxo-pyrrolidin-1-yl)-propanol (10% yield).

¹H-NMR (400 MHz, d₆-DMSO) δ 8.79 (s, 1H), 8.29 (s, 1H), 8.25 (s, 1H), 8.05 (s, 1H), 6.92 (d, 1H), 6.79 (d, 1H), 3.99 (t, 2H), 3.38 (m, 4H), 2.22 (s, 3H), 2.20 (t, 2H), 2.00 (t, 2H), 1.91 (t, 2H).

LRMS (esi, positive) m/e 392.2 (M+Na)

Compound 257:




1-[5-Methyl-2-(2-morpholin-4-yl-ethoxy)-phenyl]-3-pyrazin-2-yl-urea

Prepared according to the method of Compound 253 using 2-morpholin-4-yl-ethanol (39% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 8.79 (s, 1H), 8.29 (d, 1H), 8.25 (d, 1H), 8.05 (s, 1H), 6.95 (d, 1H), 6.79 (d, 1H), 4.19 (t, 2H), 3.59 (m, 4H), 2.80 (t, 2H), 2.49 (m, 4H), 2.22 (s, 3H).
LRMS (esi, positive) *m/e* 358.2 (M+1)

Compound 258:




1-[5-Methyl-2-(3-morpholin-4-yl-propoxy)-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea

Prepared according to the method of Compound 253 using 3-morpholin-4-yl-propanol (8% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 9.01 (s, 1H), 8.62 (br s, 1H), 8.22 (s, 1H), 8.19 (s, 1H), 8.05 (s, 1H), 6.91 (d, 1H), 6.79 (d, 1H), 4.05 (t, 2H), 3.59 (m, 4H), 2.48 (s, 3H), 2.45 (t, 2H), 2.35 (m, 4H), 2.21 (s, 3H), 2.00 (t, 2H).
LRMS (esi, positive) *m/e* 386.31 (M+1)

Compound 259:




1-[2-(3-Dimethylamino-propoxy)-5-methyl-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea

Prepared according to the method of Compound 253 3-dimethylamino-propanol (40% yield).

¹H-NMR (400 MHz, CDCl₃) δ 8.37 (s, 1H), 8.19 (s, 1H), 8.18 (s, 1H), 8.09 (br s, 1H), 6.80 (dd, 2H), 4.05 (t, 2H), 2.55 (t, 2H), 2.54 (s, 3H), 2.36 (s, 3H), 2.26 (s, 6H), 2.05 (m, 2H).
LRMS (esi, positive) *m/e* 344.20 (M+1)

Compound 260:




1-[5-Methyl-2-(2-morpholin-4-yl-ethoxy)-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea

Prepared according to the method of Compound 253 using 2-morpholin-4-yl-ethanol (10% yield).

¹H-NMR (400 MHz, CDCl₃) δ 10.79 (br s, 1H), 8.82 (s, 1H), 8.59 (s, 1H), 8.19 (s, 1H), 8.05 (s, 1H), 6.81 (dd, 2H), 4.20 (t, 2H), 3.75 (m, 4H), 2.91 (t, 2H), 2.61 (m, 4H), 2.55 (s, 3H), 2.35 (s, 3H).

LRMS (esi, positive) m/e 372.1 (M+1)

Compound 261:




1-(5-Methyl-pyrazin-2-yl)-3-[5-methyl-2-(pyridin-2-ylmethoxy)-phenyl]-urea

Prepared according to the method of Compound 253 using 2-hydroxymethyl pyridine (21% yield).

¹H-NMR (400 MHz, CDCl₃) δ 8.61 (d, 1H), 8.29 (s, 1H), 8.11 (s, 1H), 7.61 (t, 1H), 7.31 (d, 1H), 7.18 (t, 1H), 6.88 (d, 1H), 6.75 (d, 1H), 5.18 (s, 2H), 2.30 (s, 3H).

LRMS (esi, positive) m/e 372.2 (M+Na)

Compound 262:




1-(5-Methyl-pyrazin-2-yl)-3-[5-methyl-2-(pyridin-4-ylmethoxy)-phenyl]-urea

Prepared according to the method of Compound 253 using 4-hydroxymethyl pyridine (18% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 8.84 (s, 1H), 8.55 (d, 2H), 7.91 (s, 1H), 7.47 (d, 2H), 6.88 (d, 1H), 6.72 (d, 1H), 5.28 (s, 2H), 2.22 (s, 3H).

LRMS (esi, positive) m/e 350.21 (M+1)

Compound 263:




1-(5-Methyl-pyrazin-2-yl)-3-[5-methyl-2-(pyridin-3-ylmethoxy)-phenyl]-urea

Prepared according to the method of Compound 253 using 3-hydroxymethyl pyridine (10% yield).

¹H-NMR (400 MHz, CDCl₃) δ 8.82 (s, 1H), 8.68 (m, 2H), 8.25 (s, 1H), 8.20 (s, 1H), 7.84 (d, 1H), 7.38 (m, 1H), 6.99 (s, 1H), 6.88 (d, 1H), 6.80 (d, 1H), 5.10 (s, 2H), 2.38 (s, 3H), 2.35 (s, 3H).

LRMS (esi, positive) m/e 350.21 (M+1)

Compound 264:




1-[2-(2-Dimethylamino-ethoxy)-5-methyl-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea

Prepared according to the method of Compound 253 using N,N-dimethyl ethanolamine (11% yield).

¹H-NMR (400 MHz, d₆-DMSO) δ 8.69 (s, 1H), 8.18 (s, 1H), 8.03 (s, 1H), 6.95 (d, 1H), 6.79 (d, 1H), 4.11 (t, 2H), 2.72 (t, 2H), 2.43 (s, 3H), 2.25 (s, 3H), 2.22 (s, 6H).

LRMS (esi, positive) m/e 330.20 (M+1)

Compound 265:




1-(5-Methyl-pyrazin-2-yl)-3-[2-(pyridin-3-ylmethoxy)-5-trifluoromethyl-phenyl]-urea

Prepared according to the method of Compound 253 3-hydroxymethyl pyridine and 2-hydroxy-5-trifluoromethyl aniline (40% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 8.81 (s, 1H), 8.69 (s, 1H), 8.65 (s, 1H), 8.59 (br s, 1H), 8.01 (d, 1H), 7.45 (t, 1H), 7.3 (br s, 1H), 5.39 (s, 2H), 2.35 (s, 3H).

LRMS (esi, positive) *m/e* 404.10 (M+1)

Compound 266:



1-[5-Methyl-2-(6-methyl-pyridin-3-ylmethoxy)-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea

Step 1: (6-Methyl-pyridin-3-yl)-methanol. To a stirred, cooled (0 °C) solution of 6-methyl nicotinic acid (5.0 mmol) in tetrahydrofuran (10 mL) was added lithium aluminum hydride (20 mmol; 20 mL of a 1M solution in tetrahydrofuran.) dropwise. The reaction was stirred for 4 hours, treated sequentially with 1 mL of H₂O, 1 mL of 15% aqueous sodium hydroxide, and 3 mL of H₂O. The reaction was filtered and washed with tetrahydrofuran (3 x 50 mL). The filtrate was concentrated under reduced pressure to yield the alcohol as a clear viscous oil.


Steps 2-3: Mitsunobu reaction and aniline deprotection according to the method of Compound 253.

Step 4: To a stirred solution of 5-methyl-pyrazine-2-carboxylic acid (138 mg, 1.0 mmol) in toluene (3.0 mL) was added diphenylphosphoryl azide (216 µL, 1.0 mmol) and triethylamine (140 µL, 1.0 mmol). The reaction was placed under nitrogen and heated to 90 degrees C for 15 minutes. The temperature was reduced to 65 °C and 5-methyl-2-(6-methyl-pyridin-3-ylmethoxy)-phenylamine (228 mg, 1.0 mmol) was added. The reaction was stirred at that temperature for 4 hours and then stirred at room temperature for 8 hours. The precipitate that formed during the reaction was filtered, rinsed with toluene (2 x 1 mL), and dried under reduced pressure (36% yield).

¹H-NMR (400 MHz, CDCl₃) δ 11.45 (br s, 1H), 8.65 (s, 1H), 8.29 (s, 1H), 8.22 (s, 1H), 8.20 (s, 1H), 7.70 (d, 1H), 7.19 (d, 1H), 6.9 (m, 3H), 5.05 (s, 2H), 2.65 (s, 3H), 2.4 (s, 3H), 2.35 (s, 3H).

LRMS (esi, positive) *m/e* 364.16 (M+1)

Compound 267:



1-(5-Methyl-pyrazin-2-yl)-3-[5-methyl-2-(2-pyridin-2-yl-ethoxy)-phenyl]-urea


Steps 1-2: Mitsunobu reaction using 2-(2-pyridyl)-ethanol and aniline deprotection according to the method of Compound 253.

Step 3: Urea formation according to the method of Compound 266 using 5-methyl-2-(2-pyridin-2-yl-ethoxy)-phenylamine (37% yield).

¹H-NMR (400 MHz, CDCl₃) δ 10.70 (br s, 1H), 8.65 (d, 1H), 8.45 (br s, 1H), 8.21 (s, 1H), 7.85 (s, 1H), 7.59 (t, 1H), 7.29 (t, 1H), 6.80 (dd, 2H), 4.49 (t, 2H), 3.39 (t, 2H), 2.49 (s, 3H), 2.39 (s, 3H).

LRMS (esi, positive) m/e 364.14 (M+1)

Compound 268:



1-(5-Methyl-pyrazin-2-yl)-3-[5-methyl-2-(3-pyridin-2-yl-propoxy)-phenyl]-urea


Steps 1-2: Mitsunobu reaction using 3-(2-pyridyl)-propanol and aniline deprotection according to the method of Compound 253.

Step 3: Urea formation according to the method of Compound 266 using 5-methyl-2-(3-pyridin-2-yl-propoxy)-phenylamine (5% yield).

¹H-NMR (400 MHz, CDCl₃) δ 10.89 (br s, 1H), 8.59 (d, 1H), 8.49 (s, 1H), 8.45 (br s, 1H), 8.22 (s, 1H), 8.15 (s, 1H), 7.59 (t, 1H), 7.15 (d, 2H), 6.88 (dd, 2H), 4.05 (t, 2H), 3.10 (t, 2H), 2.45 (s, 3H), 2.40 (t, 2H), 2.35 (s, 3H).

LRMS (esi, positive) m/e 378.10 (M+1)

Compound 269:



1-(5-Methyl-pyrazin-2-yl)-3-[5-methyl-2-(3-pyridin-4-yl-propoxy)-phenyl]-urea


Steps 1-2: Mitsunobu reaction using 3-(4-pyridyl)-propanol and aniline deprotection according to the method of Compound 253.

Step 3: Urea formation according to the method of Compound 266 using 5-methyl-2-(4-pyridin-2-yl-propoxy)-phenylamine (28% yield).

¹H-NMR (400 MHz, CDCl₃) δ 11.15 (br s, 1H), 8.51 (d, 2H), 8.25 (s, 1H), 8.20 (s, 1H), 7.79 (s, 1H), 7.75 (s, 1H), 7.15 (d, 2H), 6.82 (d, 1H), 6.75 (d, 1H), 4.05 (t, 2H), 2.91 (t, 2H), 2.40 (s, 3H), 2.36 (s, 3H), 2.25 (m, 2H).

LRMS (esi, positive) m/e 378.16 (M+1)

Compound 270:



1-{2-[2-(Benzyl-methyl-amino)-ethoxy]-5-methyl-phenyl}-3-(5-methyl-pyrazin-2-yl)-urea


Steps 1-2: Mitsunobu reaction using N-methyl-N-benzyl ethanolamine and aniline deprotection according to the method of Compound 253.

Step 3: Urea formation according to the method of Compound 266 using 2-[2-(benzyl-methyl-amino)-ethoxy]-5-methyl-phenylamine (17% yield).

¹H-NMR (400 MHz, CDCl₃) δ 10.70 (br s, 1H), 8.49 (s, 1H), 8.20 (s, 1H), 7.95 (s, 1H), 7.85 (s, 1H), 7.32 (m, 5H), 6.85 (s, 2H), 4.15 (t, 2H), 3.62 (s, 2H), 2.92 (t, 2H), 2.49 (s, 3H), 2.33 (s, 3H), 2.29 (s, 3H).

LRMS (esi, positive) m/e 406.01 (M+1)

Compound 271:



1-[5-Methyl-2-(1-methyl-piperidin-3-ylmethoxy)-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea


Steps 1-2: Mitsunobu reaction using 3-hydroxymethyl-1-methyl piperidine and aniline deprotection according to the method of Compound 253.

Step 3: Urea formation according to the method of Compound 266 using 5-methyl-2-(1-methyl-piperidin-3-ylmethoxy)-phenylamine (16% yield).

¹H-NMR (400 MHz, CDCl₃) δ 11.25 (br s, 1H), 8.45 (br s, 1H), 8.35 (s, 1H), 8.22 (s, 2H), 6.80 (d, 1H), 6.74 (d, 1H), 3.80 (m, 2H), 3.15 (br d, 1H), 2.80 (br d, 1H), 2.51 (s, 3H), 2.35 (s, 3H), 2.30 (m, 1H), 2.22 (s, 3H), 1.50-2.00 (m, 6H), 1.00-1.25 (m, 2H).

LRMS (esi, positive) m/e 370.01 (M+1)

Compound 272:



1-{2-[2-(4-Dimethylamino-phenyl)-ethoxy]-5-methyl-phenyl}-3-(5-methyl-pyrazin-2-yl)-urea


Steps 1-2: Mitsunobu reaction using 2-(4-dimethylamino-phenyl)-ethanol and aniline deprotection according to the method of Compound 253.

Step 3: Urea formation according to the method of Compound 266 using 2-[2-(4-dimethylamino-phenyl)-ethoxy]-5-methyl-phenylamine (10% yield).

¹H-NMR (400 MHz, CDCl₃) δ 11.15 (br s, 1H), 8.35 (s, 1H), 8.25 (s, 1H), 7.80 (m, 2H), 7.25 (m, 2H), 6.82 (s, 2H), 6.75 (d, 2H), 4.25 (t, 2H), 3.20 (t, 2H), 2.99 (s, 6H), 2.55 (s, 3H), 2.41 (s, 3H).

LRMS (esi, positive) m/e 405.90 (M+1)

Compound 273:



1-(5-Methyl-pyrazin-2-yl)-3-[5-methyl-2-(3-pyridin-3-yl-propoxy)-phenyl]-urea


Steps 1-2: Mitsunobu reaction using 3-(3-pyridyl)-propanol and aniline deprotection according to the method of Compound 253.

Step 3: Urea formation according to the method of Compound 266 using 5-methyl-2-(3-pyridin-3-yl-propoxy)-phenylamine (16% yield).

¹H-NMR (400 MHz, CDCl₃) δ 10.95 (br s, 1H), 8.51 (m, 2H), 8.35 (s, 1H), 8.25 (s, 1H), 8.15 (s, 1H), 7.80 (s, 1H), 7.50 (d, 1H), 7.21 (t, 1H), 6.79 (d, 1H), 6.75 (d, 1H), 4.09 (t, 2H), 2.90 (t, 2H), 2.45 (s, 3H), 2.35 (s, 3H), 2.25 (m, 2H).

LRMS (esi, positive) m/e 377.91 (M+1)

Compound 274:



1-[2-(2-Dimethylamino-1-dimethylaminomethyl-ethoxy)-5-methyl-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea


Steps 1-2: Mitsunobu reaction using 1,3-bis-dimethylamino-propan-2-ol and aniline deprotection according to the method of Compound 253.

Step 3: Urea formation according to the method of Compound 266 using 2-(2-amino-4-methyl-phenoxy)-N,N,N',N'-tetramethyl-propane-1,3-diamine (4% yield).

¹H-NMR (400 MHz, CDCl₃) δ 9.69 (br s, 1H), 8.95 (br s, 1H), 8.19 (s, 1H), 8.05 (s, 1H), 6.80 (dd, 2H), 4.19 (m, 1H), 4.09 (m, 1H), 3.05 (m, 1H), 2.65 (m, 2H), 2.50 (s, 3H), 2.45 (s, 6H), 2.38 (s, 6H), 2.35 (s, 3H).

LRMS (esi, positive) m/e 386.92 (M+1)

Compound 275:



1-[5-Methyl-2-(2-S-1-methyl-pyrrolidin-2-ylmethoxy)-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea


Steps 1-2: Mitsunobu reaction using 1-methyl-pyrrolidin-2-S-ylmethanol and aniline deprotection according to the method of Compound 253.

Step 3: Urea formation according to the method of Compound 266 using 5-methyl-2-(1-methyl-pyrrolidin-2-S-ylmethoxy)-phenylamine (12% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 10.10 (s, 1H), 9.85 (br s, 1H), 9.65 (br s, 1H), 8.78 (s, 1H), 8.25 (s, 1H), 6.90 (s, 1H), 7.02 (d, 1H), 6.85 (d, 1H), 4.33 (br s, 2H), 3.88 (m, 1H), 3.59 (m, 1H), 3.19 (m, 1H), 2.99 (d, 2H) 2.70-2.85 (m, 2H), 2.45 (s, 3H), 2.29 (s, 3H), 1.80-2.10 (m, 3H).

LRMS (esi, positive) *m/e* 355.91 (M+1)

Compound 276:



1-[2-(2-S-1-Benzyl-pyrrolidin-2-ylmethoxy)-5-methyl-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea


Steps 1-2: Mitsunobu reaction using 1-benzyl-pyrrolidin-2-S-ylmethanol and aniline deprotection according to the method of Compound 253.

Step 3: Urea formation according to the method of Compound 266 using 5-methyl-2-(1-benzyl-pyrrolidin-2-S-ylmethoxy)-phenylamine (3% yield).

¹H-NMR (400 MHz, *d*₆-DMSO) δ 9.95 (s, 1H), 9.90 (br s, 1H), 9.59 (br s, 1H), 8.69 (s, 1H), 8.10 (s, 1H), 7.89 (s, 1H), 7.25-7.50 (m, 6H), 6.96 (d, 1H), 6.90 (d, 1H), 4.75 (d, 2H), 4.33 (m, 4H), 4.10 (m, 2H), 2.40 (s, 3H), 2.25 (s, 3H), 1.80-2.10 (m, 3H), 1.10-1.30 (m, 2H).

LRMS (esi, positive) *m/e* 432.31 (M+1)

Compound 277:




1-(5-Methyl-pyrazin-2-yl)-3-[5-methyl-2-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-urea

Prepared according to the procedure for compound 266, using 2-pyrrolidin-1-yl-ethanol. (28% yield).

¹H NMR (400 MHz, *d*₆-DMSO) δ 10.01 (s, 1H), 9.85 (br s, 1H), 9.72 (br s, 1H), 8.75 (s, 1H), 8.25 (s, 1H), 8.01 (s, 1H), 7.01 (d, 1H), 6.80 (d, 1H), 4.40 (t, 2H), 3.62 (m, 4H), 3.21 (m, 2H), 2.40 (s, 3H), 2.25 (s, 3H), 2.00 (m, 2H), 1.88 (m, 2H).

LRMS (ESI, Positive) *m/e* 356.2 (M+1).

Compound 278:




1-{5-Methyl-2-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-phenyl}-3-(5-methyl-pyrazin-2-yl)-urea

Prepared according to the procedure for compound 266, using 2-(1-methyl-pyrrolidin-2-yl)-ethanol.

(32% yield). ¹H NMR (400 MHz, *d*₆-DMSO) δ 10.01 (s, 1H), 9.85 (br s, 1H), 9.72 (br s, 1H), 8.75 (s, 1H), 8.25 (s, 1H), 8.01 (s, 1H), 7.01 (d, 1H), 6.80 (d, 1H), 4.20 (m, 3H), 3.00-4.00 (m, 11H), 2.80 (d, 2H), 2.40 (s, 3H), 2.25 (s, 3H).

LRMS (ESI, Positive) *m/e* 370.2 (M+1).

Compound 279:




1-[2-(3H-Imidazol-4-ylmethoxy)-5-methyl-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea

Prepared according to the procedure for compound 266, using (3H-imidazol-4-yl)-methanol. (24%

yield) ¹H NMR (400 MHz, *d*₆-DMSO) δ 8.51 (br s, 1H), 8.01 (s, 1H), 7.69 (s, 1H), 7.20-7.50 (m, 2H), 7.10 (d, 1H), 6.75 (d, 1H), 4.99 (s, 2H), 2.40 (s, 3H), 2.25 (s, 3H). LRMS (ESI, Positive) *m/e* 339.1 (M+1).


Compound 280:



1-(5-Methyl-pyrazin-2-yl)-3-[5-methyl-2-(2-pyridin-3-yl-ethoxy)-phenyl]-urea

Prepared according to the procedure for compound 266, using 2-pyridin-3-yl-ethanol. (16% yield). ¹H NMR (400 MHz, CDCl₃) δ 10.98 (br s, 1H), 8.65 (s, 1H), 8.49 (d, 1H), 8.35 (s, 2H), 8.20 (s, 1H), 7.75 (s, 1H), 7.65 (d, 1H), 7.19 (m, 1H), 6.82 (dd, 2H), 4.31 (t, 2H), 3.21 (t, 2H), 2.48 (s, 3H), 2.35 (s, 3H). LRMS (ESI, Positive) m/e 364.2 (M+1).


Compound 281:



1-[5-Methyl-2-(3-piperidin-1-yl-propoxy)-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea

Prepared according to the procedure for compound 266, using 3-piperidin-1-yl-propan-1-ol. (33% yield). ¹H NMR (400 MHz, CDCl₃) δ 11.21 (br s, 1H), 8.35 (s, 1H), 8.25 (s, 2H), 8.15 (s, 1H), 6.80 (dd, 2H), 4.15 (t, 2H), 2.53 (t, 2H), 2.52 (s, 3H), 2.45 (m, 4H), 2.39 (s, 3H), 2.10 (m, 2H), 1.61 (m, 4H), 1.45 (m, 2H). LRMS (ESI, Positive) m/e 384.2 (M+1).


Compound 282:



1-[5-Methyl-2-(1-methyl-piperidin-4-yloxy)-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea

Prepared according to the procedure for compound 266, using 1-methyl-piperidin-4-ol. (4% yield). ¹H NMR (400 MHz, CDCl₃) δ 11.22 (br s, 1H), 8.43 (s, 1H), 8.25 (s, 1H), 8.20 (s, 1H), 8.10 (s, 1H), 6.81 (dd, 2H), 4.25 (m, 1H), 2.8 (m, 2H), 2.59 (s, 3H), 2.39 (s, 3H), 2.36 (s, 3H), 2.19 (m, 4H), 1.90 (m, 2H). LRMS (ESI, Positive) m/e 355.9 (M+1).

Compound 283:




1-[2-(1-Benzyl-piperidin-4-yloxy)-5-methyl-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea

Prepared according to the procedure for compound 266, using 1-benzyl-piperidin-4-ol. (1% yield).

LRMS (ESI, Positive) m/e 432.0 (M+1).

Compound 284:




1-[5-Methyl-2-(3-(S)-1-methyl-piperidin-3-ylmethoxy)-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea

Step 1: (1-Methyl-piperidin-3-(S)-yl)-methanol To a stirred, cooled solution of (S)-(+)-*N*-boc nipecotic acid (5.0 mmol) in tetrahydrofuran (10 mL) lithium aluminum hydride (20 mL, 20 mmol, 1M in tetrahydrofuran.) was added dropwise. The reaction was refluxed for 12 hours and then cooled to 0 °C. The reaction was quenched with 1 mL of H₂O, 1 mL of 15% aqueous sodium hydroxide, 3 mL of H₂O. The reaction was filtered and the filter cake was washed with tetrahydrofuran (3 x 50 mL). The filtrate was concentrated under reduced pressure to yield a clear viscous oil.

Steps 2-3: 5-Methyl-2-(1-methyl-piperidin-3-(S)-ylmethoxy)-phenylamine Mitsunobu reaction and aniline deprotection according to procedure for compound 253.

Step 4: Prepared according to the procedure for compound 266 (33% yield). ¹H NMR (400 MHz, CDCl₃) δ 11.25 (br s, 1H), 8.45 (br s, 1H), 8.35 (s, 1H), 8.22 (s, 2H), 6.80 (d, 1H), 6.74 (d, 1H), 3.80 (m, 2H), 3.15 (br d, 1H), 2.80 (br d, 1H), 2.51 (s, 3H), 2.35 (s, 3H), 2.30 (m, 1H), 2.22 (s, 3H), 1.50-2.00 (m, 6H), 1.00-1.25 (m, 2H). LRMS (ESI, Positive) m/e 370.0 (M+1).

Compound 285:




1-[5-Methyl-2-(3-(R)-1-methyl-piperidin-3-ylmethoxy)-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea

Steps 1-3: **5-Methyl-2-(1-methyl-piperidin-3-(R)-ylmethoxy)-phenylamine** according to the method for compound 284, using (R)-(+)-*N*-boc nipecotic acid.

Step 4: 5-Methylpyrazine-2-carboxylic azide (1.2 eq.) dissolved in anhydrous toluene (0.1 M concentration) was heated to 90 °C. After 20 minutes N₂ evolution had subsided, and the caramel colored reaction mixture was cooled to 60 °C before adding the aniline prepared above as a solution in toluene (1 eq.). After stirring for 4 hours at 60 °C, the reaction mixture was cooled to room temperature overnight. A precipitate formed which was isolated by filtration (49% yield). ¹H NMR (400 MHz, CDCl₃) δ 11.25 (br s, 1H), 8.45 (br s, 1H), 8.35 (s, 1H), 8.22 (s, 2H), 6.80 (d, 1H), 6.74 (d, 1H), 3.80 (m, 2H), 3.15 (br d, 1H), 2.80 (br d, 1H), 2.51 (s, 3H), 2.35 (s, 3H), 2.30 (m, 1H), 2.22 (s, 3H), 1.50-2.00 (m, 6H), 1.00-1.25 (m, 2H). LRMS (ESI, Positive) m/e 370.0 (M+1).

Compound 286:




1-(5-Methyl-pyrazin-2-yl)-3-[5-methyl-2-(1-pyridin-3-yl-ethoxy)-phenyl]-urea

Step 1: **1-Pyridin-3-yl-ethanol** To a stirred, cooled (-78°C) solution of pyridine-3-carbaldehyde (15 mmol) in tetrahydrofuran (40 mL), methyl magnesium bromide (5 mL, 15 mmol, 3M in diethyl ether) was added. After stirring for 2 hours, the reaction was quenched with saturated aqueous ammonium chloride (5 mL). The pH was adjusted to ~5.0 with aqueous sodium carbonate and the product was extracted with ethyl acetate (3 x 100 mL). The ethyl acetate was washed with brine (1 x 100 mL), dried (MgSO₄), and filtered. The filtered material was concentrated under reduced pressure to yield a yellow oil.

Steps 2-3: **5-Methyl-2-(1-pyridin-3-yl-ethoxy)-phenylamine** Mitsunobu reaction and aniline deprotection according to the method for compound 253.

Step 4: Urea formation was conducted according to the procedure for compound 285. (17% yield). ¹H NMR (400 MHz, CDCl₃) δ 11.49 (br s, 1H), 8.81 (s, 1H), 8.73 (s, 1H), 8.52 (d, 1H), 8.35 (s, 1H), 8.25 (s, 1H), 8.18 (s, 1H), 7.80 (s, 1H), 7.72 (d, 1H), 7.20 (t, 1H), 6.70 (d, 1H), 6.65 (d, 1H), 5.49 (q, 1H), 2.50 (s, 3H), 2.30 (s, 3H), 1.75 (d, 3H). LRMS (ESI, Positive) m/e 363.8 (M+1).

Compound 287:




1-[5-Methyl-2-(1-methyl-piperidin-2-ylmethoxy)-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea

Steps 1-2: **5-Methyl-2-(1-methyl-piperidin-2-ylmethoxy)-phenylamine** Mitsunobu reaction and aniline deprotection according to the method for compound 253.

Step 3: Urea formation was conducted according to the procedure for compound 285. (11% yield). ¹H NMR (400 MHz, CDCl₃) δ 10.90 (br s, 1H), 8.45 (s, 1H), 8.21 (s, 1H), 8.18 (s, 1H), 7.80 (s, 1H), 6.80 (s, 2H), 4.51 (m, 1H), 2.58-3.00 (m, 4H), 2.52 (s, 3H), 2.46 (s, 3H), 2.35 (s, 3H), 1.50-2.25 (m, 7H). LRMS (ESI, Positive) m/e 369.9 (M+1).

Compound 288:




1-(5-Methyl-pyrazin-2-yl)-3-[5-methyl-2-(tetrahydro-furan-2-ylmethoxy)-phenyl]-urea

Steps 1-2: **5-Methyl-2-(tetrahydro-furan-2-ylmethoxy)-phenylamine** Mitsunobu reaction and aniline deprotection according to the method for compound 253.

Step 3: Urea formation was conducted according to the procedure for compound 285. (12% yield). ¹H NMR (400 MHz, CDCl₃) δ 11.25 (br s, 1H), 8.49 (s, 1H), 8.40 (s, 1H), 8.22 (s, 1H), 8.18 (s, 1H), 6.8 (s, 2H), 4.42 (m, 1H), 3.80-4.10 (m, 4H), 2.52 (s, 3H), 2.35 (s, 3H), 1.20-2.20 (m, 4H). LRMS (ESI, Positive) m/e 342.9 (M+1).

Compound 289:



1-[5-Methyl-2-(1-methyl-piperidin-4-ylmethoxy)-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea

Step 1: (1-Methyl-piperidin-4-yl)-methanol To a stirred, cooled solution of 1-methyl-piperidine-4-carboxylic acid (5.0 mmol) in tetrahydrofuran (10 mL) lithium aluminum hydride (20 mL, 20 mmol, 1M tetrahydrofuran.) was added dropwise. The reaction was stirred for 4 hours, quenched with 1 mL of H₂O, 1 mL of 15% aqueous sodium hydroxide, and 3 mL of H₂O. The reaction was filtered and washed with tetrahydrofuran (3 x 50 mL). The filtrate was concentrated under reduced pressure to yield a clear viscous oil.


Steps 2-3: 5-Methyl-2-(1-methyl-piperidin-4-ylmethoxy)-phenylamine Mitsunobu and aniline deprotection according to the procedure for compound 253.

Step 4: Urea formation according to the procedure for compound 285 (54% yield).

¹H NMR (400 MHz, CDCl₃) δ 11.18 (br s, 1H), 8.62 (br s, 1H), 8.38 (br s, 1H), 8.25 (s, 1H), 8.18 (s, 1H), 6.82 (d, 1H), 6.78 (d, 1H), 3.85 (d, 2H), 2.90 (br d, 2H), 2.51 (s, 3H), 2.35 (m, 6H), 1.50-2.10 (m, 7H).

LRMS (ESI, Positive) m/e 369.2 (M+1).

Compound 290:




1-[5-Methyl-2-(1-methyl-piperidin-3-yloxy)-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea

Steps 1-2: 5-Methyl-2-(1-methyl-piperidin-3-yloxy)-phenylamine Mitsunobu reaction and aniline deprotection according to the method for compound 253.

Step 3: Urea formation according to the procedure for compound 285 (3% yield). ¹H NMR (400 MHz, CDCl₃) δ 10.75 (br s, 1H), 8.59 (br s, 1H), 8.18 (s, 1H), 8.05 (s, 1H), 7.62 (s, 1H), 6.90 (d, 1H), 6.80 (d, 1H), 4.40 (m, 1H), 2.58 (s, 3H), 2.39 (s, 6H), 1.60-2.80 (m, 8H).

LRMS (ESI, Positive) m/e 356.1 (M+1).

Compound 291:



1-(5-Methyl-pyrazin-2-yl)-3-[5-methyl-2-(1,2,3,4-tetrahydro-quinolin-3-ylmethoxy)-phenyl]-urea

Step 1: Quinoline-3-carboxylic acid methyl ester. To a stirred solution of quinoline-3-carboxylic acid (346 mg, 2 mmol) dissolved in 4:1 THF:MeOH (6 mL) at 0 °C was added TMS-diazomethane (2M in hexane) portionwise until a diazomethane yellow color persisted. The reaction was concentrated to give the methyl ester as a tan solid (244 mg, 65%). ¹H-NMR (400 MHz, CDCl₃) δ 9.44 (s, 1H), 8.85 (s, 1H), 8.17 (d, 1H), 7.96 (d, 1H), 7.84 (dd, 1H), 7.62 (dd, 1H), 4.02 (s, 3H)

Step 2: 1,2,3,4-Tetrahydro-quinoline-3-carboxylic acid methyl ester and 1-Ethyl-1,2,3,4-tetrahydro-quinoline-3-carboxylic acid methyl ester. To a stirred solution of the quinoline-3-carboxylic acid methyl ester (244 mg, 1.3 mmol) in glacial acetic acid (13 mL) at room temperature was added NaBH₄ (345 mg, 9.1 mmol) portionwise (vigorous reaction). After complete addition the reaction was dark yellow. After stirring for 3 hours the color had become pale yellow. The reaction mixture was poured into 50 mL of H₂O and 50 mL of CH₂Cl₂ and stirred rapidly for 15 min. The layers were separated and the organics concentrated to a yellow oil. TLC in 15/85 EtOAc/hexane showed complete consumption of starting material, and two new lower rf spots. The compound was chromatographed using a Biotage 12M column (loaded with CH₂Cl₂) and eluted with 15/85 EtOAc/hexane. The higher rf spot corresponds to the N-ethylated product (123 mg, 43%). The lower rf spot corresponds to the desired tetrahydroquinoline-3-carboxylic acid methyl ester (93 mg, 37%). N-ethyl derivative: ¹H-NMR (400 MHz, CDCl₃) δ 7.08 (dd, 1H), 6.99 (d, 1H), 6.60 (m, 2H), 3.73 (s, 3H), 3.42 (m, 3H), 3.27 (m, 1H), 2.98 (m, 3H), 1.14 (t, 3H) N-H derivative: ¹H-NMR (400 MHz, CDCl₃) δ 6.98 (m, 2H), 6.62 (dd, 1H), 6.47 (d, 1H), 3.71 (s, 3H), 3.52 (m, 1H), 3.34 (m, 1H), 3.00 (m, 2H), 2.90 (m, 1H)

Step 3: (1,2,3,4-Tetrahydro-quinolin-3-yl)-methanol. To a stirred solution of the 1,2,3,4-Tetrahydro-quinoline-3-carboxylic acid methyl ester (93 mg, 0.49 mmol) in 1.5 mL of Et₂O at 0 °C under nitrogen was added LiAlH₄ (1M in Et₂O) dropwise with vigorous gas evolution and a white precipitate formation. After 30 min., the reaction was carefully quenched with 15% NaOH (3 mL) and 3 mL of Et₂O was added and the mixture stirred rapidly at RT for 15 min. The layers were separated and the aqueous layer extracted (1 x 10 mL) with Et₂O. The organics were combined, dried (MgSO₄), filtered and concentrated to the alcohol (64 mg, 80%). ¹H-NMR (400 MHz, CDCl₃) δ 6.97 (m, 2H), 6.62 (dd, 1H), 6.47 (d, 1H), 3.66 (m, 1H), 3.58 (m, 1H), 3.40 (m, 1H), 3.08 (m, 1H), 2.82 (m, 1H), 2.53 (m, 1H), 2.18 (m, 1H).

Step 4: [5-Methyl-2-(1,2,3,4-tetrahydro-quinolin-3-ylmethoxy)-phenyl]-carbamic acid tert-butyl ester. To a stirred solution of 2-N-Boc-amino-4-methylphenol (88 mg, 0.39 mmol), (1,2,3,4-Tetrahydro-quinolin-3-yl)-methanol (64 mg, 0.39 mmol) and triphenylphosphine (103 mg, 0.39 mmol) in 850 μ L of THF at 0 °C under nitrogen was added a solution of DIAD (77 μ L, 0.39 mmol) in 850 μ L THF. The reaction was allowed to warm to RT overnight, was concentrated and loaded directly onto a Biotage 12M column with CH_2Cl_2 and eluted with 96/4 hexane/EtOAc. The product was isolated as a yellow oil (129 mg, 89%). $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 7.92 (br s, 1H), 7.06 (br s, 1H), 6.99 (m, 2H), 6.81 (m, 1H), 6.73 (s, 2H), 6.65 (dd, 1H), 6.51 (d, 1H), 3.96 (m, 2H), 3.37 (ddd, 2H), 2.79 (ddd, 2H), 2.56 (m, 1H), 2.28 (s, 3H), 1.54 (s, 9H)

Step 5: 3-(2-tert-Butoxycarbonylamino-4-methyl-phenoxy-methyl)-3,4-dihydro-2H-quinoline-1-carboxylic acid benzyl ester. To a stirred solution of [5-Methyl-2-(1,2,3,4-tetrahydro-quinolin-3-ylmethoxy)-phenyl]-carbamic acid tert-butyl ester. To a stirred solution of 2-N-Boc-amino-4-methylphenol (129 mg, 0.35 mmol) in CH_2Cl_2 (1.5 mL) at 0 °C under nitrogen was added DIEA (61 μ L, 0.35 mmol) followed by benzyl chloroformate (50 μ L, 0.35 mmol) and DMAP (4 mg, 0.035 mmol). After 24 hours, the reaction was diluted to 30 mL with CH_2Cl_2 and washed (2 x 30 mL) with 2N HCl and (2 x 30 mL) with saturated NaHCO_3 . The organics were dried (MgSO_4), filtered and concentrated to a brown oil, which appeared to be a mixture of product and starting material.


Step 6: 3-(2-Amino-4-methyl-phenoxy-methyl)-3,4-dihydro-2H-quinoline-1-carboxylic acid benzyl ester. A solution of the crude 3-(2-tert-Butoxycarbonylamino-4-methyl-phenoxy-methyl)-3,4-dihydro-2H-quinoline-1-carboxylic acid benzyl ester (crude, 0.35 mmol) in 4N HCl in dioxane (2 mL) at room temperature was stirred under a drying tube overnight. The suspension was concentrated by rotovap, diluted to 30 mL with CH_2Cl_2 and shaken with 10% Na_2CO_3 (30 mL). The organics were isolated, dried (MgSO_4), filtered and concentrated to a brown oil corresponding to the crude aniline, which was used without purification in the urea forming reaction.

Step 7: 3-{4-Methyl-2-[3-(5-methyl-pyrazin-2-yl)-ureido]-phenoxy-methyl}-3,4-dihydro-2H-quinoline-1-carboxylic acid benzyl ester. A 0.5 M solution of the 5-methyl-pyrazine-2-carboxylic acid azide (204 μ L) was diluted with toluene (408 μ L) in a septum capped reaction vial under nitrogen and with stirring, immersed in a 90 °C oil bath. After about 20 minutes nitrogen gas evolution had stopped so the reaction was allowed to cool to RT and was treated with a solution of the crude 3-(2-Amino-4-methyl-phenoxy-methyl)-3,4-dihydro-2H-quinoline-1-carboxylic acid benzyl ester (ca. 0.101 mmol) in toluene (620 μ L). The mixture was stirred at 65 °C for 2 hours. The reaction was cooled to rt overnight and a precipitate formed. The precipitate was filtered off with toluene and appeared to be a mixture of Cbz-protected and deprotected product.

Step 8: 1-(5-Methyl-pyrazin-2-yl)-3-[5-methyl-2-(1,2,3,4-tetrahydro-quinolin-3-ylmethoxy)-phenyl]-urea. A stirred suspension of the crude 3-{4-Methyl-2-[3-(5-methyl-pyrazin-2-yl)-ureido]-phenoxy-methyl}-3,4-dihydro-2H-quinoline-1-carboxylic acid benzyl ester (6.6 mg, 12 μ mol) was

heated in 5 mL EtOAc on a heat gun until in solution. The clear solution was cooled to rt and treated with triethylamine (3.4 μ L, 24 μ mol) followed by Pearlman's catalyst (20% palladium hydroxide on carbon, 9 mg). The mixture was put through a vacuum/purge cycle three times with hydrogen gas and then held under 1 atmosphere hydrogen pressure for 1 hour. The reaction was filtered through GF/F filter paper with EtOAc and concentrated to a white solid which corresponds to the desired product (4.7 mg, 100%). $^1\text{H-NMR}$ (400 MHz, $\text{CDCl}_3/\text{CD}_3\text{OD}$) δ 8.23 (s, 1H), 8.18 (s, 1H), 7.96 (s, 1H), 7.04 (t, 1H), 6.97 (d, 1H), 6.81 (m, 2H), 6.67 (t, 1H), 6.58 (d, 1H), 4.04 (m, 2H), 3.57 (d, 1H), 3.26 (t, 1H), 2.92 (ddd, 2H), 2.64 (m, 1H), 2.34 (s, 3H), 2.31 (s, 3H) LRMS (APCI, Positive) m/e 404.2 (M+1).

Compound 292:



1-[2-(1-Ethyl-1,2,3,4-tetrahydro-quinolin-3-ylmethoxy)-5-methyl-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea


Step 1: (1-Ethyl-1,2,3,4-tetrahydro-quinolin-3-yl)-methanol. To a stirred solution of 1-Ethyl-1,2,3,4-tetrahydro-quinoline-3-carboxylic acid methyl ester (123 mg, 0.56 mmol) in 1.5 mL of Et_2O at 0 $^\circ\text{C}$ under nitrogen. LAH (1M in Et_2O) was added dropwise with vigorous gas evolution and a white precipitate formation. After 30 min., TLC in 3/7 EtOAc/hexane showed complete loss of s.m. and appearance of a clean lower rf spot. The reaction was carefully quenched with 15% NaOH (3 mL) and 3 mL of Et_2O was added and the mixture stirred rapidly at RT for 15 min. The layers were separated and the aqueous layer extracted 1 x 10 mL with Et_2O . The organics were combined, dried (MgSO_4), filtered and concentrated to a clear oil corresponding to the desired alcohol (105 mg, 95%).

Step 2: [2-(1-Ethyl-1,2,3,4-tetrahydro-quinolin-3-ylmethoxy)-5-methyl-phenyl]-carbamic acid tert-butyl ester. To a stirred solution of (1-Ethyl-1,2,3,4-tetrahydro-quinolin-3-yl)-methanol (105 mg, 0.55 mmol, prepared in step 1, compound 126xx), 2-N-Boc-amino-4-methylphenol (123 mg, 0.55 mmol), and triphenylphosphine (144 mg, 0.55 mmol) in 850 μL of THF at 0 $^\circ\text{C}$ under nitrogen was added a solution of DIAD (108 μL , 0.55 mmol) in 850 μL THF. The reaction was allowed to warm to RT overnight, was concentrated and loaded directly onto a Biotage 12M column with CH_2Cl_2 and eluted with 96/4 hexane/EtOAc to give the desired alkylated phenol as a white foam (40 mg, 18%). $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 7.92 (br s, 1H), 7.08 (dd, 1H), 7.02 (m, 1H), 6.99 (d, 1H), 6.75 (s, 2H), 6.62 (d, 1H), 6.59 (dd, 1H), 3.98 (m, 2H), 3.39 (m, 4H), 3.20 (m, 1H), 2.79 (ddd, 2H), 2.58 (m, 1H), 2.28 (s, 3H), 1.56 (s, 9H), 1.16 (t, 3H).

Step 3: 2-(1-Ethyl-1,2,3,4-tetrahydro-quinolin-3-ylmethoxy)-5-methyl-phenylamine. A solution of [2-(1-Ethyl-1,2,3,4-tetrahydro-quinolin-3-ylmethoxy)-5-methyl-phenyl]-carbamic acid tert-butyl ester (40 mg, 0.1 mmol) was stirred in 4N HCl in dioxane (2 mL) at room temperature under a drying tube overnight. The suspension was concentrated by rotovap, diluted to 30 mL with CH₂Cl₂ and shaken with 10% Na₂CO₃ (30 mL). The organics were isolated, dried (MgSO₄), filtered and concentrated to a brown oil, which was used without purification in the following reaction.

Step 4: 1-[2-(1-Ethyl-1,2,3,4-tetrahydro-quinolin-3-ylmethoxy)-5-methyl-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea. A 0.5 M solution of the acyl azide (182 µL) was diluted with 364 µL toluene in a septum capped reaction vial under nitrogen and with stirring, immersed in a 90 °C oil bath. After about 20 minutes N₂ gas evolution had stopped so the reaction was allowed to cool to rt and was treated with a solution of 2-(1-ethyl-1,2,3,4-tetrahydro-quinolin-3-ylmethoxy)-5-methyl-phenylamine (27 mg, 0.91 mmol) in 550 µL of toluene. The mixture was stirred at 65 °C for 2 hours. The reaction was cooled to rt overnight and a precipitate formed. The precipitate was filtered off with toluene and the desired urea isolated as a tan solid (7 mg, 18 %). ¹H-NMR (400 MHz, CDCl₃) δ 11.34 (br s, 1H), 8.25 (s, 1H), 8.21 (s, 1H), 8.11 (br s, 1H), 7.83 (s, 1H), 7.25 (m, 1H), 7.18 (d, 1H), 7.12 (t, 1H), 6.97 (d, 1H), 6.80 (m, 2H), 6.70 (d, 1H), 6.61 (t, 1H), 4.05 (m, 2H), 3.52 (m, 2H), 3.32 (t, 1H), 3.17 (m, 1H), 2.91 (ddd, 2H), 2.69 (m, 1H), 2.36 (s, 3H), 2.27 (s, 3H), 1.04 (t, 3H). LRMS (APCI, Positive) m/e 431.9 (M+1).

Compound 293:




1-[5-Methyl-2-(1-methyl-piperidin-3-ylmethoxy)-phenyl]-3-quinoxalin-2-yl-urea

Step 1: Quinoxaline-2-carbonyl azide. A stirred solution quinoxaline-2-carboxylic acid (348 mg, 2 mmol) in THF (6 mL) at rt under nitrogen was treated with diisopropylethylamine (365 µL, 2.1 mmol) followed by diphenylphosphoryl azide (410 µL, 1.9 mmol). After stirring overnight the reaction was diluted to 60 mL with Et₂O and washed 2 x 60 mL with sat. NaCl. There was an insoluble brown oil, which was drained off with the aqueous layer and assumed to be a diphenyl phosphate impurity. The organics were dried (MgSO₄), filtered and concentrated to a tan solid, which corresponds to the acyl azide (350 mg, 92%). ¹H-NMR (400 MHz, CDCl₃) δ 9.58 (s, 1H), 8.32 (d, 1H), 8.21 (d, 1H), 7.94 (m, 2H)

Step 2: 1-[5-Methyl-2-(1-methyl-piperidin-3-ylmethoxy)-phenyl]-3-quinoxalin-2-yl-urea. A solution of quinoxaline-2-carbonyl azide (66 mg, 0.33 mmol) in toluene (1.7 mL) was stirred under nitrogen and immersed in a 90 °C heating bath. After 20 min. the reaction was cooled to 65 °C and solid 5-methyl-

2-(1-methyl-piperidin-3-ylmethoxy)-phenylamine (70 mg, 0.3 mmol) was added. The reaction was stirred at 65 °C for 4 hours and then allowed to cool to rt overnight. The resulting precipitate was collected by filtration and washed with toluene (62 mg, 51%). ¹H-NMR (400 MHz, CDCl₃) δ 11.58 (br s, 1H), 9.27 (br s, 1H), 8.63 (s, 1H), 8.22 (s, 1H), 8.03 (d, 1H), 7.91 (d, 1H), 7.76 (t, 1H), 7.61 (t, 1H), 6.86 (m, 2H), 4.03 (m, 2H), 2.91 (d, 1H), 2.61 (d, 1H), 2.37 (s, 3H), 2.25 (m, 1H), 2.09 (s, 1H), 1.81 (m, 3H), 1.57 (m, 2H), 1.05 (m, 1H). LRMS (APCI, Positive) m/e 405.9 (M+1).


Compound 294:



1-[5-Methyl-2-(pyridin-3-ylmethoxy)-phenyl]-3-quinoxalin-2-yl-urea

Step 1: 1-[5-Methyl-2-(pyridin-3-ylmethoxy)-phenyl]-3-quinoxalin-2-yl-urea. A stirred solution of quinoxaline-2-carbonyl azide (92 mg, 0.46 mmol, prepared as above) in 1.5 mL toluene under nitrogen was immersed in a 90 °C heating bath. After 20 min. the reaction was cooled to 65 °C and treated with solid 5-methyl-2-(pyridin-3-ylmethoxy)-phenylamine (90 mg, 0.42 mmol). The reaction was stirred at 65 °C for 4 hours and then allowed to cool to rt overnight. The resulting precipitate was collected by filtration. The crude product was chromatographed on a Biotage 12M column with 2/3 EtOAc/hexane to give pure urea as a tan solid (20 mg, 12%). ¹H-NMR (400 MHz, CDCl₃) δ 11.99 (br s, 1H), 9.64 (br s, 1H), 8.76 (s, 1H), 8.61 (s, 1H), 8.48 (d, 1H), 8.24 (s, 1H), 7.98 (d, 1H), 7.79 (d, 1H), 7.55 (t, 1H), 7.42 (t, 1H), 7.17 (d, 1H), 7.12 (m, 1H), 6.93 (m, 2H), 5.26 (s, 2H), 2.41 (s, 3H). LRMS (APCI, Positive) m/e 385.9 (M+1).

Compound 295:



Step 1 Mitsunobito procedure:

1-[2-(4-Methyl-2-nitro-phenoxy)-ethyl]-aziridine. A solution of 2-nitro-4-methylphenol (505 mg, 3.3 mmol, 1.1 eq.) and 2-aziridin-1-yl-ethanol (3.0 mmol, 1eq.) in 10 mL THF was stirred at 0 °C. Triphenylphosphine (0.87 g, 3.30 mmol, 1.1 eq.) and diisopropylazodicarboxylate, (0.67 g 3.30 mmol,

1.1 eq.) were added, and the solution was allowed to warm to temperature. After 18h, the reaction mixture was diluted with 100 mL EtOAc, and was washed with water (3 x 20 mL). The organic phase was washed again with 1 N HCl (3x 20 ml). The aqueous layer was basified with 3 N NaOH to pH > 12 and extracted with EtOAc, (3x 50 mL) to give crude product. The final product was purified by flash chromatography eluting with 5 - 10 % MeOH in dichloromethane. ¹H NMR (400 MHz, CDCl₃): δ 7.66 (s, 1H), 7.32 (d, *J* = 8.61 Hz, 1H), 7.01 (d, *J* = 8.61 Hz, 1H), 4.26 (t, *J* = 5.09 Hz, 2H), 2.67 (t, *J* = 5.48 Hz, 2H), 2.34 (s, 3H), 1.79 (m, 2H), 1.34 (m, 2H).


Step 2 Nitro reduction.

2-(2-Aziridin-1-yl-ethoxy)-5-methyl-phenylamine. A solution of 3-nitro-4-alkoxy toluene (1.0 mmol) in 20 mL EtOH was hydrogenated at 2 atm over 300 mg of 10% Pd on carbon for 30 minutes. The catalyst was removed by filtration through a glass fiber filter and the filtrate was concentrated to give the desired product, which was used directly without further purification.

Step 3 Urea formation

1-[2-(2-Aziridin-1-yl-ethoxy)-5-methyl-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea. A solution of 5-methylpyrazine-2-carboxylic azide (196 mg, 1.2 mmol, 1.2 eq.) in 20 mL anhydrous toluene was heated to 90 °C. After 20 minutes N₂ evolution had subsided, and the reaction mixture was cooled to 60 °C before adding aniline (1.0 mmol, 1 eq.) as a solution in 2 mL toluene. After stirring for 4 h at 60 °C. The reaction mixture was then partitioned between 50 mL EtOAc and sat. NaHCO₃. the organic phase was washed with brine, dried over MgSO₄, filtered and concentrated. The residue was purified by flash chromatography, eluting with 5 % MeOH in dichloromethane. ¹H NMR (400 MHz, *d*₆-DMSO): δ 10.80 (s, 1H), 8.64 (s, 1H), 8.53 (s, 1H), 8.15 (s, 1H), 8.07 (s, 1H), 6.82 (m, 2H), 4.2 (m, 2H), 2.7 (m, 2H), 2.5 (s, 3H), 2.32 (s, 3H), 1.89 (s, 2H), 1.30 (m, 2H). MS APCI-Pos, *M/e* 328.0 (*M*+1)


Compound 296:



1-[2-(3-Dimethylamino-benzyloxy)-5-methyl-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea

Prepared from (3-dimethylamino-phenyl)-methanol, as described above for compound 295. ¹H NMR (400 MHz, CDCl₃): δ 11.69 (s, 1H), 8.26 (s, 1H), 8.11 (s, 1H), 7.93 (s, 1H), 7.26 (m, 1H), 6.87 (m, 6H), 5.01 (s, 2H), 2.93 (s, 6H), 2.35 (s, 6H). MS APCI-Pos, *M/e* 391.9 (*M*+1)


Compound 297:



1-[2-(1-Isopropyl-pyrrolidin-3-yloxy)-5-methyl-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea

Prepared from 3-hydroxy-1-isopropyl-pyrrolidine, as described above for compound 295. ¹H NMR (400 MHz, *d*₆-DMSO): δ 10.08 (s, 2H), 8.65 (s, 1H), 8.14 (s, 1H), 8.03 (s, 1H), 6.81 (d, *J* = 7.83 Hz, 1H), 6.74 (d, *J* = 8.61 Hz, 1H), 4.85 (s, 1H), 2.91 (m, 1H), 2.75 (m, 2H), 2.48 (m, 1H), 2.4 (s, 3H), 2.36 (m, 1H), 2.27 (m, 1H), 2.21 (s, 3H), 1.85 (m, 1H), 1.01 (m, 6H). MS APCI-Pos, *M/e* 369.9 (*M*+1)


Compound 298:



1-[5-Methyl-2-(1-methyl-pyrrolidin-3-ylmethoxy)-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea.

Prepared from (1-methyl-pyrrolidin-3-yl)-methanol as described above for compound 295. ¹H NMR (400 MHz, *d*₆-DMSO): δ 10.13 (s, 2H), 8.67 (s, 1H), 8.15 (s, 1H), 8.06 (s, 1H), 6.81 (d, *J* = 8.61 Hz, 1H), 6.75 (d, *J* = 7.83 Hz, 1H), 4.88 (s, 1H), 2.75 (m, 4H), 2.5 (m, 2H), 2.43 (s, 3H), 2.3 (s, 3H), 2.24 (s, 3H). MS APCI-Pos, *M/e* 341.9 (*M*+1)

Compound 299:



Compound 134


Step 1: 3-Hydroxymethyl-piperidine-1-carboxylic acid *tert*-butyl ester. To a stirred solution of 3-hydroxymethyl piperidine (403 mg, 3.5 mmol, 1 eq.) in 20 mL of CH₂Cl₂ and 5 mL of sat'd NaHCO₃, 0 °C was added di-*tert*-butyl dicarbonate (803 mg, 3.68 mmol, 1.05 eq.) in several portions. After stirring at 0 °C for 2 h, the solution was diluted with 10 mL of water and was extracted with 2 x 20 mL

CH₂Cl₂ the combined extracts were washed with water, then brine and were dried over MgSO₄, filtered and concentrated to give the Boc protected amine which was used in the next step.

Steps 2-4: **3-[4-Methyl-2-[3-(5-methyl-pyrazin-2-yl)-ureido]-phenoxy-methyl]-piperidine-1-carboxylic acid tert-butyl ester** was prepared from 3-hydroxymethyl-piperidine-1-carboxylic acid tert-butyl ester, as described above for compound 295. It was purified by flash chromatography eluting with 5% MeOH in CH₂Cl₂

Step 5: **1-[5-Methyl-2-(piperidin-3-ylmethoxy)-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea** Removal of the Boc group was accomplished by treatment of a 0 °C solution of the protected derivative (180 mg, 0.395 mmol) in 15 mL CH₂Cl₂ with 2 mL of TFA. After stirring for 18h at room temperature, the reaction was concentrated in *vacuo*, and the residue was taken up in 20 mL of EtOAc and was washed with 10 mL of NaHCO₃. The aqueous phase was extracted with EtOAc (2 x 30 mL), and the combined extracts were washed with 20 mL of brine, dried over MgSO₄, filtered and concentrated to give 128 mg (91%) of the desired amine. ¹H NMR (400 MHz, *d*₆-DMSO): δ 10.16 (s, 1H), 10.09 (s, 1H), 8.53 (s, 1H), 8.11 (s, 1H), 7.97 (s, 1H), 6.8 (d, *J* = 7.8 Hz, 1H), 6.69 (d, *J* = 8.6 Hz, 1H), 3.77 (s, 2H), 3.09 (m, 1H), 2.82 (m, 1H), 2.34 (m, 2H), 2.3 (s, 3H), 2.27 (m, 1H), 2.15 (s, 3H), 1.92 (m, 1H), 1.75 (m, 1H), 1.53 (m, 1H), 1.37 (m, 1H), 1.11 (m, 1H). MS APCI-Pos, *M/e* 356.0 (*M*+1)

Compound 300:




1-[5-Fluoro-2-(pyridin-3-ylmethoxy)-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea

Step 1: **3-(4-Fluoro-2-nitro-phenoxy-methyl)-pyridine**. To a stirred, cooled (about 0 °C) solution of 1,4-Difluoro-2-nitro-benzene (3.0 mmol) and 3-pyridylcarbinol (3.1 mmol) in tetrahydrofuran (8 mL) was added lithium bis(trimethylsilyl)amide (3.2 mmol; 3.2 mL of a 1.0 M solution in tetrahydrofuran). After stirring for 12 hours, the reaction was diluted with 30 mL of ethyl acetate and washed with 30 mL of 10% aqueous sodium carbonate (2 x 30 mL), brine (1 x 30 mL), then dried (MgSO₄), and filtered. The filtered solution was concentrated under reduced pressure to provide the desired crude product.

Step 2: **5-Fluoro-2-(pyridin-3-ylmethoxy)-phenylamine**. To a stirred, cooled (about 0 °C) solution of a 4-Fluoro-2-nitro-phenoxy-methyl-pyridine (1.0 mmol) in methanol (2 mL) and saturated aqueous ammonium chloride (1 mL) was added zinc dust (2.0 mmol). After stirring for 12 hours, the reaction was diluted with 30 mL of ethyl acetate and washed with 30 mL of 10% aqueous sodium carbonate (2 x 30 mL), brine (1 x 30 mL), then dried (MgSO₄), and filtered. The filtered solution was concentrated under reduced pressure to provide the desired crude product.

Step 3: Urea formation according to method for compound 295 (23% yield). ^1H NMR (400 MHz, CDCl_3) δ 11.62 (br s, 1H), 8.80 (s, 1H), 8.75 (d, 1H), 8.43 (br s, 1H), 8.25 (d, 1H), 8.15 (s, 1H), 7.85 (d, 1H), 7.43 (m, 1H), 6.95 (m, 2H), 6.68 (m, 1H), 5.15 (s, 2H), 2.43 (s, 3H). LRMS (ESI, Positive) m/e 354.10 ($M+1$).

Compound 301:




1-[5-Fluoro-2-(1-methyl-piperidin-3-ylmethoxy)-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea

Steps 1-2: According to procedure for compound 300, using 1,4-difluoro-2-nitrobenzene and 1-methyl-3-hydroxymethyl piperidine.

Step 3: Urea formation according to method for compound 285 (62% yield). ^1H NMR (400 MHz, CDCl_3) δ 8.22 (m, 3H), 7.21 (m, 2H), 6.78 (m, 2H), 3.85 (m, 2H), 3.21 (m, 1H), 2.85 (m, 1H), 2.52 (s, 3H), 2.39 (s, 3H), 1.50-2.30 (m, 8H). LRMS (ESI, Positive) m/e 374.21 ($M+1$).

Compound 302:




1-[5-Fluoro-2-(1-methyl-piperidin-4-yloxy)-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea

Steps 1-2: According to procedure for compound 300, using 1,4-difluoro-2-nitrobenzene and 1-methyl-4-hydroxypiperidine.

Step 3: Urea formation according to method for compound 295 (78% yield). ^1H NMR (400 MHz, CDCl_3) δ 11.49 (br s, 1H), 8.89 (br s, 1H), 8.35 (s, 1H), 8.22 (d, 1H), 8.10 (s, 1H), 6.80 (m, 1H), 6.70 (m, 1H), 4.25 (m, 1H), 2.90 (m, 2H), 2.55 (s, 3H), 2.38 (s, 3H), 2.35 (s, 3H), 1.80-2.30 (m, 6H). LRMS (ESI, Positive) m/e 359.91 ($M+1$).

Compound 303:



1-[4-Fluoro-2-(pyridin-3-ylmethoxy)-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea


Step 1: 3-(5-Fluoro-2-nitro-phenoxy)methyl-pyridine To a stirred, cooled (about 0 °C) solution of 2-nitro-5-fluoro-phenol (2.0 mmol), triphenyl phosphine (2.0 mmol), and 3-hydroxymethylpyridine (2.0 mmol) in dry tetrahydrofuran (5 mL) was added diisopropyl azodicarboxylate (2.0 mmol in 1 mL of tetrahydrofuran). After stirring for 12 hours, the reaction was diluted with 30 mL of ethyl acetate and washed with 30 mL of 10% aqueous sodium carbonate (2 x 30 mL), brine (1 x 30 mL), then dried (MgSO₄), and filtered. The filtered solution was concentrated under reduced pressure to provide the desired crude product.

Step 2: 4-Fluoro-2-(pyridin-3-ylmethoxy)-phenylamine Nitro reduction according to the method for compound 300.

Step 3: Urea formation according to method for compound 295 (60% yield).

¹H NMR (400 MHz, CDCl₃) δ 11.41 (br s, 1H), 8.85 (s, 1H), 8.75 (d, 1H), 8.40 (t, 1H), 8.18 (s, 1H), 7.88 (s, 1H), 7.80 (d, 1H), 7.43 (t, 1H), 7.00 (s, 1H), 6.80 (m, 2H), 5.12 (s, 2H), 2.43 (s, 3H). LRMS (ESI, Positive) m/e 354.21(M+1).

Compound 304:




1-[4-Fluoro-2-(1-methyl-piperidin-3-ylmethoxy)-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea

Prepared according to the methods for compound 303, using 2-nitro-5-fluorophenol and 1-methyl-3-hydroxymethyl piperidine.

¹H NMR (400 MHz, CDCl₃) δ 8.50 (br s, 1H), 8.19 (m, 2H), 6.65 (m, 2H), 3.85 (m, 2H), 3.60 (s, 3H), 2.80-3.20 (m, 2H), 2.54 (s, 3H), 2.39 (s, 3H), 1.60-2.10 (m 5H). LRMS (ESI, Positive) m/e 373.95 (M+1).


Compound 305:



1-[4-Fluoro-2-(1-methyl-piperidin-4-yloxy)-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea

Prepared according to the methods for compound 303, using 2-nitro-5-fluorophenol and 1-methyl-4-hydroxypiperidine. ¹H NMR (400 MHz, CDCl₃) δ 11.35 (br s, 1H), 9.49 (s, 1H), 8.35 (m, 2H), 8.05 (s, 1H), 6.65 (m, 2H), 4.35 (m, 1H), 2.90 (m, 2H), 2.54 (s, 3H), 2.35 (s, 3H), 1.80-2.30 (m, 6H). LRMS (ESI, Positive) m/e 359.93 (M+1).

Compound 306:




1-[3,4-Difluoro-2-(pyridin-3-ylmethoxy)-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea

Step 1: 3-(2,3-Difluoro-6-nitro-phenoxy)methyl-pyridine To a stirred, cooled (about 0 °C) solution of 2,3-difluoro-6-nitrophenol (2.0 mmol), triphenyl phosphine (2.0 mmol), and 3-hydroxymethylpyridine (2.0 mmol) in dry tetrahydrofuran (5 mL) was added diisopropyl azodicarboxylate (2.0 mmol in 1 mL of tetrahydrofuran). After stirring for 12 hours, the reaction was diluted with 30 mL of ethyl acetate and washed with 30 mL of 10% aqueous sodium carbonate (2 x 30 mL), brine (1 x 30 mL), then dried (MgSO₄) and filtered. The filtered solution was concentrated under reduced pressure to provide the desired crude product.

Step 2: 3,4-Difluoro-2-(pyridin-3-ylmethoxy)-phenylamine Nitro reduction according to the method for compound 300.

Step 3: Urea formation according to method for compound 295 (20% yield). ¹H NMR (400 MHz, CDCl₃) δ 11.49 (br s, 1H), 8.89 (s, 1H), 8.85 (s, 1H), 8.65 (d, 1H), 8.25 (s, 1H), 8.10 (m, 1H), 7.88 (d, 1H), 7.35 (t, 1H), 7.18 (s, 1H), 6.98 (m, 1H), 5.25 (s, 2H), 2.52 (s, 3H). LRMS (ESI, Positive) m/e 372.10 (M+1).

Compound 307:




1-(5-Methyl-pyrazin-2-yl)-3-[5-methyl-2-(pyridin-4-yloxy)-phenyl]-urea

Step 1: 5-Methyl-2-(pyridin-4-yloxy)-phenylamine To a stirred solution of 2-amino-4-methyl phenol (616 mg; 5.0 mmol) and 4-chloro pyridine (625 mg; 5.5 mmol) in dimethyl sulfoxide (5 mL) was added sodium hydroxide (600 mg; 15.0 mmol, in 1 mL of water). The reaction was heated to 100 °C and stirred for 12 hours. The reaction was cooled to room temperature, diluted with 50 mL of ethyl acetate and washed with aqueous 10% sodium carbonate (1 x 50 mL), and brine (50 mL), then dried (MgSO₄), and filtered. The crude product was purified using the biotage 40M cartridge eluting with methylene chloride:methanol:ammonia (90:8:2) to yield a light yellow oil (10% yield).

Step 2: Urea formation according to method for compound 295 (36% yield). ¹H NMR (400 MHz, CDCl₃) δ 11.41 (br s, 1H), 8.52 (m, 3H), 8.33 (s, 1H), 8.22 (s, 1H), 7.61 (s, 1H), 6.80-7.00 (m, 4H), 2.49 (s, 3H), 2.45 (s, 3H). LRMS (ESI, Positive) m/e 335.91 (M+1).

Compound 308:



1-(5-Methyl-pyrazin-2-yl)-3-[5-methyl-2-(pyridin-3-yloxy)-phenyl]-urea


Step 1: 3-(4-Methyl-2-nitro-phenoxy)-pyridine To a stirred solution of 1-chloro-4-methyl-2-nitrobenzene (686 mg; 4.0 mmol) and pyridin-3-ol (418 mg; 4.40 mmol) in dimethylformamide (5 mL) was added potassium carbonate (1.22 g, 8.80 mmol). The reaction was heated to 50 °C and stirred for 12 hours. The reaction was cooled to room temperature, diluted with 50 mL of ethyl acetate and washed with aqueous 10% sodium carbonate (1 x 50 mL), and brine (50 mL), then dried (MgSO₄), and filtered. The crude product was purified using the biotage 40M cartridge eluting with hexanes and ethyl acetate (1:1) to yield a light yellow oil (27% yield).

Step 2: 5-Methyl-2-(pyridin-3-yloxy)-phenylamine To a stirred, cooled (about 0 °C) solution of 3-(4-Methyl-2-nitro-phenoxy)-pyridine (1.0 mmol) in methanol (2 mL) and saturated aqueous ammonium chloride (1mL) was added zinc dust (2.0 mmol). After stirring for 12 hours, the reaction was diluted with 30 mL of ethyl acetate and washed with 30 mL of 10% aqueous sodium carbonate (2 x 30 mL), brine (1 x 30 mL), then dried (MgSO₄), and filtered. The filtered solution was concentrated under reduced pressure to yield a brown oil (95% yield).

Step 3: Urea formation according to method for compound 295 (45% yield). ¹H NMR (400 MHz, CDCl₃) δ 11.49 (br s, 1H), 8.55 (s, 1H), 8.39 (d, 1H), 8.35 (s, 1H), 8.15 (s, 1H), 8.05 (br s, 1H), 7.21 (m, 2H), 6.92 (m, 2H), 2.49 (s, 3H), 2.45 (s, 3H).

LRMS (ESI, Positive) m/e 335.91 (M+1).

Compound 309:



1-(5-Methyl-pyrazin-2-yl)-3-[5-methyl-2-(pyridin-2-yloxy)-phenyl]-urea

Step 1: **2-(4-Methyl-2-nitro-phenoxy)-pyridine** To a stirred solution of 1-chloro-4-methyl-2-nitrobenzene (686 mg; 4.0 mmol) and pyridin-2-ol (418 mg; 4.40 mmol) in dimethylformamide (5 mL) was added potassium carbonate (1.22 g, 8.80 mmol). The reaction was heated to 50 °C and stirred for 12 hours. The reaction was cooled to room temperature, diluted with 50 mL of ethyl acetate and washed with aqueous 10% sodium carbonate (1 x 50 mL), and brine (50 mL), then dried (MgSO₄) and filtered. The crude product was purified using the biotage 40M cartridge eluting with hexanes and ethyl acetate (1:1) to yield a light yellow oil (11% yield).


Step 2: **5-Methyl-2-(pyridin-2-yloxy)-phenylamine** To a stirred, cooled (about 0 °C) solution of 2-(4-Methyl-2-nitro-phenoxy)-pyridine (1.0 mmol) in methanol (2 mL) and saturated aqueous ammonium chloride (1mL) was added zinc dust (2.0 mmol). After stirring for 12 hours, the reaction was diluted with 30 mL of ethyl acetate and washed with 30 mL of 10% aqueous sodium carbonate (2 x 30 mL), brine (1 x 30 mL), then dried (MgSO₄) and filtered. The filtered solution was concentrated under reduced pressure to yield a white foam (77% yield).

Step 3: Urea formation according to method for compound 295 (43% yield). ¹H NMR (400 MHz, CDCl₃) δ 8.51 (br s, 1H), 8.42 (br s, 1H), 8.00 (s, 1H), 7.80 (s, 1H), 7.51 (t, 1H), 7.29 (d, 1H), 7.05 (d, 1H), 6.95 (d, 1H), 6.75 (d, 1H), 6.35 (t, 1H), 2.49 (s, 3H), 2.45 (s, 3H). LRMS (ESI, Positive) m/e 335.91 (M+1).

Substituted Aminopyrazine Ureas. General Procedure:

To a 0.3 M stirred solution of the aminopyrazine derivative (1 equiv.) in dichloroethane at room temperature under nitrogen was added 2-methoxy-5-methylphenylisocyanate (1 equiv.). The reaction was warmed to 80 °C overnight and then cooled to room temperature. In most cases, the product precipitated and was isolated by filtration. Alternatively the product could be isolated by silica gel chromatography using EtOAc/hexane or CH₂Cl₂/MeOH as eluant.

Compound 310:




3-(2-Methoxy-5-methyl-phenyl)-1-methyl-1-pyrazin-2-yl-urea

Step 1: 2-methylaminopyrazine. To a stirred solution of 2M methylamine in 1 mL of methanol, at room temperature, was added 2-chloropyrazine. The reaction was sealed and heated to 60 °C for 24 hours. The reaction was concentrated to a mixture of starting material and the desired 2-methylaminopyrazine in a 1:2 ratio. The material was used crude in a urea forming reaction. ¹H-NMR (400 MHz, CDCl₃) δ 7.97 (d, 1H), 7.85 (s, 1H), 7.73 (d, 1H), 2.96 (s, 3H)

Step 2: To a 0.3 M stirred solution of 2-methylaminopyrazine (1 equiv.) in dichloroethane at room temperature under nitrogen was added 2-methoxy-5-methylphenylisocyanate (1 equiv.). The reaction was warmed to 80 °C overnight and then cooled to room temperature. In most cases, the product precipitated and was isolated by filtration. Alternatively the product could be isolated by silica gel chromatography using EtOAc/hexane or CH₂Cl₂/MeOH as eluant. ¹H-NMR (400 MHz, CDCl₃) δ 8.57 (s, 1H), 8.32 (s, 1H), 8.31 (s, 1H), 8.17 (s, 1H), 6.80 (m, 2H), 3.90 (s, 3H), 3.56 (s, 3H), 2.32 (s, 3H). LRMS (ESI, Positive) m/e 273.2 (M+1).


Compound 311:



1-(2-Methoxy-5-methyl-phenyl)-3-(5-methyl-pyrazin-2-yl)-urea

Prepared according to general procedure described for compound 310, using 2-amino-4-methylpyrazine. ¹H-NMR (400 MHz, CDCl₃) δ 11.12 (br s, 1H), 8.28 (s, 1H), 8.17 (s, 1H), 8.09 (s, 1H), 6.81 (m, 2H), 3.91 (s, 3H), 2.54 (s, 3H), 2.35 (s, 3H). LRMS (ESI, Positive) m/e 273.2 (M+1).

Compound 312:




1-(5,6-Dimethyl-pyrazin-2-yl)-3-(2-methoxy-5-methyl-phenyl)-urea

Step 1: 2-Amino-5,6-dimethylpyrazine. Glycine amidine dihydrobromide (620 mg, 2.64 mmol) was stirred in 6 mL of MeOH at -30°C (acetonitrile/ CO_2 bath) in a capped flask. Butanedione (232 μL , 2.64 mmol) was stirred separately in 6 mL H_2O with sodium acetate (700 mg) until homogeneous. The diketone was added to the amidine solution by pipet followed by 2.5 mL of 3.6 M NaOH. The yellow solution was allowed to warm slowly to RT and was then stirred overnight. MeOH was removed by rotovap and the aqueous solution extracted 3 x 30 mL with EtOAc. The combined organic extracts were dried (MgSO_4), filtered and concentrated to a yellow solid which contained some impurities. The solid was triturated with EtOAc/ Et_2O and filtered to give pure compound (55 mg, 17%). $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 7.76 (s, 1H), 4.25 (br s, 2H), 2.40 (s, 3H), 2.37 (s, 3H)

Step 2: Prepared according to general procedure described for compound 310, using 2-amino-5,6-dimethylpyrazine. $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 11.43 (br s, 1H), 8.23 (s, 1H), 8.00 (s, 1H), 7.64 (br s, 1H), 6.81 (m, 2H), 3.95 (s, 3H), 2.59 (s, 3H), 2.52 (s, 3H), 2.35 (s, 3H). LRMS (APCI, Positive) m/e 287.1 ($\text{M}+1$).

Compound 313:




1-(2-Methoxy-5-methyl-phenyl)-3-(5-trifluoromethyl-pyrazin-2-yl)-urea

Step 1: 2-amino-5-trifluoromethylpyrazine. Prepared according to the method of Miesel, J. U.S. patent 4,293,552 (1981),

Step2: Prepared according to general procedure described for compound 310, using 2-amino-5-trifluoromethylpyrazine. $^1\text{H-NMR}$ (400 MHz, $d_6\text{-DMSO}$) δ 10.59 (s, 1H), 9.78 (br s, 1H), 9.06 (s, 1H), 8.80 (s, 1H), 8.00 (s, 1H), 6.96 (d, 1H), 6.81 (d, 1H), 3.87 (s, 3H), 2.22 (s, 3H). LRMS (ESI, Positive) m/e 327.1 ($\text{M}+1$).

Compound 314:



1-(5,6-Diphenyl-pyrazin-2-yl)-3-(2-methoxy-5-methyl-phenyl)-urea

Step 1: 2-Hydroxy-5,6-diphenylpyrazine. To a stirred suspension of glycineamide hydrochloride (1.1 gm, 10 mmol) in 20 mL MeOH at 0°C was added 20% NaOH (10 mL, 50 mmol). A clear solution

formed and was treated slowly portionwise with benzil (2.1 gm, 10 mmol) as a solid. The yellow solution was stirred at 0 °C for 2 hours and then neutralized to approximately pH=7 with concentrated HCl. The bright yellow color disappeared and a tan precipitate formed. The material was isolated by filtration with MeOH and triturated with EtOAc to give 2-hydroxy-5,6-diphenylpyrazine (2 gm, 80%). ¹H-NMR (400 MHz, CDCl₃) δ 8.24 (s, 1H), 7.42-7.31 (m, 4H), 7.39-7.21 (m, 6H).


Step 2: 2-Chloro-5,6-diphenylpyrazine. A stirred solution of 2-Hydroxy-5,6-diphenylpyrazine (430 mg, 1.7 mmol) in 5.2 mL POCl₃ in a capped reaction vial was heated to 100 °C for 4 hours. The orange solution was cooled to room temperature and stirred rapidly in a mixture of CH₂Cl₂ (100 mL) and ice cold 10% Na₂CO₃ (100 mL) for 15 minutes. The organic layer was isolated and washed 2 x 100 mL with 10% Na₂CO₃. The organics were isolated, dried (MgSO₄), filtered and concentrated to the chloropyrazine, which existed as white solid (450 mg, quantitative). ¹H-NMR (400 MHz, CDCl₃) δ 8.59 (s, 1H), 7.45-7.39 (m, 4H), 7.36-7.24 (m, 6H).

Step 3: 2-Azido-5,6-diphenylpyrazine. To a stirred solution of 2-Chloro-5,6-diphenylpyrazine (45 mg, 0.17 mmol) in 500 µL DMF at room temperature under nitrogen was added sodium azide (11 mg, 0.17 mmol) and the reaction was warmed to 100 °C. After stirring overnight, the reaction was cooled to room temperature, diluted with EtOAc (30 mL) and washed 4 x 30 mL with H₂O and 1 x 30 mL with saturated NaCl. The organics were isolated, dried (MgSO₄), filtered and concentrated to the 2-azidopyrazine, which exists as a yellow solid (45 mg, quantitative). ¹H-NMR (400 MHz, CDCl₃) δ 9.73 (s, 1H), 7.58-7.42 (m, 6H), 7.36-7.23 (m, 4H).

Step 4: 2-Amino-5,6-diphenylpyrazine. To a stirred solution of 2-Azido-5,6-diphenylpyrazine (45 mg, 0.17 mmol) in 50 mL EtOAc at room temperature was added triethylamine (100 µL) followed by Pearlman's Catalyst (50 mg). The suspension was put through a vacuum/purge cycle three times with hydrogen gas and then held under 1 atmosphere of hydrogen for 2 hours. The suspension was then filtered through GF/F filter paper with EtOAc and concentrated. The crude product was eluted through a Biotage 12S column with 1/1 EtOAc/hexane to give pure product, as a clear oil (25 mg, 59%). ¹H-NMR (400 MHz, CDCl₃) δ 8.04 (s, 1H), 7.42-7.20 (m, 10H), 4.62 (br s, 2H)

Step 5: Prepared according to general procedure described for compound 310, using 2-amino-5,6-diphenylpyrazine. ¹H-NMR (400 MHz, CDCl₃) δ 8.34 (s, 1H), 8.13 (s, 1H), 7.80 (s, 1H), 7.46 (d, 2H), 7.37-7.23 (m, 10H), 6.81 (d, 1H), 6.66 (d, 1H), 3.17 (s, 3H), 2.33 (s, 3H)


Compound 315:



1-[3-Benzyl-5-(4-methoxy-phenyl)-pyrazin-2-yl]-3-(2-methoxy-5-methyl-phenyl)-urea

Prepared according to general procedure described for compound 310, using 2-amino-3-benzyl-4-(4-methoxyphenyl)pyrazine ¹H-NMR (400 MHz, CDCl₃) δ 8.51 (s, 1H), 8.11 (s, 1H), 7.96 (d, 2H), 7.34 (m, 5H), 7.03 (d, 2H), 6.80 (m, 2H), 4.28 (s, 2H), 3.95 (s, 3H), 3.90 (s, 3H), 2.30 (s, 3H). LRMS (ESI, Positive) m/e 477.2 (M+1).

Compound 316:




1-(6-Azido-pyrazin-2-yl)-3-(2-methoxy-5-methyl-phenyl)-urea

Step 1: Tetrazolo[1,5-a]pyrazin-5-ylamine. Prepared according to the method of Shaw, J.T.; et al. J. Heterocyclic Chem. 1980, 17, 11.

Step 2: Prepared using *p*-nitrophenyl carbamate general procedure described for compound 166 (step 2) using Tetrazolo[1,5-a]pyrazin-5-ylamine. ¹H-NMR (400 MHz, CDCl₃) δ 9.72 (br s, 1H), 8.23 (s, 1H), 7.97 (s, 1H), 7.87 (s, 1H), 7.80 (br s, 1H), 6.88 (d, 1H), 6.80 (d, 1H), 3.84 (s, 3H), 2.36 (s, 3H). LRMS (ESI, Positive) m/e 300.0 (M+1).

Compound 317:




1-(6-Amino-pyrazin-2-yl)-3-(2-methoxy-5-methyl-phenyl)-urea

To a stirred solution of 1-(6-azido-pyrazin-2-yl)-3-(2-methoxy-5-methyl-phenyl)-urea (8 mg, 27 μmol) 95% EtOH (2mL) at room temperature was added concentrated NH₄OH (10 μL) and 10% Pd on C (25 mg). The suspension was put through a vacuum/purge cycle three times with hydrogen gas and then held under 50 psi of hydrogen pressure and shaken on a Parr Shaker. After 2 hours the vacuum/purge cycle was repeated and the reaction held under hydrogen for another 2 hours. The suspension was then filtered through GF/F filter paper with EtOH and concentrated to a yellow film (3 mg, 41%). ¹H-NMR

(400 MHz, CDCl₃) δ 8.19 (s, 1H), 7.61 (s, 1H), 7.56 (s, 1H), 6.83 (s, 2H), 3.95 (s, 3H), 2.33 (s, 3H). LRMS (ESI, Positive) m/e 274.2 (M+1).


Compound 318:



1-(6-Chloro-pyrazin-2-yl)-3-(2-methoxy-5-methyl-phenyl)-urea

To a stirred solution of 2-amino-6-chloropyrazine (130 mg, 1 mmol) in 3 mL THF at 0 °C under nitrogen was added methyl magnesium iodide (3M in Et₂O, 330 μ L, 1 mmol) to give a yellow suspension that was stirred at 0 °C for 15 minutes. The suspension was treated with the isocyanate neat (147 μ L, 1 mmol) and allowed to warm to room temperature overnight. The reaction was partitioned between EtOAc (30 mL) and 10% Na₂CO₃ (30 mL). The organics were isolated and washed 1 x 30 mL with 10% Na₂CO₃ and 1 x 30 mL with saturated NaCl. The organics were dried (MgSO₄), filtered and concentrated to a crude residue that was triturated with EtOAc to give, after filtration, the urea product as a white solid (27 mg, 9%). ¹H-NMR (400 MHz, CDCl₃) δ 8.26 (s, 1H), 8.23 (s, 1H), 8.17 (s, 1H), 8.09 (br s, 1H), 6.84 (d, 1H), 6.81 (d, 1H), 3.96 (s, 3H), 2.35 (s, 3H). LRMS (ESI, Positive) m/e 293.0 (M+1).

Compound 319:



1-(5-Bromo-pyrazin-2-yl)-3-(2-methoxy-5-methyl-phenyl)-urea

Step 1: 2-amino-5-bromopyrazine. To a stirred, cooled (0°C) solution of amino pyrazine (5.0 g, 52.6 mmol) in methylene chloride (200 mL) was added *N*-bromosuccinimide (9.39 g, 52.8 mmol). After stirring for 24 hours, the reaction was washed with aqueous 10% sodium carbonate (3 x 50 mL), water (50 mL), then dried (MgSO₄), and filtered. The filtered material was concentrated under reduced pressure, taken up in minimal ethyl acetate (5 mL) followed by hexanes (200 mL). Yellow crystals formed which were filtered and dried. (56 % yield).

Step 2: Prepared according to general procedure described for compound 310, using 2-amino-4-bromopyrazine. ¹H-NMR (400 MHz, CDCl₃/CD₃OD) δ 8.55 (s, 1H), 8.32 (s, 1H), 8.03 (s, 1H), 6.81 (m, 2H), 3.92 (s, 3H), 2.34 (s, 3H).

COc1ccc(NC(=O)Nc2nc(Br)nc(Br)n2)cc1

Prepared according to general procedure described for compound 310, using 2-amino-4,6-dibromopyrazine. ¹H-NMR (400 MHz, CDCl₃) δ 7.98 (s, 1H), 7.13 (s, 1H), 6.79 (m, 2H), 3.83 (s, 3H), 2.32 (s, 3H)

COc1ccc(NC(=O)Nc2ncnc3c2CN(C3)c4c(=O)c5ccccc5c4=O)cc1


Step 1: **(5-Bromomethyl-pyrazin-2-yl)-carbamic acid tert-butyl ester**. To a stirred solution of 2-Boc-amino-5-methyl pyrazine (1.34 gm, 6.4 mmol), in 20 mL CCl_4 at room temperature under nitrogen was added *N*-bromosuccinimide (1.14 gm, 6.4 mmol) followed by benzoyl peroxide (125 mg). The solution was irradiated with a 100 watt flood lamp, which caused the reaction to reflux vigorously. After 2 hours, the reaction was cooled to room temperature, diluted to 125 mL with CH_2Cl_2 and washed 1 x 125 mL with 10% sodium bisulfite solution and 1 x 125 mL with saturated NaCl. The organics were dried (MgSO_4), filtered and concentrated to a brown oil, which was directly loaded onto a Biotage 40S column with CH_2Cl_2 and eluted with 15/85 EtOAc/hexane to give the desired benzylic bromide as a yellow solid (954 mg, 51%). $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 9.22 (s, 1H), 8.29 (s, 1H), 7.37 (br s, 1H), 4.54 (s, 2H), 1.55 (s, 9H).

Step 2: [5-(1,3-Dioxo-1,3-dihydro-isindol-2-ylmethyl)-pyrazin-2-yl]-carbamic acid tert-butyl ester. To a stirred solution of phthalimide (971 mg, 6.6 mmol) and powdered K_2CO_3 (1.37 gm, 9.9 mmol) in acetonitrile (9.9 mL) at room temperature under nitrogen was added the bromide (954 mg, 3.3 mmol) as a solid. The suspension was heated to 65°C for 4 hours. After cooling to room temperature the reaction was partitioned between EtOAc (60 mL) and H_2O (60 mL). The organics were isolated and washed 2 x 50 mL with H_2O and 1 x 50 mL with saturated NaCl. The organics were dried ($MgSO_4$), filtered and concentrated. The crude product was triturated with CH_2Cl_2 and filtered to remove solid excess phthalimide and the filtrate partially concentrated and loaded directly onto a Biotage 40S column and eluted with 3/7 EtOAc/hexane to give the desired phthalimide as a white solid (495 mg, 42%). 1H -NMR (400 MHz, $CDCl_3$) δ 9.17 (s, 1H), 8.23 (s, 1H), 7.85 (m, 2H), 7.71 (m, 2H), 7.39 (br s, 1H), 4.97 (s, 2H), 1.52 (s, 9H).

Step 3: 2-(5-Amino-pyrazin-2-ylmethyl)-isoindole-1,3-dione. To a stirred solution of the phthalimide (495 mg, 1.4 mmol) in 7 mL CH_2Cl_2 at room temperature in a capped flask was added trifluoroacetic acid (7 mL). After stirring overnight, the reaction was concentrated to remove excess trifluoroacetic acid and was then dissolved in 200 mL 10/1 $\text{CH}_2\text{Cl}_2/\text{MeOH}$, stirred rapidly, and treated with a solution of 10% Na_2CO_3 (200 mL). The organics were isolated, dried (MgSO_4), filtered and concentrated to give the free aminopyrazine as a yellow solid (260 mg, 73%). $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 8.25 (s, 1H), 7.85 (m, 2H), 7.77 (s, 1H), 7.74 (m, 2H), 4.83 (s, 2H)

Step 4: Prepared according to general procedure described for compound 310 using 2-(5-Amino-pyrazin-2-ylmethyl)-isoindole-1,3-dione. $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 8.38 (s, 1H), 8.29 (s, 1H), 8.03 (s, 1H), 7.86 (m, 2H), 7.76 (m, 2H), 6.81 (m, 2H), 4.98 (s, 2H), 3.91 (s, 3H), 2.31 (s, 3H). LRMS (ESI, Positive) m/e 418.1 (M+1).


Compound 322:



1-(5-Aminomethyl-pyrazin-2-yl)-3-(2-methoxy-5-methyl-phenyl)-urea

To a stirred solution of 1-[5-(1,3-dioxo-1,3-dihydro-isoindol-2-ylmethyl)-pyrazin-2-yl]-3-(2-methoxy-5-methyl-phenyl)-urea (16 mg, 38 μmol) in 380 μL 95% EtOH and 100 μL DMF at room temperature in a capped reaction vial was added hydrazine monohydrate (3.8 μL , 76 μmol). After stirring overnight at room temperature, a white precipitate formed. The precipitate was filtered off, dried and triturated with EtOAc to remove phthalimide based impurities to provide the product as a white solid (7.9 mg, 72%). $^1\text{H-NMR}$ (400 MHz, $d_6\text{-DMSO}$) δ 9.54 (s, 1H), 8.80 (s, 1H), 8.46 (s, 1H), 8.02 (s, 1H), 6.93 (d, 1H), 6.79 (d, 1H), 4.51 (d, 2H), 4.39 (br s, 2H), 3.89 (s, 3H), 2.23 (s, 3H). LRLCMS (ESI, Positive) m/e 288.2 (M+1).

Compound 323:




1-(2-Methoxy-5-methyl-phenyl)-3-(6-methoxy-pyrazin-2-yl)-urea

Step 1: 2-amino-6-methoxypyrazine. To a stirred solution of methanol (89 μL ; 2.2 mmol) in dioxane (1 mL) was added sodium hydride (53 mg; 2.2 mmol). After stirring for 30 minutes, 2-amino-6-chloropyrazine (258 mg; 2.0 mmol) was added and the reaction was heated to 90 $^\circ\text{C}$. After stirring for 12 hours, the reaction was cooled to room temperature, diluted with 30 mL of ethyl acetate and washed with aqueous 10% sodium carbonate (1 x 30 mL), and brine (30 mL), then dried (MgSO_4), and filtered.

The crude product was purified using the Biotage 12i cartridge eluting with hexane and ethyl acetate (3:1) to yield a white solid (11% yield).

Step 2: Prepared according to general procedure described for compound 310 using 2-amino-6-methoxypyrazine. (8% yield). ¹H NMR (400 MHz, CDCl₃) δ 8.15 (s, 1H), 8.05 (br s, 1H), 7.92 (s, 1H), 7.82 (s, 1H), 6.89 (d, 1H), 6.80 (d, 1H), 4.05 (s, 3H), 3.81 (s, 3H), 2.38 (s, 3H). LRMS (ESI, Positive) m/e 289.10 (M+1).

Compound 324:




1-(6-Benzyloxy-pyrazin-2-yl)-3-(2-methoxy-5-methyl-phenyl)-urea

Step 1: **2-amino-6-benzyloxy-pyrazine**. To a stirred solution of benzyl alcohol (432 µL; 4.0 mmol) in dioxane (2 mL) was added sodium hydride (96 mg; 4.0 mmol). After stirring for 30 minutes, 2-amino-6-chloropyrazine (258 mg; 2.0 mmol) was added and the reaction was heated to 90 °C. After stirring for 12 hours, the reaction was cooled to room temperature, diluted with 30 mL of ethyl acetate and washed with aqueous 10% sodium carbonate (1 x 30 mL), and brine (30 mL), then dried (MgSO₄) and filtered. The crude product was purified using the biotage 12i cartridge eluting with hexane and ethyl acetate (3:1) to yield a white solid (33% yield).

Step 2: Prepared according to general procedure described for compound 310 using 2-amino-6-benzyloxy-pyrazine. (34% yield). ¹H NMR (400 MHz, d₆-DMSO) δ 9.99 (s, 1H), 9.18 (s, 1H), 8.62 (s, 1H), 7.99 (s, 1H), 7.95 (s, 1H), 7.52 (d, 2H), 7.41 (m, 3H), 6.92 (d, 1H), 6.80 (d, 1H), 5.39 (s, 2H), 3.80 (s, 3H), 2.21 (s, 3H). LRMS (ESI, Positive) m/e 365.10 (M+1).

Compound 325:




1-(2-Methoxy-5-methyl-phenyl)-3-(5-methoxy-pyrazin-2-yl)-urea

To a stirred solution of 1-(5-bromo-pyrazin-2-yl)-3-(2-methoxy-5-methyl-phenyl)-urea (47 mg; 0.14 mmol) in *N*-methyl pyrrolidinone (300 µL) was added sodium methoxide (0.5 mmol). The reaction was heated to 100 °C. After stirring for 12 hours, the reaction was cooled to room temperature, diluted with 30 mL of ethyl acetate and washed with aqueous 10% sodium carbonate (1 x 30 mL), brine (30 mL), then dried (MgSO₄) and filtered. The crude product was purified using a 0.5 mm prep plate eluting with hexane and ethyl acetate (1:1) to yield a yellow solid (13% yield). ¹H NMR (400 MHz,

CDCl₃) δ 8.12 (s, 1H), 7.99 (s, 1H), 7.91 (s, 1H), 6.80 (dd, 2H), 3.95 (s, 3H), 3.89 (s, 3H), 2.38 (s, 3H).
LRMS (ESI, Positive) m/e 289.10 (M+1).

Compound 326:




1-(5-Ethynyl-pyrazin-2-yl)-3-(2-methoxy-5-methyl-phenyl)-urea

Step1: 2-Amino-5-alkynylpyrazine. To a stirred solution of 5-bromo-2-aminopyrazine (432 mg; 2.5 mmol), Pd(Ph₃P)₂Cl₂ (91 mg; 0.13 mmol), CuI (1.2 g, 6.5 mmol) in triethylamine (8 mL) was added TMS-acetylene. The reaction was stirred at 60 °C for 12 hours. The reaction was cooled to room temperature, diluted with 30 mL of ethyl acetate and washed with aqueous 10% sodium carbonate (1 x 30 mL), brine (30 mL), then dried (MgSO₄), and filtered. The crude product was diluted in 1 mL of methanol and sodium hydroxide (10 mL of a 1N aqueous solution). After stirring for 12 hours the reaction was diluted with 30 mL of ethyl acetate and washed with aqueous 10% sodium carbonate (1 x 30 mL), brine (30 mL), then dried (MgSO₄), and filtered. The crude product was purified using a biotage 12L column eluting with methylene chloride and methanol (98:2) to yield an off white solid (40% yield).

Step 2: Prepared according to general procedure described for compound 310 using 2-Amino-5-alkynylpyrazine. (20% yield). ¹H NMR (400 MHz, d₆-DMSO) δ 10.25 (s, 1H), 9.80 (br s, 1H), 8.90 (s, 1H), 8.50 (s, 1H), 8.00 (s, 1H), 6.95 (d, 1H), 6.82 (d, 1H), 4.42 (s, 1H), 3.82 (s, 3H), 2.22 (s, 3H). LRMS (ESI, Positive) m/e 283.10 (M+1).

Compound 327:




1-(5-Ethyl-pyrazin-2-yl)-3-(2-methoxy-5-methyl-phenyl)-urea

Step 1: 2-Amino-5-ethylpyrazine. To a stirred solution of 5-ethynyl-2-aminopyrazine (18 mg; 0.151 mmol), in ethyl acetate (500 μ L) was added triethylamine (63 μ L; 0.45 mmol) and Pd(OH)₂ (0.01 mmol; 20% wt on carbon). The reaction was placed under a hydrogen atmosphere at 45 psi and shook for 6 hours. The reaction was filtered and concentrated under reduced pressure to yield an off white solid (84% yield).

Step 2: Prepared according to general procedure described for compound 310 using 2-Amino-5-ethylpyrazine. (27% yield). ¹H NMR (400 MHz, CDCl₃) δ 8.65 (s, 1H), 8.38 (s, 1H), 8.18 (s, 1H),

8.08 (s, 1H), 6.80 (dd, 2H), 3.92 (s, 3H), 2.81 (q, 2H), 2.39 (s, 3H), 1.39 (t, 3H). LRMS (ESI, Positive) m/e 287.21 (M+1).

Compound 328:




1-(5-Cyano-pyrazin-2-yl)-3-(2-methoxy-5-methyl-phenyl)-urea

Step 1: 2-amino-5-cyanopyrazine. To a stirred solution of 5-bromo-2-aminopyrazine (1.0 g, 5.8 mmol), CuI (2.76 g, 14.5 mmol), 18-crown-6 (121 mg; 0.46 mmol), potassium cyanide (943 mg; 14.5 mmol) in dimethylformamide (20 mL) was added Pd(PPh₃)₄ (196 mg; 0.17 mmol). After stirring at room temperature for 20 minutes the reaction was placed in an oil bath at 155 °C for 2 hours. The reaction was allowed to cool to room temperature and then poured into chloroform (300 mL). A precipitate formed that was filtered and triturated with hexanes to yield an off white solid (60% yield).

Step 2: Prepared according to general procedure described for compound 310 using 2-amino-5-cyanopyrazine. (30% yield). ¹H NMR (400 MHz, d₆-DMSO) δ 8.89 (s, 1H), 8.79 (s, 1H), 8.05 (s, 1H), 6.91 (d, 1H), 6.80 (d, 1H), 3.85 (s, 3H), 2.22 (s, 3H). LRMS (ESI, Positive) m/e 283.91 (M+1).

Compound 329:




1-(5-Benzoyl-pyrazin-2-yl)-3-(2-methoxy-5-methyl-phenyl)-urea

Step 1: 5-Benzoyl-pyrazine-2-carboxylic acid. To a stirred, cooled (0 °C) solution of 2-pyrazine carboxylic acid (3.0 g, 24.2 mmol) and benzaldehyde (7.4 mL; 73 mmol) in a 50% aqueous solution of sulfuric acid (40 mL) and 25 mL of acetic acid was added FeSO₄ · 7 H₂O (20.3 g, 73 mmol dissolved in 50 mL of water) and *t*-butyl peroxide (9.2 mL; 73 mmol) simultaneously. After stirring for 1 hour, the reaction was treated with 200 mL of water. A precipitate formed which was filtered and washed with methylene chloride (3 x 100 mL) to yield a tan solid (36% yield).

Step 2: 1-(5-Benzoyl-pyrazin-2-yl)-3-(2-methoxy-5-methyl-phenyl)-urea. To a stirred solution of 5-Benzoyl-pyrazine-2-carboxylic acid (912 mg; 4.0 mmol) and triethylamine (584 µL; 4.2 mmol) in toluene (12 mL) was added diphenyl phosphoryl azide (860 µL; 4.0 mmol). The reaction was stirred for 30 minutes followed by the addition of *t*-butanol (764 µL; 8.0 mmol). The reaction was heated to 90 °C and stirred for 3 hours. The reaction was cooled to room temperature, diluted with 30 mL of ethyl acetate and washed with aqueous 10% sodium carbonate (1 x 30 mL), brine (30 mL), then dried

(MgSO₄), and filtered. The material was purified using a biotage 40M cartridge eluting with hexane and ethyl acetate (1:1) to yield an off white solid (14% yield). ¹H NMR (400 MHz, CDCl₃) δ 9.41 (s, 1H), 9.01 (s, 1H), 8.45 (s, 1H), 8.18 (s, 1H), 8.08 (d, 2H), 7.61 (t, 1H), 7.52 (t, 2H), 6.90 (d, 1H), 6.82 (d, 1H), 3.92 (s, 3H), 2.39 (s, 3H). LRMS (ESI, Positive) m/e 363.21 (M+1).


Compound 330:



1-[5-(Hydroxy-phenyl-methyl)-pyrazin-2-yl]-3-(2-methoxy-5-methyl-phenyl)-urea

To a stirred solution of 1-(5-benzoyl-pyrazin-2-yl)-3-(2-methoxy-5-methyl-phenyl)-urea (22 mg; 0.061 mmol) in methanol (1 mL) was added sodium borohydride (10 mg; 0.3 mmol). After stirring for 12 hours, the reaction diluted with 30 mL of ethyl acetate and washed with aqueous 10% sodium carbonate (1 x 30 mL), brine (30 mL), then dried (MgSO₄), and filtered. The filtered material was concentrated under reduced pressure to yield a white solid (91% yield). ¹H NMR (400 MHz, CDCl₃) δ 8.48 (br s, 1H), 8.39 (s, 1H), 8.22 (s, 1H), 8.12 (s, 1H), 7.25-7.45 (m, 5H), 6.89 (d, 1H), 6.80 (d, 1H), 5.85 (d, 1H), 3.88 (s, 3H), 2.37 (s, 3H). LRMS (ESI, Positive) m/e 365.24 (M+1).

Compound 331:




6-Phenyl-pyrazin-2-ylamine

Suzuki procedure

Step 1: To a stirred solution of 2-amino-6-chloro pyrazine (400 mg; 3.1 mmol) and phenyl boronic acid (415 mg; 3.4 mmol) in dioxane (6 mL) and ethanol (3 mL) was added cesium carbonate (2.28 g, 7.0 mmol in 3 mL of water) followed by Pd(PPh₃)₄ (185 mg; 0.16 mmol). The reaction was heated to 75 °C and stirred for 12 hours. The reaction was cooled to room temperature, diluted with 50 mL of ethyl acetate and washed with aqueous 10% sodium carbonate (1 x 50 mL), brine (50 mL), then dried (MgSO₄), and filtered. The material was purified using a biotage 40M cartridge eluting with ethyl acetate to yield an off white solid (84% yield).

Step 2: Prepared according to general procedure described for compound 310 using 2-amino-6-phenylpyrazine. (33% yield). ¹H NMR (400 MHz, CDCl₃) δ 11.11 (br s, 1H), 8.612 (s, 1H), 8.29 (s, 1H), 8.24 (s, 1H), 8.14 (s, 1H), 8.03 (m, 2H), 7.51 (m, 3H), 6.87 (d, 1H), 6.77 (d, 1H). LRMS (ESI, Positive) m/e 355.6 (M+1).

Compound 332:




1-(3-Bromo-5-phenyl-pyrazin-2-yl)-3-(2-methoxy-5-methyl-phenyl)-urea

Step 1: 2-amino-3-bromo-5-phenylpyrazine. To a stirred solution of 3,5-dibromo-2-aminopyrazine (200 mg; 0.79 mmol) and phenyl boronic acid (106 mg; 0.87 mmol) in dioxane (4 mL) and ethanol (2 mL) was added cesium carbonate (571 mg; 1.75 mmol in 2 mL of water) followed by $\text{Pd}(\text{PPh}_3)_4$ (46 mg; 0.04 mmol). The reaction was heated to 75 °C and stirred for 12 hours. The reaction was cooled to room temperature, diluted with 50 mL of ethyl acetate and washed with aqueous 10% sodium carbonate (1 x 50 mL), brine (50 mL), then dried (MgSO_4), and filtered. The material was purified using a biotage 12L cartridge eluting with hexanes and ethyl acetate (3:1) to yield an off white solid (88% yield).

Step 2: Prepared according to general procedure described for compound 310 using 2-amino-3-bromo-5-phenylpyrazine. (18% yield). ^1H NMR (400 MHz, CDCl_3) δ 8.36 (s, 1H), 8.09 (s, 1H), 7.63 (d, 2H), 7.59 (m, 3H), 6.85 (dd, 2H), 3.92 (s, 3H), 2.39 (s, 3H). LRMS (ESI, Positive) m/e 413.2 415.2 (M+1).

Compound 333:




1-(2-Methoxy-5-methyl-phenyl)-3-(5-phenyl-pyrazin-2-yl)-urea

Step 1: 2-amino-5-phenylpyrazine. To a stirred solution of 3-bromo-5-phenyl-2-amino pyrazine (80 mg; 0.32 mmol), in ethyl acetate (1 mL) was added triethylamine (139 μL ; 1.0 mmol) and $\text{Pd}(\text{OH})_2$ (10 mg; 20% wt on carbon). The reaction was placed under a hydrogen atmosphere at 45 psi and shook for 6 hours. The reaction was filtered and concentrated under reduced pressure. The product was purified using a biotage 12L eluting with ethyl acetate to yield an off white solid (75% yield).

Step 2: Prepared according to general procedure described for compound 310 using 2-amino-5-phenylpyrazine. (25% yield). ^1H NMR (400 MHz, CDCl_3) δ 8.36 (s, 1H), 8.22 (s, 1H), 8.18 (s, 1H), 7.63 (m, 2H), 7.59 (m, 3H), 7.28 (br s, 1H), 6.82 (m, 2H), 3.92 (s, 3H), 2.33 (s, 3H). LRMS (ESI, Positive) m/e 335.21 (M+1).

Compound 334:




Quinoxalin-2-ylamine

Step 1: To 2-chloroquinoxaline (1.0 g, 6 mmol) was added ammonia in methanol (8 mL of a 2M solution). The reaction was sealed in a vial, heated to 80 °C, and stirred for 12 hours. The reaction was cooled to room temperature and concentrated under reduced pressure. The residue was taken up in methylene chloride and filtered. Hexane was added until a precipitate formed which was filtered and found to be the desired product (5% yield).

Step 2: Prepared according to general procedure described for compound 310 using Quinoxalin-2-ylamine. (26% yield). ¹H NMR (400 MHz, D₆-DMSO) δ 11.63 (br s, 1H), 10.59 (br s, 1H), 8.80 (s, 1H), 8.15 (s, 1H), 7.97 (d, 1H), 7.81 (m, 2H), 7.64 (m, 1H), 6.98 (d, 1H), 6.82 (d, 1H), 3.97 (s, 3H), 2.23 (s, 3H). LRMS (ESI, Positive) m/e 309.4 (M+1).

Compound 335:




1-(3,6-Dimethyl-pyrazin-2-yl)-3-(2-methoxy-5-methyl-phenyl)-urea

Step 1: **2-Azido-3,6-dimethylpyrazine.** To a stirred solution of 2-chloro-3,5-dimethyl pyrazine (1.0 mL; 8.3 mmol) in dimethylformamide (10 mL) was added sodium azide (539 mg; 8.3 mmol). The reaction was heated to 100 °C and stirred for 12 hours. The reaction was cooled to room temperature, diluted with 50 mL of ethyl acetate and washed with aqueous 10% sodium carbonate (1 x 50 mL), brine (50 mL), then dried (MgSO₄), and filtered. The material was purified using a biotage 12L cartridge eluting with hexanes and ethyl acetate (3:1) to yield an off white solid (42% yield).

Step 2: **2-Amino-3,6-dimethylpyrazine.** To a stirred solution of 3-azido-2,5-dimethyl pyrazine (100 mg; 0.66 mmol) in methanol (800 µL) was added 12N HCl (100 µL) and tin chloride dihydrate (149 mg; 0.66 mmol). The reaction was heated to 60 °C and stirred for 12 hours. The reaction was cooled to room temperature, diluted with 50 mL of ethyl acetate and washed with aqueous 10% sodium carbonate (1 x 50 mL), brine (50 mL), then dried (MgSO₄), and filtered. The material was purified using a biotage 12i cartridge eluting with ethyl acetate to yield an off white solid (38% yield).

Step 3: Prepared according to general procedure described for compound 310 using 2-Amino-3,6-dimethylpyrazine. (15% yield). ¹H NMR (400 MHz, CDCl₃) δ 8.22 (br s, 1H), 8.01 (s, 1H), 6.82 (m, 2H), 3.92 (s, 3H), 2.59 (s, 3H), 2.55 (s, 3H), 2.38 (s, 3H). LRMS (ESI, Positive) m/e 287.20 (M+1).

Compound 336:




Step 1: 2-Methoxy-4-methoxymethyl-1-nitro-benzene To a 250 mL round bottomed flask containing 5.4 g (39 mmol) of 3-methoxy-4-nitrobenzyl alcohol in 30 mL THF and 30 mL DMF was added, 38 g (117 mmol, 3 eq.) of finely powdered cesium carbonate followed by 24 mL (390 mmol, 10 eq.) iodomethane. The mixture was stirred at room temperature for 18 h, and was then partitioned between 100 mL water and 100 mL diethyl ether. The aqueous phase was extracted with ether (2 x 100 mL), and the combined organic extracts were washed with brine (2 x 50 mL), dried over MgSO₄, filtered through a short plug of silica, and concentrated. The residue was purified by flash chromatography, eluting with 1:1 EtOAc-Hexane, to give 6.76 g (88%) of the methyl ether as a yellow oil. ¹H NMR (400 MHz, CDCl₃): δ 7.84, (d, J=8.2 Hz, 1H) 7.10 (s, 1H), 6.94 (d, J=9.1 Hz, 1H), 4.51 (s, 2H), 3.98 (s, 3H), 3.44 (s, 3H).

Step 2: 2-Methoxy-4-methoxymethyl-phenylamine. In a 250 mL Parr apparatus, 2.1 g (10.6 mmol) of 2-methoxy-4-methoxymethyl-1-nitro-benzene, in 40 mL of ethanol was hydrogenated at 2 atm over 300 mg of 10% Pd on carbon for 2.5 h. The catalyst was removed by filtration through a glass fiber filter and the filtrate was concentrated to give 1.61 g (91%) of product as a light yellow oil. ¹H NMR (400 MHz, CDCl₃): δ 6.80 (s, 1H), 6.74 (d, J=7.8 Hz, 1H), 6.67 (d, J=7.8 Hz), 4.35 (s, 2H), 3.86 (s, 3H), 3.78 (br d, 2H), 3.35 (s, 3H). MS ESI-pos, M+1 = 168.1

Step 3: 1-(2-Methoxy-4-methoxymethyl-phenyl)-3-(5-methylpyrazin-2-yl)-urea.

General diphenylphosphoryl azide coupling method: To a solution of 5-methylpyrazine-2-carboxylic acid (365 mg, 2.64 mmol) in 20 mL of anhydrous toluene, was added, diisopropylethylamine (483 μL, 2.77 mmol) and the mixture was stirred at room temperature until the solid dissolved. Then diphenylphosphoryl azide was added and the solution was heated to 90 °C. After 20 minutes N₂ evolution had subsided, and the caramel colored reaction mixture was cooled to 60 °C before adding 2-methoxy-4-methoxymethylaniline as a solution in 4 mL toluene. After stirring for 6 hr at 60 °C, the mixture was cooled to room temperature and diluted with 20 mL of 5% NH₄OH, and extracted with EtOAc (3 x 50 mL). The combined extracts were washed with 20 mL water and 20 mL brine then dried over MgSO₄, filtered and concentrated. The brown residue was purified by flash chromatography (eluting with 5% MeOH in CH₂Cl₂) to give 219 mg (27%) of the desired product. ¹H NMR (400 MHz, CDCl₃): δ 11.36 (s, 1H), 9.47, (s, 1H) 8.40, (s, 1H), 8.31 (s, 1H), 8.08 (s, 1H), 6.95 (s, 1H), 6.94 (d, J= 7.8 Hz, 1H), 4.45 (s, 2H), 3.97 (s, 3H), 3.39 (s, 3H), 2.52 (s, 3H). MS ESI-pos M+1=303.2

Compound 337:




1-(4-Benzyloxymethyl-2-methoxy-phenyl)-3-(5-methyl-pyrazin-2-yl)-urea

Step 1: 4-Benzyloxymethyl-2-methoxy-1-nitro-benzene. To a stirred suspension of finely powdered cesium carbonate (8.0 g, 24.5 mmol) was added 3-methoxy-4-nitrobenzyl alcohol (1.5 g, 8.18 mmol), followed by benzyl bromide, (2 mL, 16.4 mmol). After stirring at room temperature for 18 h, the suspension diluted with 100 mL diethyl ether, and washed with 3 x 50 mL water, then 50 mL brine. The organic phase was dried over MgSO_4 , filtered through a plug of silica, and concentrated. The resulting orange oil was purified by flash chromatography, eluting with 2:1 hexane-EtOAc, to give 1.87 g, (84%) of the benzyl ether. ^1H NMR (400 MHz, CDCl_3): δ 7.85 (d, J = 8.2 Hz, 1H), 7.3-7.4 (m, 5H), 7.26 (s, 1H), 6.97 (d, J = 8.2 Hz, 1H), 4.61 (s, 2H), 4.59 (s, 2H), 3.96 (s, 3H).

Step 2: 4-Benzyloxymethyl-2-methoxy-phenylamine. A solution of 4-nitro-3-methoxybenzylbenzyl ether (2.2 g, 8.1 mmol) and ammonium acetate (2.46 g, 32 mmol, 4 eq.) in 30 mL MeOH was stirred at 0 °C and 1.3 g (20 mmol, 2.5 eq.) of zinc dust was added in several portions. After 1 h, the reaction mixture was partitioned between 40 mL water and 40 mL ethyl acetate. The organic phase was dried over MgSO_4 and concentrated. The residue was used directly in next step without further purification. ^1H NMR (400 MHz, CDCl_3): δ 7.35 (m, 5H), 6.82 (s, 1H), 6.77 (d, J = 6.3 Hz, 1H), 6.67 (d, J = 7.8 Hz, 1H), 4.51 (s, 2H), 4.45 (s, 2H), 3.85 (s, 3H), 3.79 (s, 2H). MS ESI-pos, $M+1$ = 244.2.

Step 3: 1-(4-Benzyloxymethyl-2-methoxy-phenyl)-3-(5-methyl-pyrazin-2-yl)-urea. Prepared according to the general diphenylphosphoryl azide coupling method described above for compound 336. ^1H NMR (400 MHz, CDCl_3): δ 11.36, (s, 1H), 9.23 (s, 1H), 8.39 (s, 1H), 8.32 (d, J = 7.8 Hz, 1H), 8.09 (s, 1H), 7.38 (m, 5H), 6.97, (s, 2H), 4.56 (s, 4H), 3.96 (s, 3H), 2.53 (s, 3H). MS APCI-pos, $M+1$ = 379.3

Compound 338:



1-{4-[(Benzy-methyl-amino)-methyl]-2-methoxy-phenyl}-3-(5-methyl-pyrazin-2-yl)-urea

Step 1: 3-Methoxy-4-nitro-benzyl bromide. To a 250 mL round bottomed flask containing 10 g (54.6 mmol) of 4-nitro-3-methoxybenzyl alcohol in 30 mL THF was added, 36 g (109 mmol, 2 eq.) of carbon tetrabromide followed by 15.9 g (60 mmol, 1.1 eq.) triphenylphosphine at 0 °C. The mixture was stirred at 0 °C for 3 hours. Upon removal of the solvent, the residue was purified by flash chromatography, eluting with 10:90 EtOAc-Hexane, to give 11 g (82%) of the product as a yellow solid. ¹H NMR (400 MHz, CDCl₃): δ 7.86 (s, 1H), 7.84 (m, 1H), 7.12 (s, 1H), 7.07 (m, 1H), 4.47 (s, 2H), 4.00 (s, 3H).


Step 2: *N*-Benzyl-*N*-(3-methoxy-4-nitro-benzyl)-amine. To a 150 mL round bottomed flask containing 1.97 g (8.0 mmol) of 3-methoxy-4-nitro-benzyl bromide in 20 mL THF was added, 2.4 g (24 mmol, 3eq.) of triethylamine followed by 2.5 g (24 mmol, 3 eq.) of benzylamine. The mixture was stirred at room temperature for 2 h, and was then partitioned between 50 mL ethyl acetate and brine. The organic phase was dried over MgSO₄ and concentrated. The residue was purified by flash chromatography, eluting with 1-4% MeOH in dichloromethane to give 1.6 g (73%) of the benzyl amine as a yellow oil. ¹H NMR (400 MHz, CDCl₃): δ 7.84 (d, *J* = 8.61 Hz, 1H), 7.34 (m, 5H), 7.16 (s, 1H), 6.99 (d, *J* = 8.61 Hz, 1H), 3.97 (s, 3H), 3.86 (s, 2H), 3.81 (s, 2H).

Step 3: Benzyl-(3-methoxy-4-nitro-benzyl)-carbamic acid tert-butyl ester. To a 150 mL round bottomed flask containing 0.92 g (3.4 mmol, 1 eq.) of *N*-benzyl-*N*-(3-methoxy-4-nitro-benzyl)-amine in 2 mL dichloromethane was added Boc anhydride (0.74 g, 3.4 mmol, 1 eq.) then stirred at room temperature for 18 h. The reaction mixture was then partitioned between 40 mL water and 40 mL ethyl acetate. The organic phase was dried over MgSO₄ and concentrated. No further purification was necessary. ¹H NMR (400 MHz, CDCl₃): δ ¹H NMR (400 MHz, CDCl₃): δ 7.81 (d, *J* = 8.61 Hz, 1H), 7.2-7.3 (m, 6H), 6.93 (m, 1H), 6.82 (s, 1H), 4.39 (m, 4H), 3.88 (s, 3H), 1.53 (s, 9H).

Steps 4-6: 1-[4-[(Benzyl-methyl-amino)-methyl]-2-methoxy-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea. Benzyl-(3-methoxy-4-nitro-benzyl)-carbamic acid tert-butyl ester was reduced to the corresponding aniline according to the general hydrogenation procedure detailed above for compound 336. The crude aniline was used in the coupling step as follows: A solution of 5-methylpyrazine-2-carboxylic acid (34.5 mg, 0.25 mmol) and added triethylamine (28 mg, 0.275 mmol) in 5 mL of anhydrous toluene was stirred at room temperature until the solid dissolved. Diphenylphosphoryl azide (62 mg, 0.225 mmol) was added and the solution heated to 90 °C for 20 min. The reaction pot was then transferred to a 60 °C oil bath, and the aniline (0.25 mmol) was added as a solution in 2 mL toluene. After stirring for 4.5 hr at 60 °C, the mixture was cooled to room temperature, diluted with EtOAc, washed with sat'd NaHCO₃, then brine. The organic phase was dried over MgSO₄, filtered and concentrated. The resulting residue was purified by preparative TLC, eluting with 5% MeOH in CH₂Cl₂, to give the desired urea. The Boc group was removed by treatment of the Boc protected amine in 15 mL of CH₂Cl₂ with 3 mL TFA and stirring at room temperature for 3h. The mixture was diluted with EtOAc (50 mL), washed with 20 mL of sat'd NaHCO₃ followed by 20 mL of brine. The organic

phase was then dried over MgSO_4 , filtered and concentrated to give the free amine ^1H NMR (400 MHz, d_6 -DMSO): δ 11.35 (s, 1H), 9.68 (s, 1H), 8.41 (s, 1H), 8.28 (d, J = 8.21 Hz, 1H), 8.06 (s, 1H), 7.36 (s, 6H), 6.96 (s, 1H), 6.94 (d, J = 8.21 Hz, 1H), 3.95 (s, 3H), 3.84 (s, 2H), 3.81 (s, 2H), 2.52 (s, 3H), 2.13 (s, 1H). MS APCI-pos, $M+1$ = 377.9

Compound 339:




1-(2-Methoxy-4-methylaminomethyl-phenyl)-3-(5-methyl-pyrazin-2-yl)-urea

Step 1+2: (3-Methoxy-4-nitro-benzyl)-methyl-carbamic acid tert-butyl ester. In a fashion similar to that described above for the analogous benzyl derivative compound 338, 3-methoxy-4-nitro-benzyl bromide was alkylated with methylamine, and the resulting secondary amine was protected as the Boc derivative. ^1H NMR (400 MHz, CDCl_3): δ 7.84 (d, J = 8.61 Hz, 1H), 6.98 (s, 1H), 6.87 (d, J = 8.61 Hz, 1H), 4.46 (s, 2H), 3.95 (s, 7H), 2.85 (s, 3H), 1.53 (s, 9H).

Step 3: (4-Amino-3-methoxy-benzyl)-methyl-carbamic acid tert-butyl ester. In a 250 mL Parr apparatus, 0.98 g (3.5 mmol) of (3-methoxy-4-nitro-benzyl)-methyl-carbamic acid tert-butyl ester in 40 mL of ethanol was hydrogenated at 2 atm over 300 mg of 10% Pd on carbon for 15 minutes. The catalyst was removed by filtration through a glass fiber filter and the filtrate was concentrated to give a crude product as light yellow oil. ^1H NMR (400 MHz, CDCl_3): δ 6.69 (m, 3H), 3.95 (s, 2H), 3.84 (s, 3H), 2.8 (m, 2H), 2.75 (s, 3H), 1.51 (s, 9H).

Step 4+5: 1-(2-Methoxy-4-methylaminomethyl-phenyl)-3-(5-methyl-pyrazin-2-yl)-urea. A solution of (4-amino-3-methoxy-benzyl)-methyl-carbamic acid tert-butyl ester was converted to the urea according to the general diphenylphosphoryl azide coupling method detailed for compound 336. The Boc group was removed as described for above for compound 338. ^1H NMR (400 MHz, d_6 -DMSO): δ 9.91 (s, 2H), 8.78 (s, 1H), 8.21 (s, 1H), 8.07 (d, J = 8.61 Hz, 1H), 7.01 (s, 1H), 6.85 (d, J = 8.61 Hz, 1H), 3.89 (s, 3H), 3.59 (s, 1H), 2.42 (s, 3H), 2.26 (s, 3H). MS APCI-Pos, $M+1$ = 301.8.

Compound 340:



1-{4-[(Benzyl-methyl-amino)-methyl]-2-methoxy-phenyl}-3-(5-methyl-pyrazin-2-yl)-urea

Step 1: N-Benzyl-N-(3-methoxy-4-nitro-benzyl)-methyl-amine. A solution of N-methyl-benzyl amine was alkylated with 3-methoxy-4-nitro-benzyl bromide as described above for compound 338.

^1H NMR (400 MHz, CDCl_3): δ 7.83 (d, J = 8.6 Hz, 1H), 7.35 (s, 4H), 7.26 (s, 1H), 7.17 (s, 1H), 7.01 (d, J = 7.04 Hz, 1H), 3.97 (s, 3H), 3.55 (s, 4H), 2.22 (s, 3H).


Step 2: 4-[(Benzyl-methyl-amino)-methyl]-2-methoxy-phenylamine.

Nickel boride Reduction General Method: To a stirred solution of nickel chloride hexahydrate (820 mg, 3.45 mmol) in 12 mL EtOH and 3 mL THF, at 0 °C, NaBH₄ (130 mg, 3.45 mmol) was added. The resulting black suspension was stirred at 0 °C while *N*-benzyl-(3-methoxy-4-nitro-benzyl)-methylamine was added as a solution in 5 mL THF. After several minutes, 260 mg of NaBH₄ was added in several portions over 10 minutes, and the reaction mixture was subsequently allowed to warm to room temperature. After 2 h, TLC indicated complete conversion to a new, more polar product. At this point, 1.5 mL of 5% NH₄OH was added and the reaction was stirred for about 10 min, until the black solids achieved a granular consistency. The reaction was filtered through a glass fiber filter, rinsing with THF. The clear colorless filtrate was concentrated to about ¼ volume, diluted with 30 mL water and extracted with EtOAc (3 x 30 mL). The combined extracts were washed with brine, dried over MgSO₄ and filtered through a short plug of silica and concentrated *in vacuo* to afford 575 mg (65%) of the desired product as an off-white solid. ¹H NMR (400 MHz, CDCl₃): δ 7.2-7.4 (m, 5H), 7.85 (s, 1H), 6.85 (s, 1H), 6.74 (d, *J* = 8.2 Hz, 1H), 6.66 (d, *J* = 8.2 Hz, 1H), 3.87 (s, 3H), 3.73 (br. s, 2H) 3.49 (s, 2H), 3.45 (s, 2H), 2.18 (s, 3H). MS (APCI-pos)M+1= 256.9

Step 3: 1-{4-[(Benzyl-methyl-amino)-methyl]-2-methoxy-phenyl}-3-(5-methyl-pyrazin-2-yl)-urea.

In a 50 mL round bottom flask, 5-methylpyrazine-2-carboxylic acid (250 mg, 1.8 mmol) and diisopropylethylamine (330 µL, 1.9 mmol) in 20 mL toluene, was stirred under a nitrogen atmosphere until the acid dissolved. Diphenylphosphoryl azide (523 mg, 1.9 mmol) was added and the solution was heated to 90 °C. After 20 minutes, nitrogen evolution had subsided, and the solution had darkened to a caramel color. The reaction was cooled to 65 °C, and 4-[(benzyl-methyl-amino)-methyl]-2-methoxy-phenylamine (486 mg, 1.9 mmol) was added as a solution in 5 mL toluene. The reaction was allowed to stir at 65 °C for 6 h, and was then cooled to room temperature, diluted with 30 mL EtOAc, and washed with 15 mL 5% NH₄OH. The aqueous phase was extracted with EtOAc (2 x 20 mL) and the combined organics were washed with 30 mL brine, dried over MgSO₄, filtered, and concentrated. The resulting residue was purified by flash chromatography to give 285 mg (40%) of the desired product, which was further purified by trituration with diethyl ether. Mp=142-143 °C. ¹H NMR (400 MHz, CDCl₃): δ 11.33 (s, 1H), 9.51 (s, 1H), 8.41 (s, 1H), 8.27 (d, *J* = 8.61 Hz, 1H), 8.08 (s, 1H), 7.2-7.4 (m, 5H), 6.96 (d, *J* = 7.83 Hz, 1H) 3.97 (s, 3H), 3.53 (s, 4H), 2.53 (s, 3H), 2.22 (s, 3H). ¹³C NMR (400 MHz, CDCl₃): δ 153.70, 149.03, 147.48, 147.4, 145.99, 138.73, 137.38, 134.81, 129.19, 128.42, 127.16, 121.89, 119.81, 111.05, 104.49, 94.98, 87.22, 61.81, 56.37, 42.5. MS (APCI-pos)M+1=392.0.

Compound 341:




1-(4-Dimethylaminomethyl-2-methoxy-phenyl)-3-(5-methyl-pyrazin-2-yl)-urea

Step 1: (3-Methoxy-4-nitro-benzyl)-dimethyl-amine According to the method described above for compound 338, 3-methoxy-4-nitro-benzyl bromide was treated with dimethyl amine to give the desired product. ¹H NMR (400 MHz, CDCl₃): δ 7.86 (d, *J* = 7.8 Hz, 1H), 7.3 (s, 1H), 6.98 (d, *J* = 8.6 Hz, 1H), 4.01 (s, 3H), 3.5 (s, 2H), 2.3 (s, 6H).

Step 2: 4-Dimethylaminomethyl-2-methoxy-phenylamine According to the nickel-boride method described for compound 340 above, (3-methoxy-4-nitro-benzyl)-dimethyl-amine, was reduced to the corresponding aniline and was used in the next step without characterization.

Step 3: 1-(4-Dimethylaminomethyl-2-methoxy-phenyl)-3-(5-methyl-pyrazin-2-yl)-urea. According to the diphenylphosphoryl azide method described above for compound 336, 4-dimethylaminomethyl-2-methoxy-phenylamine was converted to the (5-methyl-pyrazin-2-yl)-urea. The crude product was purified by preparative TLC, eluting with 5% MeOH in CH₂Cl₂. ¹H NMR (400 MHz, *d*₆-DMSO): δ 11.21 (s, 1H), 8.95 (s, 1H), 8.39 (s, 1H), 8.21 (d, *J* = 7.83 Hz, 1H), 8.04 (s, 1H), 7.01 (s, 1H), 6.89 (d, *J* = 7.83 Hz, 1H), 3.98 (s, 3H), 3.92 (s, 2H), 2.26 (s, 3H), 2.24 (s, 6H). MS APCI-Pos, *M*+1 = 316.0.

Compound 342:



1-(4-Aminomethyl-2-methoxy-phenyl)-3-(5-methyl-pyrazin-2-yl)-urea

Step 1: 2-(3-Methoxy-4-nitro-benzyl)-isoindole-1,3-dione. To a 250 mL round bottomed flask containing 10 g (55 mmol) of 4-nitro-3-methoxybenzyl alcohol in 150 mL THF was added, diethylazodicarboxylate (8.03 g, 54.6 mmol, 1 eq.) and triphenylphosphine (15.0 g, 57.3 mmol) followed by 11.6 g of phthalimide (57.3 mmol) at 0 °C. Reaction was then allowed to gradually warm to room temperature over night. A white precipitate formed and was collected by suction filtration. Recrystallization from acetonitrile gave 13.8 g (81%) of the desired product. ¹H NMR (400 MHz, *d*₆-DMSO): δ 7.79 (m, 5H), 7.19 (s, 1H), 7.08 (d, *J* = 8.80 Hz, 1H), 4.91 (s, 2H), 3.87 (s, 3H).

Step 2: 2-(4-Amino-3-methoxy-benzyl)-isoindole-1,3-dione. In a 500 mL Parr vessel, a partially dissolved suspension of 2-(3-methoxy-4-nitro-benzyl)-isoindole-1,3-dione (1.50 g, 4.80 mmol) in 100 mL EtOH and 30 mL THF was hydrogenated over 250 mg 10 % Pd-C at 2.5 atm for 1 h. The catalyst was removed by filtration through a glass fiber filter, and the clear light yellow filtrate was

concentrated *in vacuo*. The product was washed with 30 mL of diethyl ether and collected by suction filtration to give 1.28 g (95%) of the aniline as fine, light green needles. ¹H NMR (400 MHz, CDCl₃): δ 7.82 (m, 2H), 7.81 (m, 2H), 6.93 (s, 1H), 6.9 (d, *J* = 7.8 Hz, 1H), 6.63 (d, *J* = 7.8 Hz, 1H), 4.74 (s, 2H), 3.84 (s, 3H).

Step 3: Acyl Azide coupling General Method


1-[4-(1,3-Dioxo-1,3-dihydro-isoindol-2-ylmethyl)-2-methoxy-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea. In a 50 mL round bottom flask, a solution of 5-methyl-pyrazine-2-carbonyl azide (510 mg, 3.15 mmol) in 15 mL of anhydrous toluene was stirred under a nitrogen atmosphere. The reaction flask was immersed in a 90 °C oil bath, and as the internal temperature approached 90 °C, N₂ release was evident and the solution started to darken. After 20 min, effervescence had subsided, and the solution had darkened to a caramel color. The reaction flask was moved to a 65 °C bath, and 2-(4-amino-3-methoxy-benzyl)-isoindole-1,3-dione (884 mg, 3.15 mmol) suspended in toluene (5 mL) was added. The reaction was stirred at 65 °C for 6 h, then cooled to room temperature. The product precipitated from solution after cooling to room temperature, and was collected by suction filtration.

In cases where the product does not precipitate from the reaction mixture, the following work-up is applied: After cooling to room temperature, the brown solution is diluted with 5% aq. NH₄OH and the extracted with EtOAc (3x). The combined extracts are washed with brine, dried over MgSO₄, filtered and concentrated. The residue is then purified by flash chromatography in an appropriate solvent system.

Step 4: 1-(4-Aminomethyl-2-methoxy-phenyl)-3-(5-methyl-pyrazin-2-yl)-urea In a 100 mL round bottom flask, a suspension of 1-[4-(1,3-dioxo-1,3-dihydro-isoindol-2-ylmethyl)-2-methoxy-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea (620 mg, 1.48 mmol) in 22 mL of EtOH, under a nitrogen atmosphere, was warmed, with stirring, to 70 °C. Hydrazine monohydrate (1.4 mL) was added, and the reaction was stirred at 70 °C. After 10 minutes, the reaction had become a completely homogenous caramel colored solution. After several minutes more, product started to precipitate from the solution. After 20 min, the reaction was cooled to room temperature, and the white solid product was collected by suction filtration. The crude product, which contained some phthalhydrazide by-product, was taken up in 80 mL of EtOAc, and washed with water (3 x 20 mL). The washings were back extracted with 30 mL EtOAc, and the combined organics were washed with brine, then dried over MgSO₄, filtered, and concentrated *in vacuo* to afford 400 mg, (94%) of the desired amine. Mp=168-169 °C. ¹H NMR (400 MHz, *d*₆-DMSO): δ 9.88 (br.s, 2H), 8.78 (s, 1H), 8.21 (s, 1H), 8.05 (d, *J* = 8.2 Hz, 1H), 7.05 (s, 1H), 6.85 (d, *J* = 8.1 Hz, 1H), 3.89 (s, 3H), 3.68 (s, 2H), 2.42 (s, 3H) 1.80 (br.s, 2H, NH₂). MS (APCI-pos)M-17 (-NH₃) = 270.1 apci-neg M-1=285.8.

Alkyl Derivatives of Compound 342.

Compound 343:




A solution of 1-(4-aminomethyl-2-methoxy-phenyl)-3-(5-methyl-pyrazin-2-yl)-urea (0.25 mmol, 1.0 eq.) and 1 mL trimethylorthoformate in 4 mL MeOH was stirred at room temperature, then thiophene-2-carboxaldehyde (2.5 mmol, 10 eq.) was added and the mixture was heated at 80 °C. After 18 h the reaction was cooled to room temperature and was concentrated *in vacuo*. The resulting imine was taken up in 5 mL of anhydrous MeOH and stirred at 0 °C. Sodium borohydride (0.75 mmol, 3 eq.) was added, and the reaction was stirred at 0 °C for 30 minutes and was then diluted with 2 mL of water and partitioned between 50 mL EtOAc and 30 mL sat. NaHCO₃. The organic phase was washed with brine, dried over MgSO₄, filtered and concentrated. If needed, the product was purified by flash chromatography, eluting with an appropriate MeOH-CH₂Cl₂ mixture.

1-(2-Methoxy-4-(((thiophen-2-ylmethyl)-amino)-methyl)-phenyl)-3-(5-methyl-pyrazin-2-yl)-urea

¹H NMR (400 MHz, *d*₆-DMSO): δ 10.01 (s, 2H), 8.79 (s, 1H), 8.23 (s, 1H), 8.17 (d, *J* = 8.61 Hz, 1H), 7.59 (d, *J* = 6.26 Hz, 1H), 7.33 (s, 1H), 7.15 (m, 2H), 7.05 (s, 1H), 6.97 (d, *J* = 8.61 Hz, 1H), 4.02 (s, 2H), 3.91 (s, 3H), 3.88 (s, 1H), 3.78 (s, 1H), 2.43 (s, 3H). MS APCI-Pos no detectable molecular ion.

Compound 344:




Prepared according to the general procedure described for compound 343 using thiophene-3-carboxaldehyde.

1-(2-Methoxy-4-(((thiophen-3-ylmethyl)-amino)-methyl)-phenyl)-3-(5-methyl-pyrazin-2-yl)-urea

¹H NMR (400 MHz, *d*₆-DMSO): δ 9.92 (s, 2H), 8.78 (s, 1H), 8.21 (s, 1H), 8.07 (d, *J* = 7.81 Hz, 1H), 7.47 (s, 1H), 7.3 (s, 1H), 7.11 (d, *J* = 4.88 Hz, 1H), 7.04 (s, 1H), 6.87 (d, *J* = 7.81 Hz, 1H), 3.9 (s, 3H), 3.67 (s, 2H), 3.65 (s, 2H), 2.42 (s, 3H). MS APCI-Neg, *M*-1 = 382.0.


Compound 345:



Prepared according to the general procedure described for compound 343 using furfural.


1-(4-(((Furan-2-ylmethyl)-amino)-methyl)-2-methoxy-phenyl)-3-(5-methyl-pyrazin-2-yl)-urea

¹H NMR (400 MHz, MeOD): δ 8.57 (s, 1H), 8.2 (s, 1H), 8.1 (d, *J* = 8.61 Hz, 1H), 7.46 (s, 1H), 7.03 (s, 1H), 6.88 (d, *J* = 8.61 Hz, 1H), 6.36 (s, 1H), 6.27 (s, 1H), 3.96 (s, 3H), 3.74 (s, 2H), 3.71 (s, 2H), 2.48 (s, 3H). MS APCI-Neg, *M*-1 = 366.0.

Compound 346:


Prepared according to the general procedure described for compound 343 using furan-3-carboxaldehyde.

1-(4-((Furan-3-ylmethyl)-amino)-methyl)-2-methoxy-phenyl)-3-(5-methyl-pyrazin-2-yl)-urea ^1H NMR (400 MHz, MeOD): δ 8.58 (s, 1H), 8.21 (s, 1H), 8.13 (d, $J = 7.83$ Hz, 1H), 7.49 (s, 1H), 7.05 (s, 1H), 6.91 (d, $J = 7.04$ Hz, 1H), 6.5 (s, 1H), 3.97 (s, 3H), 3.78 (s, 2H), 3.69 (s, 2H), 2.49 (s, 3H). MS APCI-Neg, $M-1 = 366.0$

Compound 347:

Prepared according to the general procedure described for compound 343 using 2-methoxybenzaldehyde.

1-(2-Methoxy-4-((2-methoxy-benzylamino)-methyl)-phenyl)-3-(5-methyl-pyrazin-2-yl)-urea ^1H NMR (400 MHz, CDCl_3): δ 11.3 (s, 1H), 9.97 (s, 1H), 8.44 (s, 1H), 8.22 (d, $J = 7.83$ Hz, 1H), 7.98 (s, 1H), 7.28 (m, 2H), 7.06 (s, 1H), 6.88 (m, 3H), 4.08 (s, 1H), 3.93 (s, 5H), 3.81 (s, 5H), 2.5 (s, 3H). MS APCI-Pos, $M+1 = 407.8$.

Compound 348:


Prepared according to the general procedure described for compound 343 using 3-methoxybenzaldehyde.

1-(2-Methoxy-4-((3-methoxy-benzylamino)-methyl)-phenyl)-3-(5-methyl-pyrazin-2-yl)-urea ^1H NMR (400 MHz, d_6 -DMSO): δ 8.57 (s, 1H), 8.20 (s, 1H), 8.11 (d, $J = 8.61$ Hz, 1H), 7.22 (t, $J = 7.83$ Hz, 1H), 7.04 (s, 1H), 6.91 (m, 3H), 6.83 (d, $J = 8.61$ Hz, 1H), 3.96 (s, 3H), 3.79 (s, 3H), 3.75 (s, 4H), 2.48 (s, 3H). MS APCI-Neg, $M-1 = 406.0$

COc1ccc(cc1)CNCCc2ccc(NC(=O)Nc3cc(C)nc(C)n3)c(OC)c2

1-{2-Methoxy-4-[(4-methoxy-benzylamino)-methyl]-phenyl}-3-(5-methyl-pyrazin-2-yl)-urea ¹H
NMR (400 MHz, *d*6-DMSO): δ 9.91 (s, 2H), 8.78 (s, 1H), 8.21 (s, 1H), 8.07 (d, *J* = 7.81 Hz, 1H), 7.25
(d, *J* = 8.78 Hz, 2H), 7.03 (s, 1H), 6.88 (m, 2H), 3.89 (s, 3H), 3.73 (s, 3H), 3.62 (s, 2H), 3.6 (s, 2H),
2.42 (s, 3H). MS APCI-Pos, M+1 =407.9.

Compound 350:




N-{3-Methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzyl}-acetamide ¹H NMR (400 MHz, d6-DMSO): δ 9.92 (s, 2H), 8.75 (s, 1H), 8.28 (t, *J* = 5.87 Hz, 1H), 8.19 (s, 1H), 8.07 (d, *J* = 8.61 Hz, 1H), 6.92 (s, 1H), 6.78 (d, *J* = 8.61 Hz, 1H), 4.19 (s, 1H), 4.18 (s, 1H), 3.87 (s, 3H), 2.4 (s, 3H), 1.85 (s, 3H). MS ESI-pos, M+1 = 330.2.

COCC(=O)NCc1ccc(NC(=O)Nc2cc(C)ncn2)c(OC)c1

Prepared according to the general procedure described for compound 350 using methoxyacetyl chloride.

2-Methoxy-N-{3-methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzyl}-acetamide ^1H NMR (400 MHz, d_6 -DMSO): δ 9.51 (s, 2H), 8.34 (s, 1H), 7.89 (t, J = 6.26 Hz, 1H), 7.78 (s, 1H), 7.65 (d, J = 7.83 Hz, 1H), 6.54 (s, 1H), 6.38 (d, J = 7.83 Hz, 1H), 4.2 (s, 1H), 4.19 (s, 1H), 3.82 (s, 3H), 3.79 (s, 2H), 3.29 (s, 3H), 2.35 (s, 3H). MS APCI-Pos, $M+1$ = 359.9.


Compound 352:



Prepared according to the general procedure described for compound 350 using dimethylamino-acetyl chloride.

2-Dimethylamino-N-{3-methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzyl}-acetamide ^1H NMR (400 MHz, MeOD): δ 8.56 (s, 1H), 8.18 (s, 1H), 8.09 (d, J = 7.83 Hz, 1H), 6.97 (s, 1H), 6.86 (d, J = 7.83 Hz, 1H), 4.37 (s, 2H), 3.94 (s, 3H), 3.03 (s, 2H), 2.47 (s, 3H), 2.3 (s, 6H). MS APCI-Neg, $M-1$ = 370.9.


Compound 353:



Prepared according to the general procedure described for compound 350 using 2-(2-pyridyl)-acetyl chloride.

N-{3-Methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzyl}-2-pyridin-2-yl-acetamide ^1H NMR (400 MHz, d_6 -DMSO): δ 9.94 (s, 2H), 8.77 (s, 1H), 8.63 (t, J = 5.87 Hz, 1H), 8.49 (s, 1H), 8.44 (d, J = 6.26 Hz, 1H), 8.22 (s, 1H), 8.08 (d, J = 7.83 Hz, 1H), 7.7 (d, J = 10.17 Hz, 1H), 7.35 (q, J = 5.74 Hz, 1H), 6.87 (s, 1H), 6.79 (d, J = 9.39 Hz, 1H), 4.25 (s, 1H), 4.24 (s, 1H), 3.82 (s, 3H), 3.53 (s, 2H), 2.42 (s, 3H). MS APCI-Pos $M+1$ = 407.1.


Compound 354:



Prepared according to the general procedure described for compound 350 using 2-(4-methoxyphenyl)-acetyl chloride.

***N*-{3-Methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzyl}-2-(4-methoxy-phenyl)-acetamide** ¹H NMR (400 MHz, *d*₆-DMSO): δ 9.94 (s, 2H), 8.77 (s, 1H), 8.54 (t, *J* = 4.7 Hz, 1H), 8.21 (s, 1H), 8.08 (d, *J* = 7.83 Hz, 2H), 7.89 (m, 1H), 7.22 (t, *J* = 7.83 Hz, 1H), 6.89 (s, 3H), 4.25 (s, 1H), 4.23 (s, 1H), 3.8 (s, 3H), 3.73 (s, 2H), 3.45 (s, 2H), 2.42 (s, 3H). MS (APCI-Pos) *M*+1 = 436.2.


Compound 355:



Prepared according to the general procedure described for compound 350 using benzoyl chloride.

***N*-{3-Methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzyl}-benzamide** ¹H NMR (400 MHz, *d*₆-DMSO): δ 9.95 (s, 2H), 9.02 (t, *J* = 5.48 Hz, 1H), 8.77 (s, 1H), 8.22 (s, 1H), 8.1 (d, *J* = 8.61 Hz, 1H), 7.9 (d, *J* = 7.83 Hz, 2H), 7.48 (m, 3H), 7.04 (s, 1H), 6.88 (d, *J* = 10.17 Hz, 1H), 4.46 (s, 1H), 4.44 (s, 1H), 3.89 (s, 3H), 2.42 (s, 3H). MS APCI-Pos, *M*+1 = 391.9.


Compound 356:



Prepared according to the general procedure described for compound 350 using pyridine-2-carboxylic acid chloride.

Pyridine-2-carboxylic acid 3-methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzylamide ¹H NMR (400 MHz, *d*₆-DMSO): δ 9.94 (s, 2H), 9.29 (t, *J* = 6.26 Hz, 1H), 8.77 (m, 1H), 8.65 (m, 1H), 8.21 (s, 1H), 8.05 (m, 3H), 7.61 (m, 1H), 7.07 (s, 1H), 6.89 (d, *J* = 8.61 Hz, 1H), 4.47 (s, 1H), 4.45 (s, 1H), 3.88 (s, 1H), 2.42 (s, 3H). MS APCI-Pos no detectable molecular ion.


Compound 357:



Prepared according to the general procedure described for compound 350 using nicotinoyl chloride.

***N*-{3-Methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzyl}-nicotinamide** ¹H NMR (400 MHz, *d*₆-DMSO): δ 10 (s, 2H), 9.26 (t, *J* = 5.87 Hz, 1H), 9.11 (s, 1H), 8.82 (s, 1H), 8.77 (d, *J* = 4.7 Hz, 1H), 8.27 (s, 2H), 8.16 (d, *J* = 7.83 Hz, 1H), 7.58 (m, 1H), 7.09 (s, 1H), 6.95 (d, *J* = 8.61 Hz, 1H), 4.53 (s, 1H), 4.51 (s, 1H), 3.95 (s, 3H), 2.47 (s, 3H). MS APCI-Pos, *M*-1 = 397.9.


Compound 358:



Prepared according to the general procedure described for compound 350 using isonicotinoyl chloride.

***N*-[3-Methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzyl]-isonicotinamide** ¹H NMR (400 MHz, *d*₆-DMSO): δ 9.92 (s, 2H), 9.27 (t, *J* = 5.87 Hz, 1H), 8.74 (s, 1H), 8.71 (d, *J* = 6.3 Hz, 2H), 8.18 (s, 1H), 8.08 (d, *J* = 8.6 Hz, 1H), 7.77 (d, *J* = 6.3 Hz, 2H), 7 (s, 1H), 6.86 (d, *J* = 8.6 Hz, 1H), 4.44 (s, 1H), 4.43 (s, 1H), 3.86 (s, 3H), 2.39 (s, 3H). MS APCI-pos no detectable molecular ion.


Compound 359:



Prepared according to the general procedure described for compound 350 using thiophene-2-carbonyl chloride.

Thiophene-2-carboxylic acid 3-methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzylamide ¹H NMR (400 MHz, *d*6-DMSO): δ 9.96 (s, 2H), 9.03 (t, *J* = 6.26 Hz, 1H), 8.78 (s, 1H), 8.22 (s, 1H), 8.11 (d, *J* = 7.83 Hz, 1H), 7.82 (d, *J* = 4.7 Hz, 1H), 7.77 (d, *J* = 3.91 Hz, 1H), 7.16 (d, *J* = 3.91 Hz, 1H), 7.03 (s, 1H), 6.88 (d, *J* = 10.17 Hz, 1H), 4.43 (s, 1H), 4.42 (s, 1H), 3.89 (s, 3H), 2.42 (s, 3H). MS APCI-*pos*, *M*-1 = 397.9.


Compound 360:



Prepared according to the general procedure described for compound 350 using 4-dimethylamino-2-carbonyl chloride.

3-Dimethylamino-N-{3-methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzyl}-benzamide ¹H
NMR (400 MHz, *d*6-DMSO): δ 9.94 (s, 2H), 8.9 (t, *J* = 5.87 Hz, 1H), 8.77 (s, 1H), 8.21 (s, 1H), 8.09 (d, *J* = 8.61 Hz, 1H), 7.26 (t, *J* = 7.83 Hz, 1H), 7.2 (s, 2H), 7.02 (s, 1H), 6.87 (d, *J* = 8.61 Hz, 2H), 4.44 (s, 1H), 4.42 (s, 1H), 3.88 (s, 3H), 2.93 (s, 6H), 2.42 (s, 3H). MS APCI-pos, *M*+1 = 434.9.

Compound 361:




Prepared according to the general procedure described for compound 350 using 1-phenyl-4-trifluoromethyl-1H-pyrazole-3-carboxyl chloride.

1-Phenyl-4-(3-methoxy-4-((5-methylpyrazin-2-yl)ureido)benzyl)-1H-pyrazole-3-carboxylic acid 3-methoxy-4-[3-(5-methylpyrazin-2-yl)ureido]benzylamide ^1H NMR (400 MHz, d_6 -DMSO): δ 9.96 (s, 2H), 9.09 (t, J = 6.26 Hz, 1H), 8.78 (s, 1H), 8.22 (s, 1H), 8.19 (s, 1H), 8.12 (d, J = 7.83 Hz, 1H), 7.59 (m, 6H), 7.02 (s, 1H), 6.89 (d, J = 8.61 Hz, 1H), 4.43 (s, 1H), 4.42 (s, 1H), 3.9 (s, 3H), 2.42 (s, 3H). MS TIC-pos $M+1$ = 526.2.

Sulfonylated Derivatives


Compound 362:



A solution of 1-(4-aminomethyl-2-methoxy-phenyl)-3-(5-methylpyrazin-2-yl)-urea (0.25 mmol, 1.0 eq.), 4-dimethylaminopyridine (5 mg), diisopropylethylamine (36 mg, 0.275 mmol, 1.1 eq.) in THF (5 mL) was prepared and thiophene-2-sulfonyl chloride (0.275 mmol, 1.1 eq.) was added. The mixture was stirred at room temperature for 24 h. The reaction mixture was partitioned between EtOAc (75 mL) and sat. NaHCO_3 . After separation of the layers, the organic layer was washed with water, saturated brine, dried over MgSO_4 , filtered and concentrated. The residue was purified by flash chromatography eluting with 5 % MeOH in dichloromethane and trituration with ether to give pure products.

Thiophene-2-sulfonic acid 3-methoxy-4-[3-(5-methylpyrazin-2-yl)ureido]benzylamide ^1H NMR (400 MHz, d_6 -DMSO): δ 9.96 (s, 2H), 8.77 (s, 1H), 8.33 (t, J = 7.04 Hz, 1H), 8.31 (s, 1H), 8.07 (d, J = 7.83 Hz, 1H), 7.93 (d, J = 4.7 Hz, 1H), 7.59 (s, 1H), 7.18 (t, J = 3.91 Hz, 1H), 6.9 (s, 1H), 6.8 (t, J = 7.04 Hz, 1H), 4.05 (s, 1H), 4.03 (s, 1H), 3.85 (s, 3H), 2.51 (s, 3H). MS APCI-Pos, $M+1$ = 433.9.


Compound 363:



Prepared according to the general procedure described for compound 362 using benzenesulfonyl chloride.

***N*-{3-Methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzyl}-benzenesulfonamide** ¹H NMR (400 MHz, *d*₆-DMSO): δ 9.95 (s, 2H), 8.77 (s, 1H), 8.21 (s, 1H), 8.13 (t, *J* = 6.26 Hz, 1H), 8.05 (d, *J* = 8.61 Hz, 1H), 7.81 (d, *J* = 7.04 Hz, 2H), 7.58 (m, 3H), 6.85 (s, 1H), 6.77 (d, *J* = 8.61 Hz, 1H), 3.96 (s, 1H), 3.95 (s, 1H), 3.81 (s, 3H), 2.42 (s, 3H). MS APCI-Pos, *M*+1 = 428.2.


Compound 364:



Prepared according to the general procedure described for compound 362 using 2-trifluoromethoxybenzenesulfonyl chloride.

***N*-{3-Methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzyl}-2-trifluoromethoxybenzenesulfonamide** ¹H NMR (400 MHz, *d*₆-DMSO): δ 9.94 (s, 2H), 8.77 (s, 1H), 8.38 (s, 1H), 8.21 (s, 1H), 8.02 (d, *J* = 8.61 Hz, 1H), 7.89 (m, 1H), 7.72 (t, *J* = 7.83 Hz, 1H), 7.51 (d, *J* = 7.83 Hz, 2H), 6.88 (s, 1H), 6.75 (d, *J* = 10.17 Hz, 1H), 4.11 (s, 2H), 3.82 (s, 3H), 2.42 (s, 3H). MS APCI-Pos, *M*+1 = 512.0.


Compound 365:



Prepared according to the general procedure described for compound 362 using 3-methoxybenzenesulfonyl chloride.

3-Methoxy-*N*-{3-methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzyl}-benzenesulfonamide ¹H NMR (400 MHz, *d*₆-DMSO): δ 9.92 (s, 2H), 8.73 (s, 1H), 8.17 (s, 1H), 8.08 (s, 1H), 8.01 (d, *J* = 7.83 Hz, 1H), 7.45 (t, *J* = 7.83 Hz, 1H), 7.34 (d, *J* = 7.83 Hz, 1H), 7.24 (s, 1H), 7.14 (m, 1H), 6.81 (s, 1H), 6.74 (d, *J* = 7.83 Hz, 1H), 3.93 (s, 2H), 3.77 (s, 3H), 3.76 (s, 3H), 2.38 (s, 3H). MS APCI-Pos, *M*+1 = 458.0.


Compound 366:



Prepared according to the general procedure described for compound 362 using 4-methoxybenzenesulfonyl chloride.

4-Methoxy-N-{3-methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzyl}-benzenesulfonamide ^1H NMR (400 MHz, d_6 -DMSO): δ 9.93 (s, 2H), 8.74 (s, 1H), 8.18 (s, 1H), 8.02 (d, $J = 8.61$ Hz, 1H), 7.92 (s, 1H), 7.69 (d, $J = 8.61$ Hz, 2H), 7.05 (d, $J = 9.39$ Hz, 2H), 6.81 (s, 1H), 6.74 (d, $J = 7.83$ Hz, 1H), 3.88 (s, 2H), 3.79 (s, 6H), 2.39 (s, 3H). MS APCI-Pos, $M+1 = 458.2$.

Compound 367:



Prepared according to the general procedure described for compound 362 using pyridine-2-sulfonyl chloride.

Pyridine-2-sulfonic acid 3-methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzylamide ^1H NMR (400 MHz, d_6 -DMSO): δ 9.91 (s, 2H), 8.74 (s, 1H), 8.68 (d, $J = 4.7$ Hz, 1H), 8.17 (s, 1H), 7.99 (m, 2H), 7.86 (d, $J = 7.83$ Hz, 1H), 7.6 (s, 1H), 6.86 (s, 1H), 6.74 (d, $J = 8.61$ Hz, 1H), 4.1 (s, 2H), 3.80 (s, 3H), 3.32 (s, 1H), 2.38 (s, 3H). MS APCI-Pos, $M+1 = 428.9$

Additional preferred compounds of the present invention include

N-(2-dimethylamino-1-phenyl-ethyl)-3-methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzamine;

5 N-(1-aza-bicyclo[2.2.2]oct-3-yl)-3-methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzamide;

N-(3-R-1-cyclohexylmethyl-pyrrolidin-3-yl)-3-methoxy-4-[3-(5-methyl-pyrazin-2-yl)-ureido]-benzamide;

10 1-[2-(2-dimethylamino-ethoxy)-5-methyl-phenyl]-3-pyrazin-2-yl-urea;

1-[2-(3-dimethylamino-propoxy)-5-methyl-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea;

1-(5-methyl-pyrazin-2-yl)-3-[5-methyl-2-(pyridin-3-ylmethoxy)-phenyl]-urea;

15 1-[2-(2-dimethylamino-1-dimethylaminomethyl-ethoxy)-5-methyl-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea;

1-[5-methyl-2-(2-S-1-methyl-pyrrolidin-2-ylmethoxy)-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea;

20 1-[5-methyl-2-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea;

1-[5-methyl-2-(1-methyl-piperidin-4-yloxy)-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea;

1-[5-methyl-2-(3-(S)-1-methyl-piperidin-3-ylmethoxy)-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea;

25 1-[5-methyl-2-(3-(R)-1-methyl-piperidin-3-ylmethoxy)-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea;

1-[5-methyl-2-(1-methyl-piperidin-2-ylmethoxy)-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea;

30 1-[5-methyl-2-(1-methyl-piperidin-3-yloxy)-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea;

1-[5-methyl-2-(1-methyl-piperidin-3-ylmethoxy)-phenyl]-3-quinoxalin-2-yl-urea;

1-[5-methyl-2-(piperidin-3-ylmethoxy)-phenyl]-3-(5-methyl-pyrazin-2-yl)-urea;

- 203 -

1-[5-fluoro-2-(1-methyl-piperidin-3-ylmethoxy)-
phenyl]-3-(5-methyl-pyrazin-2-yl)-urea;
1-[5-fluoro-2-(1-methyl-piperidin-4-yloxy)-phenyl]-
3-(5-methyl-pyrazin-2-yl)-urea;
5 1-[4-fluoro-2-(1-methyl-piperidin-4-yloxy)-phenyl]-
3-(5-methyl-pyrazin-2-yl)-urea;
1-(2-methoxy-4-methylaminomethyl-phenyl)-3-(5-
methyl-pyrazin-2-yl)-urea;
1-(4-{[(furan-3-ylmethyl)-amino]-methyl}-2-methoxy-
10 phenyl)-3-(5-methyl-pyrazin-2-yl)-urea; and
1-{2-methoxy-4-[(4-methoxy-benzylamino)-methyl]-
phenyl}-3-(5-methyl-pyrazin-2-yl)-urea.

Example 15

Identification of Chk1 Inhibitors

15 The human Chk1 cDNA was identified and
cloned as described previously in International
Application No. PCT/US98/18558, filed September 4,
20 1998. A FLAG® tag was inserted in frame with the
amino terminus of the full-length Chk1. The 5'
primer contains an EcoRI site, a Kozak sequence, and
also encodes a FLAG® tag for affinity purification
using the M2 Antibody (Sigma, Saint Louis, IL). The
25 3' primer contains a SalI site. The PCR-amplified
fragment was cloned into pCI-Neo as an EcoRI-SalI
fragment (Invitrogen, Carlsbad, CA), then subcloned
as an EcoRI-NotI fragment into pFastBacI (Gibco-BRL,
Bethesda, MD). Recombinant baculovirus was prepared
30 as described in the Gibco-BRL Bac-to-Bac manual and
used to infect Sf-9 cells grown in CCM3 medium
(HyClone Laboratories, Logan, UT) for expression of
FLAG®-tagged Chk1 protein.

35 FLAG®-tagged Chk1 was purified from frozen
pellets of baculovirus-infected SF9 cells. Frozen

The protein kinase was used to develop an assay for Chk1 kinase activity that includes 100 ng purified FLAG®-Chk1 (150 pmol of ATP/min), 20 μ M Cdc25C peptide (H-leu-tyr-arg-ser-pro-ser-met-pro-glu-asn-leu-asn-arg-arg-arg-arg-OH) (SEQ ID NO: 1), 400 μ M ATP, 2 μ Ci [³²P]-gATP, 20 mM Hepes pH 7.2, 5 mM MgCl₂, 0.1% NP40 and 1 mM DTT. This assay was used to screen approximately 100,000 small molecule

- 205 -

inhibitors. Reactions were initiated by the addition of ATP-containing reaction mix and carried out at room temperature for 10 min. Reactions were stopped by the addition of phosphoric acid (150 mM final concentration) and transferred to phosphocellulose discs. The phosphocellulose discs were washed five times with 150 mM phosphoric acid and air-dried. Scintillation fluid was added and discs were counted in a Wallac scintillation counter. The screen identified a number of Chk1 inhibitors having IC₅₀ values in the range of 1 to 100 μ M.

Example 16

Chk1 Kinase Inhibitors are Selective

Chk1 inhibitors of the invention were tested for selectivity as against one or more other protein kinases, i.e., DNA-PK, Cdc2, Casein Kinase I (CKI), Chk2, p38 MAP kinase, Protein Kinase A (PKA), and calcium-calmodulin protein kinase II (CaM KII). Assay procedures for all of these kinases except Chk2 have been previously described in the literature, including U.S. provisional patent application 60/229,899, filed September 1, 2000, and U.S. patent application 08/184,605, filed January 21, 1994, both of which are herein incorporated by reference. Activity of the compounds against Chk2 was assayed as follows: 128 ng of purified His-tagged Chk2 was incubated with up to 100 mM Chk1 inhibitor in the presence of 4 mM ATP, 1 mCi [32P]g-ATP, 20 mM Hepes pH 7.5, 5 mM MgCl₂, and 0.25% NP40 for 20 minutes at room temperature. Reactions were stopped with a final concentration of 150 mM phosphoric acid, and 5/8 of the reaction mixture was transferred to phosphocellulose discs. The discs were washed five

- 206 -

times with 150 mM phosphoric acid, and air-dried. Scintillant was added and radioactivity was counted using a Wallac beta counter. p38 MAP kinase, PKA, CaM KII, and Cdc2 were purchased from New England Biolabs, and assays were performed according to the manufacturer's instructions using 4-50 μ M ATP and testing Chk1 inhibitor concentrations as high as 100 μ M. All inhibitors tested were showed at least 5-fold selectivity for Chk1 over the other enzymes.

Example 17

Chk1 Inhibitors Block Chk1 Function in Cells

Chk 1 is activated in response to ionizing radiation and certain chemical DNA damaging agents. In the presence of DNA damage, Chk1 is activated and causes a cell cycle arrest. In mammalian cells, the best-characterized cell cycle arrest invoked by Chk1 is a G2 arrest. Activation of Chk 1 by DNA damage results in the phosphorylation and inactivation of Cdc25C, the dual specificity phosphatase that normally dephosphorylates cyclin B/cdc2 as cells progress into mitosis (Funari et al., *Science*, Sep. 5, 1997; 277(5331)1495-7; Sanchez et al.; Matsuoka et al.; and Blasina et al.). This negative regulation of Cdc2 activity causes cell cycle arrest in order to prevent cells from entering mitosis in the presence of DNA damage or unreplicated DNA. Inhibition of Chk1, therefore, allows cells to progress through the cell cycle in the presence of DNA damage of unreplicated DNA.

To establish that the Chk1 inhibitors prevented Chk1 function in cells, inhibitors were tested in molecular cell-based assays. Since mammalian Chk1 has been shown to phosphorylate

- 207 -

Cdc25C *in vitro*, suggesting that it negatively regulates cyclin B/cdc2 in response to DNA damage, the ability of the Chk1 inhibitors to enhance the activity of CyclinB/cdc2 was analyzed. The experiment was designed as follows: HeLa cells were irradiated with 800 rads and incubated for 7 hours at 37°C. Because these cells are functionally p53 negative, they arrest exclusively in G2. Then, nocodazole was added to a concentration of 0.5 µg/mL and incubated for 15 hours at 37°C. The addition of nocodazole was designed to trap any cells that progressed through the G2 arrest into M. Finally, a Chk1 inhibitor was added for 8 hours, the cells harvested, lysed and immunoprecipitated equal amounts of protein with an antibody to Cyclin B1 (New England Biolabs) as suggested by the manufacturer. IPs then were analyzed for CyclinB-associated cdc2 kinase activity by assaying histone H1 kinase activity (Yu et al., J Biol Chem. Dec. 11, 1998;273(50):33455-64). The results demonstrated that Compound 29 overrides the IR-induced inactivation of Cyclin B/Cdc2.

In addition, whether the Chk1 inhibitors abrogate the IR-induced G2 DNA damage checkpoint as assayed by mitotic index experiments was tested. HeLa cells (approximately 1×10^6) were treated as described above. Cells were harvested by centrifugation, washed once with PBS, then resuspended in 2.5 mL 75 mM KCl and centrifuged again. The cells then were fixed in 3 mL of freshly prepared cold, acetic acid: methanol (1:3) and incubated on ice for 20 minutes. Cells were pelleted, fix solution aspirated and resuspended in 0.5 mL of PBS. Mitotic spreads were prepared by pipeting 100 µL of the fixed cells onto a glass microscope slide and flood-

Abrogation of the IR-induced G2 checkpoint allows the cells to continue through the cell cycle, presumably in the presence of DNA damage, as demonstrated by analysis of DNA content by FACS profile. 293T cells were treated with 800 rads of ionizing radiation and increasing concentrations (up to 80 mM) of some of the Chk1 inhibitors. The cells then were harvested and fixed with 5 mL of cold 70% ethanol at -20°C overnight. The cells then were pelleted by centrifugation at 1000 x g for 10 minutes, and stained with 1 mL of solution containing 50 mg/mL propidium iodide and 250 mg/mL RNase for 30 minutes at room temperature. Stained cells were then analyzed by FACS on FL2 using a Becton-Dickinson apparatus. These experiments demonstrated that, while the cells treated with radiation and vehicle alone remained arrested in G2, the Chk1 inhibitor treated cells were distributed in G1 and S phase. These data, taken together with the data above, suggest that the Chk1 inhibitors allow cells to continue cycling in the presence of ionizing radiation.

Example 18

**Chk1 Inhibitors enhance killing
of cells by cancer treatments**

To test the hypothesis that inhibition of Chk1 potentiates the killing effect of DNA-damaging agents, cells were incubated in the presence of selective Chk1 inhibitors and either irradiation or chemical DNA-damaging agents. Cells plated at a density of 1000-2000 per well in 96-well microtitre plates were grown in RMPI 1640 containing 10% FBS, 100 U/mL penicillin and 100 µg/mL streptomycin for 18 hours at 37°C in a humidified incubator with 5% CO₂. Cells tested included HeLa, ACHN, 786-0, HCT116, SW620, HT29, Colo205, SK-MEL-5, SK-MEL-28, A549, H322, OVCAR-3, SK-OV-3, MDA-MB-231, MCF-7, PC-3, HL-60, K562, and MOLT4. All cell line designations refer to human cell lines and refer to the following:

HeLa	cervical adenocarcinoma
ACHN	renal adenocarcinoma
786-0	renal adenocarcinoma
HCT116	colon carcinoma
SW620	colon carcinoma, lymph node metastasis
HT-29	colonrectal adenocarcinoma
Colo205	colon adenocarcinoma
SK-MEL-5	melanoma
SK-MEL-28	malignant melanoma
A549	lung carcinoma
H322	broncholoalveolar carcinoma
OVCAR-3	ovarian adenocarcinoma
SK-OV-3	ovarian adenocarcinoma
MDA-MB-231	breast adenocarcinoma
MCF-7	breast adenocarcinoma

- 210 -

PC-3	prostate adenocarcinoma, from metastasis to bone
HL-60	acute promyelocytic leukemia
K562	chronic myelogenous leukemia
MOLT4	acute lymphoblastic leukemia; T lymphoblast

5

Cells were treated with media containing chemotherapeutic drugs alone or chemotherapeutic drugs and Compounds 12 and 29. Cells were incubated for approximately 5 days before growth was measured by determination of levels of ³H-thymidine uptake. Chemotherapeutic drugs included etoposide, doxorubicin, cisplatin, chlorambucil, 5-fluorouracil (5-FU). The drug concentration necessary to inhibit cell growth to 90% of untreated control cells was defined as the GL₉₀. At concentrations less than 100 μM, Compounds 12 and 29 enhanced the killing of 5-FU from 2- to 10-fold.

10

15

20

Compounds 2 and 12 were tested with additional antimetabolites, including methotrexate, hydroxyurea, 2-chloroadenosine, fludarabine, azacytidine, and gemcitabine for an ability to enhance killing of the agents. These Chk1 inhibitors were found to enhance the killing of cells to hydroxyurea, fludarabine, 5-azacytidine, and methotrexate suggesting that the combination of inhibition of Chk1 and blocking of DNA synthesis leads to increased cell death by these agents.

25

In addition, the ability of the Chk1 inhibitor to enhance killing by irradiation was tested. In HeLa cells, Compounds 12 and 29 were found to enhance killing by irradiation 2-3 fold.

30

Animal Tumor Models

Mice were randomized (5 mice/group) into four treatment groups and used when tumors reached a weight of 75-100 mg (usually 7-11 days post-inoculation). Tumors were measured with vernier calipers and tumor weights were estimated using the empirically derived formula: tumor weight (mg) = tumor length (mm) x tumor width (mm)²/3.3. Treatment consisted of i) 100 μ L intraperitoneal (i.p). injection of 5-FU at 50 mg/kg, 100 mg/kg, or 150 mg/kg. A dose-dependent delay in tumor growth was observed in the mice treated with 5-FU. Tumor size was monitored every other day for the duration of the experiment.

Obviously, many modifications and variations of the invention as hereinbefore set forth can

- 212 -

be made without departing from the spirit and scope thereof, and, therefore, only such limitations should be imposed as are indicated by the appended claims.